arXiv:1008.3587v2 [math.SP] 7 Dec 2010

$2D$- and -$3D$Magnetic Schrödinger Operators: Short Loops, Pointwise Spectral Asymptotics and Asymptotics of Dirac Energy

Victor Ivrii

December 8, 2010

Abstract

We consider 2- and 3-dimensional Schrödinger or generalized Schrödinger-Pauli operators with the non-degenerating magnetic field in the open domain under certain non-degeneracy assumptions we derive pointwise spectral asymptotics.

We also consider asymptotics of some related expressions (see below). For all asymptotics loops rather than periodic trajectories play important role.

0 Introduction

In this paper we consider 2D- and 3D-magnetic Schrödinger operator (13.1.1) satisfying assumptions (13.1.2)–(13.1.5) and consider pointwise asymptotics of $e(x,x,0)$ and also asymptotics of expression (6.3.4):

$$\mathsf{I} := \iint e(x,y,0)e(y,x,0)\omega(x,y)\psi_2(x)\psi_1(y)\,dx\,dy \tag{0.1}$$

as well as of

$$\mathsf{J} := \iint e(x,x,0)e(y,y,0)\omega(x,y)\psi_2(x)\psi_1(y)\,dxdy \tag{0.2}$$

with a function $\omega(x,y)$ satisfying assumption (6.3.5):

(0.3) $\omega(x,y) := \Omega(x,y;x-y)$ where function Ω is smooth in $B(0,1)\times B(0,1)\times B(\mathbb{R}^2\setminus 0)$ and homogeneous of degree $-\kappa$ $(0<\kappa<2)$ with respect to its third argument,

and with smooth cut-off functions ψ_1,ψ_2.

These two expressions play a role in the applications to the multiparticle quantum theory of Part VIII. Actually, instead of asymptotics of (0.2) we consider related estimates of

$$\mathsf{K} := \iint \bigl(e(x,x,\tau)-h^{-d}\mathcal{N}_x(\tau)\bigr)\bigl(e(y,y,\tau)-h^{-d}\mathcal{N}_x(\tau)\bigr)\omega(x,y)\,dxdy \tag{0.4}$$

where $\mathcal{N}_x(\tau)$ denotes some approximation to $e(x,x,\tau)$.

We assume that V/F satisfies non-degeneracy conditions (13.3.45) and (13.3.54) i.e.

$$V\le -\epsilon_0 \qquad \text{in}\quad B(0,1), \tag{0.5}$$

$$|F|\ge \epsilon \qquad \text{in}\quad B(0,1), \tag{0.6}$$

$$|\nabla \frac{V}{F}|\asymp 1 \qquad \text{in}\quad B(0,1); \tag{0.7}$$

in asymptotics of expressions (0.1)–(0.2) we will be able to drop (0.5) and replace (0.7) by

$$|\nabla \frac{V}{F}|\le \epsilon_0 \implies |\operatorname{det}\operatorname{Hess}\frac{V}{F}|\ge \epsilon_0 \tag{0.8}$$

or

$$|\nabla \frac{V}{F}|\le \epsilon_0 \implies \operatorname{det}\operatorname{Hess}\frac{V}{F}\ge \epsilon_0. \tag*{$(0.8)^+$}$$

As we consider Schrödinger-Pauli operator (13.5.3) and $\mu h\gtrsim 1$ conditions (0.7)–$(0.8)^+$ will be modified to

$$|\frac{V}{F}+(2m+1-\varsigma)|+|\nabla\frac{V}{F}|\asymp 1 \qquad \text{in}\quad B(0,1)\ \forall m\in\mathbb{Z}^+; \tag*{$(0.7)'$}$$

$$|\frac{V}{F}+(2m+1-\varsigma)|+|\nabla\frac{V}{F}|\le \epsilon_0 \implies |\operatorname{det}\operatorname{Hess}\frac{V}{F}|\ge\epsilon_0, \tag*{$(0.8)'$}$$

$$|\frac{V}{F}+(2m+1-\varsigma)|+|\nabla\frac{V}{F}|\le \epsilon_0 \implies \operatorname{det}\operatorname{Hess}\frac{V}{F}\ge\epsilon_0 \tag*{$(0.8)^{+\prime}$}$$


(a) prolate cycloid (b) perturbed prolate cycloid

Figure 1: Classical trajectories

respectively.

Non-degeneracy assumptions (0.7)–(0.8)$^+$ eliminate periodicity at least as $\mu h \ll 1$ but there are plenty of loops as on figures 1a and 1b and we know that loops play the same important role in the pointwise asymptotics (of $e(x,x,0)$) as periodic trajectories play in the local asymptotics (of $\int e(x,x,0)\psi(x)\,dx$). There also will be many paths from x to y and back and these paths play important role in the asymptotics of expression (0.1).

Fortunately, even if there are plenty of looping points they are rather exceptional on the classical trajectory. Surely, one needs to remember that if we study $e(x,x,0)$ point x is fixed but direction ξ varies and thus trajectory[1] moves so it passes through x and remains on the energy level 0 (see figures 2a–2b below). However for the geometrical simplicity we will sometimes talk about x moving along trajectory. As we study expression (0.1) x, y both move.

In fact, situation is more complicated than this due to the uncertainty principle: as μ increases not cyclotrons becomes smaller and tighter but the lines retain their width and become wider in comparison with the cyclotron, and symbolically we can consider figures 3a–3c below.

We also generalize section 6.3 and we consider expression (0.1). It follows from section 6.3 that $\mathsf{I} = \mathcal{I}^{\mathsf{W}} + O(h^{1-d-\kappa})$ as $\mu = O(1)$ with $\mathcal{I}^{\mathsf{W}}$ defined by the same formula but with $e(x,y,\tau)$ replaced by

$$e^{\mathsf{W}}_y(x,y,\tau) := (2\pi h)^{-d} \int_{g(y,\xi)\le V(y)+\tau} e^{ih^{-1}\langle x-y,\xi\rangle}\,d\xi. \tag{0.9}$$

---

[1] Even if one calls them trajectories they are only *x-projections* of actual trajectories which live in (x,ξ)-space.

Then the standard rescaling technique implies the same asymptotics but with the remainder estimate $O(\mu h^{1-d-\kappa})$ provided $1\le \mu = o(h^{-1})$.

Let $d=2$. Then *in the general case* it is the best remainder estimate possible while $O(\mu h^{-1})$ is the best possible remainder estimate for

$$\int e(x,x,\tau)\psi(x)\,dx; \tag{0.10}$$

recall that this estimate cannot be improved in the case of constant magnetic field F and $g^{jk}=\mathsf{const}$, $V=\mathsf{const}$.

On the other hand, *in the generic case* the remainder estimate for (0.10) is $o(\mu^{-1}h^{-1})$ and the principal part is $\int e^{\mathsf{MW}}(x,\tau)\psi(x)\,dx$ if F does not vanish and $\mu\le h^{-1}$.

Meanwhile for $d=3$ the remainder estimate for I' is $O\bigl(h^{-2}+\mu h^{-1-\delta}\bigr)$ and under very mild non-degeneracy assumptions it is $O\bigl(h^{-2}+\mu h^{-1}\bigr)$.

Therefore our purpose is to get the sharper remainder estimate for (0.1) under the same non-degeneracy assumptions. This is a very daunting task since for (0.10) *periodic trajectories* were the main source of trouble and they were broken in the generic case; for (0.1) *loops* are also the source of trouble, and in the generic case former periodic trajectories generate a lot of loops.

It is rather easy to derive asymptotics with the sharp remainder estimate $O(\mu^{-1}h^{-1-\kappa})$ as $d=2$ and $O(h^{-2})$ as $d=3$ but with the principal part given by very implicit Tauberian formula when I^{T} is defined by (0.1) but with $e(x,y,\tau)$ replaced by its Tauberian approximation

$$e^{\mathsf{T}}(x,y,\tau):=h^{-1}\int_{-\infty}^{\tau} F_{t\to h^{-1}\tau}\bigl(\bar\chi_T(t)u(x,y,t)\bigr)\,d\tau; \tag{0.11}$$

such remainder estimate is a rather easy corollary of the results of Chapter 13. However deriving of an asymptotics with the sharp remainder estimate and rather explicit principal part is much more difficult.

Plan of the paper

In sections 1 and 5 we consider respectively 2D- and 3D-pilot-models with the constant g^{jk} and F and linear V and derive representations of solutions and pointwise asymptotics as magnetic field is weak enough. We also derive asymptotics of expression (0.10) with $\psi(x)$ replaced by $\psi_\gamma(x)$ which is $\psi(x)$ scaled with a spatial scaling parameter γ; these asymptotics are instrumental in the asymptotics of (0.2) and estimates of (0.4).

In sections 2 and 6 we prove the same asymptotics respectively for the general 2D- and 3D-operators; as magnetic field is strong we use explicit expressions for the pilot-model operators as approximations.

In sections 3 and 7 we prove important preliminary results and derive sharp asymptotics for expression (0.1) but with the Tauberian principal part.

In sections 4 and 8 we pass from the Tauberian approximations to Weyl, magnetic Weyl or pilot-model approximations.

Finally, in section 9 we consider estimates of (0.4) with different approximations, leaving to the reader very similar asymptotics of expression (0.2).

The results of this paper are not always sharp or very explicit, but could be made either sharp or completely explicit.

1 Pointwise asymptotics: 2D-pilot-model

1.1 Pilot-model in $\mathbb{R}^2$: propagator

Consider the *pilot-model operator*

$$A=\bar{A}:=h^2D_1^2+(hD_2-\mu x_1)^2+2\alpha x_1. \tag{1.1}$$

We are interested in the Schwartz kernel $U(x,y,t)$ of the propagator $e^{ih^{-1}tA}$. Making h-Fourier transform with respect to $x_2\mapsto \xi_2$ and rescaling $x_1\mapsto \mu x_1$, $t\mapsto \mu t$ we arrive to

$$U(x,y,t)=(2\pi h)^{-1}\mu\int \mathsf{u}(x_1,y_1;\eta,t)e^{ih^{-1}(x_2-y_2)\eta}\,d\eta \tag{1.2}$$

with $\mathsf{u}(x_1,y_1;\eta,t)$ the Schwartz kernel of $e^{i\hbar^{-1}t\mathbf{a}}$ with 1D-Harmonic oscillator

$$\begin{multline}\mathbf{a}=\hbar^2D_1^2+(x_1-\eta)^2+2\alpha\mu^{-1}x_1=\\ \underbrace{\hbar^2D_1^2+(x_1-\eta+\alpha\mu^{-1})^2}_{\bar{\mathbf{a}}}+\mu^{-1}\alpha\underbrace{(2\eta-\alpha\mu^{-1})}_{\zeta(\eta)}.\end{multline} \tag{1.3}$$

Recall that for the Harmonic oscillator $\mathbf{b}=D^2+x^2$ the Schwartz kernel of $e^{it\mathbf{b}}$ [2] is

$$(2\pi)^{-\frac{1}{2}}e^{\frac{i\pi}{4}\sigma(t)}|\sin(2t)|^{-\frac{1}{2}}\exp\Bigl(-\frac{i}{2}\bigl(\cot(2t)(x^2+y^2)-2xy\csc(2t)\bigr)\Bigr) \tag{1.4}$$

where $\sigma(t)=1,2,3,4$ at $(0,\frac{\pi}{2}),(\frac{\pi}{2},\pi),(\pi,\frac{3\pi}{2}),(\frac{3\pi}{2},2\pi)$ respectively and it is 2π-periodic and therefore for Harmonic oscillator $\mathbf{b}'=\hbar^2D^2+x^2$ the Schwartz kernel of $e^{it\hbar^{-1}\mathbf{b}'}$ is

$$\begin{multline}(2\pi\hbar)^{-\frac{1}{2}}e^{\frac{i\pi}{4}\sigma(t)}|\sin(2t)|^{-\frac{1}{2}}\times\\ \exp\Bigl(-\frac{i}{2}\hbar^{-1}\bigl(\cot(2t)(x^2+y^2)-2xy\csc(2t)\bigr)\Bigr);\end{multline} \tag{1.5}$$

one can prove it easily by rescaling $x\mapsto x\hbar^{\frac{1}{2}}$.

For operator (1.3) the Schwartz kernel of $e^{it\hbar^{-1}\mathbf{a}}$ is obtained by plugging $x:=x_1-\eta+\alpha\mu^{-1}$, $y:=y_1-\eta+\alpha\mu^{-1}$ and multiplication by $\exp\bigl(i\hbar^{-1}\mu^{-1}\alpha\zeta(\eta)\bigr)$. Therefore, multiplying (1.5) by

$$(2\pi\hbar)^{-1}\mu^2\exp\Bigl(i\hbar^{-1}\bigl(\mu^{-1}t\alpha(2\eta-\alpha\mu^{-1})+(x_2-y_2)\eta\bigr)\Bigr) \tag{1.6}$$

and integrating by η we arrive to[3]

$$U(x,y,t)=(2\pi\hbar)^{-\frac{3}{2}}\mu^2\int e^{\frac{i\pi}{4}\sigma(t)}|\sin(2t)|^{-\frac{1}{2}}e^{i\hbar^{-1}\bar{\varphi}(x,y,\eta,t)}\,d\eta \tag{1.7}$$

with

$$\begin{multline}\bar{\varphi}:=-\frac{1}{2}\cot(2t)\bigl((x_1-\eta+\alpha\mu^{-1})^2+(y_1-\eta+\alpha\mu^{-1})^2\bigr)+\\ \csc(2t)(x_1-\eta+\alpha\mu^{-1})(y_1-\eta+\alpha\mu^{-1})+(x_2-y_2)\eta+\\ t\mu^{-1}\alpha(2\eta-\alpha\mu^{-1}).\end{multline} \tag{1.8}$$

We can rewrite (1.7)–(1.8) after integration by η as

$$U(x,y,t)=i(4\pi\hbar)^{-1}\mu^2\csc(t)\,e^{i\hbar^{-1}\bar{\phi}(x,y,t)} \tag{1.9}$$

---

[2] This is a metaplectic operator, see 1.2.3 or H. ter Morsche and P. J. Oonincx [MO].

[3] After rescaling $x\mapsto\mu x$, $y\mapsto\mu y$, $t\mapsto\mu t$, $\mu\mapsto 1$, $h\mapsto\hbar=\mu h$; however we treat $U(x,y,t)$ and $e(x,y,\tau)$ as functions, not as densities with respect to y and it leads to the factor μ^2.

with

$$\begin{aligned}(1.10)\quad \bar{\phi} := -\frac{1}{4}\cot(t)(x_1-y_1)^2\\ +\frac{1}{2}(x_1+y_1+2\alpha\mu^{-1})(x_2-y_2+2t\alpha\mu^{-1})-\frac{1}{4}\cot(t)(x_2-y_2+2t\alpha\mu^{-1})^2-t\alpha^2\mu^{-2};\end{aligned}$$

here the critical point with respect to η is

$$\eta = \frac{1}{2}(x_1+y_1+2\alpha\mu^{-1}) - \frac{1}{2}\cot(t)(x_2-y_2+2t\alpha\mu^{-1}). \tag{1.11}$$

So, we have proven

Proposition 1.1.1. *For the pilot-model operator* (1.1) ${}^{3)}$ *in* $\mathbb{R}^2$ *the Schwartz kernel* $\bar{U}(x,y,t)$ *of* $e^{i\hbar^{-1}t\bar{A}}$ *is given by* (1.9)–(1.11) *and the Schwartz kernel* $e(x,y,\tau)$ *of the spectral projector is given by*

$$\begin{aligned}(1.12)\quad \partial_\tau e(x,y,\tau) = (2\pi\hbar)^{-1}F_{t\to\hbar^{-1}\tau}U(x,y,.) =\\ (2\pi\hbar)^{-1}\int e^{-i\hbar^{-1}t\tau'}U(x,y,t)\,dt\end{aligned}$$

and

$$\begin{aligned}(1.13)\quad e(x,y,\tau) = (2\pi\hbar)^{-1}\int^{\tau}\Bigl(\int e^{-i\hbar^{-1}t\tau'}U(x,y,t)\,dt\Bigr)\,d\tau =\\ (2\pi)^{-1}\int (-it)^{-1}e^{-i\hbar^{-1}t\tau}U(x,y,t)\,dt\end{aligned}$$

with the last integral taken in the sense of the essential value at 0*.*

In the case of the strong magnetic field we will need an alternative representation. Starting from formula for 1D-harmonic oscillator

$$\mathsf{e}(x_1,y_1,\tau) = \sum_{m\in\mathbb{Z}^+}\upsilon_m(x_1)\upsilon_m(y_1)\theta\bigl(\tau-(2m+1)\bigr) \tag{1.14}$$

with Hermite functions υ_m we after the same rescaling and transition to the pilot-model as before arrive to

Proposition 1.1.2. *For the pilot-model operator* (1.1) $^{3)}$ *in* $\mathbb{R}^2$ *the Schwartz kernel of the spectral projector is defined by*

$$\begin{multline}
e(x,y,\tau) = \\
(2\pi)^{-1}\mu^2\hbar^{-1}\sum_{m\in\mathbb{Z}^+}\int \upsilon_m\bigl(\eta+\hbar^{-\frac{1}{2}}(x_1-y_1)\bigr)\upsilon_m\bigl(\eta-\hbar^{-\frac{1}{2}}(x_1-y_1)\bigr)\times \\
\theta\Bigl(\tau-\alpha\mu^{-1}(x_1+y_1)-2\alpha\mu^{-1}\hbar^{\frac{1}{2}}\eta-\alpha^2\mu^{-2}-(2m+1)\hbar\Bigr)e^{i\hbar^{-\frac{1}{2}}(x_2-y_2)\eta}\,d\eta
\end{multline}\tag{1.15}$$

and thus

$$\begin{multline}
\partial_\tau e(x,y,\tau) = \\
(4\pi\alpha)^{-1}\mu^3\hbar^{-\frac{3}{2}}\sum_{m\in\mathbb{Z}^+}\int \upsilon_m\bigl(\eta+\hbar^{-\frac{1}{2}}(x_1-y_1)\bigr)\upsilon_m\bigl(\eta-\hbar^{-\frac{1}{2}}(x_1-y_1)\bigr)\times \\
e^{i\hbar^{-\frac{1}{2}}(x_2-y_2)\eta}\Bigr|_{\eta=\eta_m}
\end{multline}\tag{1.16}$$

with

$$\eta_m := (2\alpha)^{-1}\mu\hbar^{-\frac{1}{2}}\bigl(\tau-\alpha\mu^{-1}(x_1+y_1)-\alpha^2\mu^{-2}-(2m+1)\hbar\bigr).\tag{1.17}$$

1.2 Tauberian estimate

Consider now pilot-model (1.1) in $B(0,1)\subset X\subset\mathbb{R}^2$. As dynamics (classical or microlocal) starts in $B(0,\epsilon)$ it is confined to $B(0,1)$ for $|t|\le T^*=\epsilon\mu$ and thus in this time interval we can use formulae of the previous subsection (modulo negligible term).

1.2.1 Preparatory estimate

Let us rescale$^{3)}$ and set $x=y=0$ in (1.9)–(1.11):

$$U(0,0,t)\equiv i(4\pi\hbar)^{-1}\mu^2\csc(t)\,e^{i\hbar^{-1}\bar\phi(t)}\tag{1.18}$$

with

$$\bar\phi(t):=t^2\alpha^2\mu^{-2}\cot(t)+\alpha^2\mu^{-2}t;\tag{1.19}$$

$$\eta=-t\alpha\mu^{-1}\cot(t)+\alpha\mu^{-1}.\tag{1.20}$$

We are interested in

$$F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)U(0,0,t) \tag{1.21}$$

with $T=T^*$. Plugging (1.18) we arrive to

$$\begin{gathered} F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)U(0,0,t)\equiv\\ i(4\pi\hbar)^{-1}\mu^2\int\bar{\chi}_T(t)\csc(t)e^{i\hbar^{-1}(\bar{\phi}(t)-\tau)}\,dt=\\ (4\pi\hbar)^{-1}\mu^2\int\bar{\chi}_T(t)\csc(t)\exp\Bigl(i\hbar^{-1}\bigl(-t^2\alpha^2\mu^{-2}\cot(t)+t\mu^{-2}\alpha^2-t\tau\bigr)\Bigr)\,dt. \end{gathered}\tag{1.22}$$

Applying stationary phase to

$$\varphi(t):=-t^2\mu^{-2}\alpha^2\cot(t)+t\mu^{-2}\alpha^2-t\tau \tag{1.23}$$

we get

$$\mu^{-2}\alpha^2(t^2-t\sin(2t))=(\tau-\mu^{-2}\alpha^2)\sin^2(t). \tag{1.24}$$

Remark 1.2.1. Consider corresponding classical trajectory:

$$x_1=\rho\cos(2s)+\eta-\alpha\mu^{-1},\qquad \xi_1=\rho\sin(2s)+\alpha\mu^{-1}s \tag{1.25}$$

and there is a self-intersection iff

$$\left\{\begin{aligned} &\cos(2s_1)=\cos(2s_2),\\ &-\rho\sin(2s_1)+2\alpha\mu^{-1}s_1=-\rho\sin(2s_2)+2\alpha\mu^{-1}s_2; \end{aligned}\right. \tag{1.26}$$

then $s_2+s_1=\pi k$ with $k\in\mathbb{Z}$ (because we cannot fulfill the second equation as $s_1\ne s_2$ and $s_2-s_1=\pi k$); then $2s_1=\pi k-t$, $2s_2=\pi k+t$ and the second equation is $\rho\sin(\pi k-t)=-\alpha\mu^{-1}t$ or, equivalently, $\rho=-\alpha\mu^{-1}t\csc(\pi k-t)$.

Further, we need to satisfy $\rho\cos(\pi k-t)+\eta-\alpha\mu^{-1}=0$ (as $x_1=0$ is a level of intersection) and $(\eta-\alpha\mu^{-1})^2+2\alpha\mu^{-1}(\eta-\alpha\mu^{-1})+\alpha^2\mu^{-2}=\tau$ (to be on the energy level τ) and then $\rho^2\cos^2 t-2\alpha\mu^{-1}\rho\cos(\pi k-t)=\tau-\alpha^2\mu^{-2}$.

Plugging $\rho=-\alpha\mu^{-1}t\csc(\pi k-t)$ we conclude that this equation becomes (1.24). We can find η to satisfy condition $\rho\cos(2s_1)+\eta-\alpha\mu^{-1}=0$ (as $(0,.)$ is a point of intersection, or equivalently, $\eta=-\alpha\mu^{-1}t\cot(t)+\alpha\mu^{-1}$.

We need to justify the stationary phase method. Note first that the spacing between successive stationary points is $\sim \pi$. Therefore

$$t_k = -t_{-k}, \quad t_k \sim \pi k, \quad \sin(t_k) \sim \alpha\mu^{-1}\tau^{-\frac{1}{2}}\pi k \tag{1.27}$$

and

(1.28) The number of stationary points is $\sim 2\pi^{-1}|\alpha|^{-1}\mu\tau^{-\frac{1}{2}}$.

Recall that

$$\varphi'(t) = \alpha^2\mu^{-2}\bigl(t^2\csc^2(t) - 2t\cot(t) + 1\bigr) - \tau \tag{1.29}$$

and

$$\varphi''(t) = -2\alpha^2\mu^{-2}\csc(t)\Bigl(t^2\csc^2(t)\cos(t) - 2t\csc(t) + \cos(t)\Bigr); \tag{1.30}$$

therefore

(1.31) Let $k \ne 0$; then $\varphi''(t_k) \asymp -2\tau\cot(t_k)$ as $|\cos(t_k)| \ge C_0\varepsilon$ and moreover $\varphi''(t_k) \sim -2\tau\cot(t_k)$ as $|\cos(t_k)| \gg \varepsilon$.

Therefore as $|\sin(t_k)| \ge \epsilon$, $|\cos(t_k)| \ge \epsilon$ stationary points are non-degenerate and we can use the stationary phase method.

In the *near-pole zone* $\{|\sin(t)| \le \epsilon\}$ we need to remember about singularity as $\sin(t) = 0$; so we need to introduce a scale $\ell \asymp |\sin(t)|$ and the stationary phase method is expected to work only as $|\varphi''|\ell^2 \ge C_0\hbar$ or equivalently $|\sin(t_k)| \ge C_0\hbar$ or, finally

$$|k| \ge \bar{k} := C_0\varepsilon^{-1}\hbar \tag{1.32}$$

with

$$\varepsilon := \alpha\mu^{-1}(\tau - \alpha^2\mu^{-2})^{-\frac{1}{2}}. \tag{1.33}$$

Summarizing, we arrive to

(1.34) Condition (1.32) is fulfilled for all $k \ne 0$ as $\varepsilon \ge C_0\hbar$; otherwise stationary phase method fails for $|k| \le \bar{k}$. In particular, $\bar{k} = C_0\mu^2 h$ as $|\alpha| \asymp 1$, $\tau \asymp 1$ and condition (1.32) is fulfilled for all $k \ne 0$ as $\mu \le \epsilon_0 h^{-\frac{1}{2}}$.

Meanwhile, there is no singularity in the *near-equator zone* $\{|\cos(t)| \le \epsilon\}$ but there is a degeneration as $\cos(t) = 0$ and therefore we need to introduce a a scale $\ell \asymp |\cos(t)|$ and the stationary phase method is expected to work only as $|\varphi''|\ell^2 \ge C_0\hbar$ or equivalently $|\cos(t)| \ge C_0\max(\varepsilon, \hbar^{\frac{1}{3}})$.

Definition 1.2.2. *(i)* $\{|\sin(t)| \le \max(\epsilon_0\varepsilon, C_0\hbar)\}$ *is a* near-pole singular zone*;*

(ii) $\{|\cos(t)| \le C_0\max(\varepsilon, \hbar^{\frac{1}{3}})\}$ *is a* near-equator singular zone*;*

(iii) $\{|\sin(t)| \le \max(\epsilon_0\varepsilon, C_0\hbar), |\cos(t)| \ge C_0\max(\varepsilon, \hbar^{\frac{1}{3}})\}$ *is a* regular zone*.*

We apply the stationary phase method in the regular zone and justify it and we estimate contributions of the singular zones.

Note that the simple integration by parts brings factors $\hbar/(\varphi'\sin(t))$ and $\hbar\varphi''/(\varphi')^2$ and contribution of zones where these expressions are less than h^δ are negligible. However at this moment we want just to estimate.

Remark 1.2.3. As long as $\mu h \lesssim 1$ we will assume that $\tau \asymp 1$; while we can always achieve $\alpha \asymp 1$ by rescaling $x \mapsto |\alpha|x$, $\mu \mapsto \mu|\alpha|^{-1}$, $\hbar \mapsto \hbar$ we will need many intermediate results and it is simpler to deal with the general case $|\alpha| \le 1$ from the very beginning. Without any loss of the generality one can assume that

$$0 < \alpha \le 1 \qquad (\implies \varepsilon \le \mu^{-1}). \tag{1.35}$$

First, consider zone

$$\bigl\{t:\ |\sin t| \le (1-\epsilon_0)\varepsilon|t|,\ |t| \ge \epsilon_0\bigr\}. \tag{1.36}$$

Then $|\varphi'| \asymp \varepsilon^2 t^2 |\sin(t)|^{-2}$ and $|\varphi''| \lesssim \varepsilon^2 t^2|\sin(t)|^{-3}$ and both factors $|\hbar/\varphi'\sin(t)|$ and $|\hbar\varphi''/(\varphi')^2|$ do not exceed $\hbar|\sin(t)|/\varepsilon^2t^2 \le \hbar/\varepsilon|t|$ and are less than 1 as $\hbar|\sin(t)| \le \varepsilon^2t^2$. This always holds in zone (1.36) if $\hbar \le \varepsilon$.

Then multiple integration by parts brings factor $C(\hbar|\sin(t)|/\varepsilon^2t^2)^l$ with an arbitrarily large exponent l and contribution of k-th "tick"[4] to (1.22) does not exceed

$$C\mu^2\hbar^{-1}(\hbar|\sin(t)|/\varepsilon^2t^2)^l|\sin(t)|^{-1}.$$

Therefore summation with respect to t along zone (1.36) intersected with k-th tick returns $C\mu h^{-1}(\hbar/\varepsilon|k|)^l$ as $|k| \ge \bar{k}$ with $\bar{k}$ defined by (1.32); then summation by k results in $C\mu h^{-1}(\hbar/\varepsilon)^l$ as $\hbar \le \varepsilon$ $(1 \le |k|)$ and $C\mu h^{-1}\bar{k} \asymp C\mu^2\varepsilon^{-1}$ as $\hbar \ge \varepsilon$ $(\bar{k} \le |k|)$.

In the latter case as $1 \le |k| \le \bar{k}$ we integrate by parts only as $|\sin(t)| \le \hbar^{-1}\varepsilon^2t^2$ and contribution of k-th tick to (1.22) does not exceed

---

[4] I.e. interval $\{t:\ |t-t_k| \le \pi - \epsilon_0\}$.

$C\mu h^{-1}\log(\bar{k}/|k|)$ where logarithmic factor is an integral of $|\csc(t)|$ along k-th tick intersected with $\{t:\,\hbar^{-1}\varepsilon^2t^2\le|\sin(t)|\le\varepsilon|t|\}$. Then summation with respect to $1\le|k|\le\bar{k}$ returns $O(\mu h^{-1}\bar{k})$ again.

Therefore we arrive to

(1.37) Contribution of zone (1.36) to expression (1.22) does not exceed $C\mu h^{-1}(\hbar/\varepsilon)^l$ as $\hbar\le\varepsilon$ and $C\mu^2\varepsilon^{-1}$ as $\hbar\ge\varepsilon$.

In the same time in the zone

$$\bigl\{t:\ |\sin(t)|\ge(1+\epsilon_0)\varepsilon|t|,\ |t|\ge\epsilon_0\bigr\}\tag{1.38}$$

$|\varphi'|\asymp 1$ and $|\varphi''|\lesssim|\sin(t)|^{-1}$ and both factors do not exceed $\hbar/|\sin(t)|$. Again summation with respect to t along k-th tick intersected with zone (1.38) returns $C\mu h^{-1}(\hbar/\varepsilon|k|)^l$ as $|k|\ge\bar{k}$ and summation by k results in $C\mu h^{-1}(\hbar/\varepsilon)^l$ as $\hbar\le\varepsilon$ $(1\le|k|)$ and $\mu^2\varepsilon^{-1}$ as $\hbar\ge\varepsilon$ $(\bar{k}\le|k|)$.

In the latter case as $|k|\le\bar{k}$ we do not integrate by parts and contribution of k-th tick to (1.22) does not exceed $C\mu h^{-1}$; summation with respect to $1\le|k|\le\bar{k}$ returns $O(\mu^2\varepsilon^{-1})$. Therefore

(1.39) Contribution of zone (1.38) to expression (1.22) does not exceed $C\mu h^{-1}(\hbar/\varepsilon)^l$ as $\hbar\le\varepsilon$ and $C\mu^2\varepsilon^{-1}$ as $\hbar\ge\varepsilon$.

Note that as $\hbar\le\epsilon\varepsilon$ these considered zones (1.36) and (1.38) cover completely the singular near-pole zone. On the other hand, as $\hbar\ge\epsilon\varepsilon$ we did not integrate by parts in the second zone as $|k|\le\bar{k}$ and therefore it can be extended to $\{t:\,|\sin(t)|\ge\epsilon_0\varepsilon|t|\}$ and we arrive to

(1.40) Contribution of singular near-pole zone to expression (1.22) does not exceed $C\mu h^{-1}(\hbar/\varepsilon)^l$ as $\hbar\le\varepsilon$ and $C\mu^2\varepsilon^{-1}$ as $\hbar\ge\varepsilon$.

Claims (1.37), (1.39), (1.39) imply that as $|\cos(t_k)|\ge\epsilon$ we can apply the stationary phase method.

Then contribution of k-th stationary point to (1.22) does not exceed $C\mu h^{-1}(\hbar/\varepsilon|k|)^{\frac{1}{2}}$ and summation with respect to $k:\,\max(1,\bar{k})\le|k|\le C_0\varepsilon^{-1}$ returns $C\mu h^{-1}(\hbar/\varepsilon|k|)^{\frac{1}{2}}\times|k|$ calculated as $|k|=\varepsilon^{-1}$ i.e. $C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}$ which is larger than what we got in (1.37)–(1.40) thus resulting in

(1.41) Contribution of zone $\{t:\,|\cos(t)|\ge\epsilon_0,\ |t|\ge\epsilon_0\}$ to expression (1.22) does not exceed $C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}$ as $1\le\mu\le h^{-1}$.

Consider now the part of the near-equator zone $\{|\cos(t)| \le \epsilon_0,\ |t| \ge \epsilon_0\}$ not covered by (1.37), (1.39); there $(1-\epsilon)t^* \le |t| \le (1+\epsilon)t^*$ where

(1.42) $t^* = (k^* + \frac{1}{2})\pi$ is the closest to ε^{-1} number of the form $(k + \frac{1}{2})\pi$.

Plugging $t = t^* - s$ with $|s| \le \epsilon t^*$, $|\sin(s)| \le \epsilon$ we find that

$$\tau^{-1}\varphi'(t) = -\varepsilon^2 s(2t^* - s) + \sin^2(s) + O(\varepsilon^2) + O(\varepsilon|\sin(s)|). \tag{1.43}$$

Consider first zone

$$\bigl\{s:\ C_0 \le |s| \le \epsilon\varepsilon^{-1},\ |\sin(s)|^2 \le 2(1-\epsilon_1)\varepsilon|s|\bigr\}$$

where $|\varphi'| \gtrsim \varepsilon|s|$. Integrating by parts we acquire factor $(\hbar/\ell|\varphi'|) \asymp (\hbar/(\varepsilon|s|)^{\frac{3}{2}})$ because $\ell = (\varepsilon|s|)^{\frac{1}{2}}$ is the scale; thus we integrate multiple times by parts as $|s| \gtrsim \varepsilon^{-1}\hbar^{2/3}$; so contribution of k-th tick to (1.22) does not exceed

$$C\mu\varepsilon^{\frac{1}{2}}h^{-1}(\hbar^{2/3}/\varepsilon|k-k^*|)^{\frac{3}{2}l}|k-k^*|^{\frac{1}{2}}.$$

Then, as $\varepsilon \ge \hbar^{\frac{2}{3}}$ summation with respect to $k : |k-k^*| \ge C_0$ returns $C\mu\varepsilon^{\frac{1}{2}}h^{-1}(\hbar^{\frac{2}{3}}/\varepsilon)^l$. On the other hand, as $\varepsilon \le \hbar^{\frac{2}{3}}$ summation with respect to $k : |k-k^*| \ge \varepsilon^{-1}\hbar^{\frac{2}{3}}$ returns $C\mu\varepsilon^{\frac{1}{2}}h^{-1} \times \hbar^{\frac{2}{3}}\varepsilon^{-1} = C\mu^{\frac{5}{3}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{3}}$.

Further, the contribution of the zone

$$\bigl\{s:\ C_0 \le |s| \le \epsilon\varepsilon^{-1},\ |\sin(s)|^2 \ge 2(1+\epsilon_1)\varepsilon|s|\bigr\}$$

is estimated in the same way. Finally, as $s<0$ both terms in the right-hand expression of (1.43) have the same sign and we can extend zones in question to cover $\{s:\ \sin^2(s) \asymp \varepsilon|s|\}$ as well and we arrive to

(1.44) Contribution to (1.22) of the zone $\{s:\ |\sin(s)| \le \epsilon,\ |s| \le \epsilon\varepsilon^{-1}\}$ without subzone

$$\bigl\{s:\ s \ge C_0\max(1, \varepsilon^{-1}\hbar^{\frac{2}{3}}),\ (2-\epsilon)\varepsilon s \le \sin^2(s) \le (2+\epsilon)\varepsilon s\bigr\} \tag{1.45}$$

and the near-equator singular zone

$$\bigl\{s:\ |s| \le C_0\max(1, \varepsilon^{-1}\hbar^{\frac{2}{3}}),\ |\sin(s)| \le C_0\max(\varepsilon, \hbar^{\frac{1}{3}})\bigr\} \tag{1.46}$$

does not exceed $C\mu\varepsilon^{\frac{1}{2}}h^{-1}(\hbar^{\frac{2}{3}}/\varepsilon)^l$ as $\varepsilon \ge \hbar^{\frac{2}{3}}$ and $C\mu^{\frac{5}{3}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{3}}$ as $\varepsilon \le \hbar^{\frac{2}{3}}$.

Obviously,

(1.47) Contribution to expression (1.22) of zone (1.46) does not exceed $C\mu h^{-1}\times \max(1,\varepsilon^{-1}\hbar^{\frac{2}{3}})\times \max(\varepsilon,\hbar^{\frac{1}{3}}) = C\mu h^{-1}\times \max(\varepsilon, \varepsilon^{-1}\hbar, \hbar^{\frac{1}{3}})$.

Finally, in zone (1.45) we can use the stationary phase method. Recall that $\varphi''\asymp \sin(s)\asymp (\varepsilon|k-k^*|)^{\frac{1}{2}}$; therefore the contribution of k-th tick does not exceed $C\mu h^{-1}(\hbar/|\sin(s)|)^{\frac{1}{2}}\asymp C\mu h^{-1}(\hbar/(\varepsilon|k-k^*|)^{\frac{1}{2}})^{\frac{1}{2}}$ and summation with respect to $k:\ |k-k^*|\le \varepsilon^{-1}$ returns $C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}$.

Combining all these estimates we arrive to estimate (1.49) below; adding

$$|F_{t\to \hbar^{-1}\tau}\bigl(\bar\chi_{\bar T}(t)U(0,0,t)\bigr)|\le C\mu h^{-1} \tag{1.48}$$

which is the standard semiclassical estimate $O(\hbar^{-1})$ with an extra factor μ^2 due to rescaling we arrive to estimate (1.50):

Proposition 1.2.4. *For the pilot-model operator* (1.1) *with* $\tau\asymp 1$, $\varepsilon\le\mu^{-1}$, $\mu\le h^{-1}$ *in* $\mathbb{R}^2$

$$|F_{t\to \hbar^{-1}\tau}\bigl(\bigl(\bar\chi_T(t)-\bar\chi_{\bar T}\bigr)U(0,0,t)\bigr)|\le C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}} \tag{1.49}$$

as $\bar T=\epsilon_0\le T\le\infty$ *and*

$$|F_{t\to \hbar^{-1}\tau}\bigl(\bar\chi_T(t)U(0,0,t)\bigr)|\le C\mu h^{-1}+C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}. \tag{1.50}$$

1.2.2 Tauberian estimate

Remark 1.2.5. (i) Recall that for the pilot-model operator (1.1) in domain X, $B(0,\ell)\subset X\subset\mathbb{R}^2$ with $\ell\ge C_0\mu^{-1}$, fundamental solution U coincides modulo negligible with one in $\mathbb{R}^2$ as $T\le T^*=\epsilon_0\mu\varepsilon^{-1}\ell$.

(ii) We have $T^*=\epsilon_0\mu\varepsilon^{-1}\ell$ rather than $T^*=\epsilon_0\varepsilon^{-1}\ell$ as in Chapter 13 only because we rescaled $t\mapsto\mu t$; without this rescaling we would have $F_{t\to h^{-1}\tau}$ rather than $F_{t\to\hbar^{-1}\tau}$ and the right-hand expressions (1.49), (1.50) should be multiplied by μ^{-1}.

Due to above we arrive to the statement (i) of proposition 1.2.6 below; then the standard Tauberian arguments yield that the Tauberian error does not exceed the right-hand expression of (1.50) divided by T^*:

Proposition 1.2.6. *For a self-adjoint operator in* $\mathscr{L}^2(X)$ *in the domain* X, $B(0,\ell)\subset X\subset\mathbb{R}^2$, $\ell\ge C_0\mu^{-1}$, *coinciding in* $B(0,\ell)$ *with the pilot-model operator* (1.1) *with* $\tau\asymp 1$, $\varepsilon\le\mu^{-1}$, $\mu\le h^{-1}$,

(i) Estimates (1.49), (1.50) *hold as* $T\le T^*=\epsilon_0\mu\varepsilon^{-1}\ell$;

(ii) Tauberian estimate

$$|e(0,0,\tau)-e^{\mathsf{T}}(0,0,\tau)|\le C\bigl(\mu h^{-1}+\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}\bigr)\mu^{-1}\varepsilon\ell^{-1}= C\bigl(\varepsilon h^{-1}+\mu^{\frac{1}{2}}h^{-\frac{1}{2}}\bigr)\ell^{-1} \tag{1.51}$$

holds with the Tauberian expression

$$e^{\mathsf{T}}(x,y,\tau):=\hbar^{-1}\int_{-\infty}^{\tau}\Bigl(F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)U(x,y,t)\Bigr)\,d\tau \tag{1.52}$$

with any $T\in(C_0\varepsilon^{-1},\epsilon_0\mu\varepsilon^{-1}\ell)$.

1.2.3 Micro-averaging

Note that the second terms in the right-hand expressions of (1.50), (1.51) are due to the loops and according to Chapter 13 these expressions would not appear if instead of $\Gamma_x e(.,.,\tau)$ we looked at $\Gamma(e(.,.,\tau)\psi)$ with the uniformly smooth function ψ. Now we want to consider $\Gamma\bigl(e(.,.,\tau)\psi_\gamma\bigr)$ where ψ_γ is γ-admissible function and $\gamma\le\ell$.

So, let us consider $\Gamma\bigl(e(.,.,\tau)\psi_\gamma\bigr)$. First of all note that all the above estimates will acquire factor γ^2 due to the integration, but there is more: integrating by parts with respect to x_1 and remembering that due to rescaling $x\mapsto\mu x$ we also rescale $\gamma\mapsto\mu\gamma$ we conclude that the contribution of k-th tick acquires factor $(\hbar/\varepsilon|k|\mu\gamma)^l=(h/\varepsilon\gamma|k|)^l$ as

$$|k|\ge\hat{k}(\gamma):=h/\varepsilon\gamma. \tag{1.53}$$

This factor "forces" $|k|=\max(\hat{k}(\gamma),1)$ in the estimate of the Fourier transform

$$F_{t\to\hbar-1}\Gamma(U\psi_\gamma) \tag{1.54}$$

and there are four cases:

(a) $\gamma \ge h/\varepsilon$; then $\hat{k}(\gamma) = 1$ and expression (1.54) does not exceed $C\mu h^{-1}\min\bigl((\hbar/\varepsilon)^{\frac{1}{2}},1\bigr)\times (h/\varepsilon\gamma)^l\gamma^2$;

(b) $h/\varepsilon \ge \gamma \ge \mu^{-1}$. Then $1 \le \hat{k}(\gamma) \le \bar{k} = \hbar/\varepsilon$ and (1.54) does not exceed $C\mu h^{-1}\hat{k}(\gamma)\gamma^2 = C\mu\varepsilon^{-1}\gamma$;

(c) $\min(h/\varepsilon,\mu^{-1}) \ge \gamma \ge h$. Then $\bar{k} \le \hat{k}(\gamma) \le \varepsilon^{-1}$ and (1.54) does not exceed $C\mu h^{-1}(\hbar/\varepsilon)^{\frac{1}{2}}|\hat{k}(\gamma)|^{\frac{1}{2}}\gamma^2 = C\mu^{\frac{3}{2}}\varepsilon^{-1}\gamma^{\frac{3}{2}}$;

(d) $\gamma \le h$. Then $\hat{k}(\gamma) \ge \varepsilon^{-1}$ and micro-averaging brings no improvement and estimates (1.55)–(1.58) below are due to estimates (1.49)–(1.51) without micro-averaging.

Therefore we arrive to

Proposition 1.2.7. *For the pilot-model operator* (1.1) *with* $\tau \asymp 1$, $\varepsilon \le \mu^{-1}$, $\mu \le h^{-1}$ *in* $\mathbb{R}^2$ *estimates*

$$\gamma^{-2}|F_{t\to\hbar^{-1}\tau}\bigl((\bar{\chi}_T(t)-\bar{\chi}_{\bar{T}})\Gamma(U\psi_\gamma)\bigr)| \le C\mu\varepsilon^{-1}R^{\mathsf{T}}(\gamma) \tag{1.55}$$

and

$$\gamma^{-2}|F_{t\to\hbar^{-1}\tau}\bigl(\bar{\chi}_T(t)\Gamma(U\psi_\gamma)\bigr)| \le C\mu h^{-1} + C\mu\varepsilon^{-1}R^{\mathsf{T}}(\gamma) \tag{1.56}$$

hold as $\bar{T} = \epsilon_0 \le T \le \infty$ *where*

$$R^{\mathsf{T}}(\gamma) = \begin{cases} \mu^{\frac{1}{2}}h^{-\frac{1}{2}} & \text{as } \gamma \le h,\\ \mu^{\frac{1}{2}}\gamma^{-\frac{1}{2}} & \text{as } \min(h/\varepsilon,\mu^{-1}) \ge \gamma \ge h,\\ \gamma^{-1} & \text{as } h\varepsilon^{-1} \ge \gamma \ge \mu^{-1},\\ \varepsilon h^{-1}\min\bigl((\hbar/\varepsilon)^{\frac{1}{2}},1\bigr)\times(h/\varepsilon\gamma)^l & \text{as } \gamma \ge h\varepsilon^{-1}. \end{cases} \tag{1.57}$$

Then the standard Tauberian arguments imply

Corollary 1.2.8. *In frames of proposition 1.2.6*

$$\gamma^{-2}|\Gamma\Bigl(\bigl(e(.,.,\tau)-e^{\mathsf{T}}(.,.,\tau)\bigr)\psi_\gamma\Bigr)| \le C\bigl(\varepsilon h^{-1}+R^{\mathsf{T}}(\gamma)\bigr)\ell^{-1}. \tag{1.58}$$

1.3 Calculations

1.3.1 Correction term

We want however more explicit expression for (1.52) which we rewrite as

$$\hbar^{-1}\int_{-\infty}^{\tau}\Bigl(F_{t\to\hbar^{-1}\tau}\bar{\chi}_{\bar{T}}(t)U(x,x,t)\Bigr)\,d\tau+\\ \int_{-\infty}^{\tau}\Bigl(F_{t\to\hbar^{-1}\tau}(-it)^{-1}\bigl(\bar{\chi}_{T}(t)-\bar{\chi}_{\bar{T}}(t)\bigr)U(x,x,t)\Bigr)\,d\tau \tag{1.59}$$

and as $\mu \le h^{\delta-1}$ the first term modulo negligible is a Weyl expression $h^{-2}\mathcal{N}_x^{\mathsf{W}} := (4\pi)^{-1}h^{-2}(\tau - V(x))$ (provided $V(x)\le \tau-\epsilon$) and the second expression is a *correction term* $h^{-2}\mathcal{N}_{x,\mathsf{corr}}$ and our purpose is either to estimate or calculate it.

Let us apply the stationary phase method whenever in makes sense; otherwise we just temporarily skip the corresponding zone. Then the main part of the answer is

$$\sum_{k\in\mathfrak{Z}}\frac{1}{4\pi\sqrt{\pi}}\mu^2\hbar^{-\frac{1}{2}}(t_k\sin(t_k))^{-1}|\varphi''(t_k)|^{-\frac{1}{2}}\times\\ \exp\Bigl(\frac{i\pi}{4}\operatorname{sign}\varphi''(t_k)+i\hbar^{-1}\varphi(t_k)\Bigr) \tag{1.60}$$

with

$$\varphi''(t_k)\sim 2\tau\cot(t_k)\qquad \text{as }\ |\cos(t_k)|\ge C\varepsilon^{\frac{1}{2}} \tag{1.61}$$

and

$$\mathfrak{Z} := \bigl\{k\ne 0:\ |\sin(t_k)|\ge\hbar,\ |\cos(t_k)|\ge C\max\bigl(\varepsilon^{\frac{1}{2}},\hbar^{\frac{1}{3}}\bigr)\bigr\} \tag{1.62}$$

where dimmed restriction is just temporary.

1.3.2 Estimating correction term

First, let us estimate the correction term. As $|\sin(t_k)|\ge\hbar$, $|\cos(t_k)|\ge\epsilon$ we have $t_k\asymp k$, $\sin(t_k)\asymp\varepsilon k$, $\varphi''\asymp\varepsilon^{-1}k^{-1}$ and the contribution of k-th tick does not exceed $C\mu^{\frac{3}{2}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}}|k|^{-\frac{3}{2}}$.

In comparison with the previous subsection we acquired factor $|k|^{-1}$ due to the factor $(-it)^{-1}$ in (1.59) and it is a game-changer: summation with respect to k returns $C\mu^{\frac{3}{2}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}}|k|^{-\frac{1}{2}}$ with the *smallest* possible k: as $\hbar \le \varepsilon$ we sum from $|k|=1$ and get $C\mu^{\frac{3}{2}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}}$; as $\hbar \ge \varepsilon$ we sum from $|k| = \bar{k} := \hbar\varepsilon^{-1}$ and get $C\mu h^{-1}$ which is the trivial estimate.

Further, contribution of k-th tick to the error is $C\mu h^{-1} \times (\hbar/\varepsilon|k|)^{r+\frac{1}{2}}|k|^{-1}$ if we use r-term stationary phase approximation. Then summation with respect to $k : |k| \ge 1$ returns $C\mu h^{-1}(\hbar/\varepsilon)^{r+\frac{1}{2}}$ as $\hbar \le \varepsilon$.

On the other hand, as $\hbar \ge \varepsilon$ summation with respect to $k : |k| \ge \bar{k}$ returns $C\mu h^{-1}$ and thus this approximation does not make any sense. Therefore

(1.63) We assume until the end of this subsection that $\hbar \le \varepsilon$.

This assumption takes care of many problems: it eliminates the near-pole singular zone and away from the stationary points we can many times integrate by parts; one can see easily that contribution of zones (1.37) and (1.39) does not exceed $C\mu h^{-1}(\hbar/\varepsilon)^{l}$ with arbitrarily large l.

Therefore we need an additional analysis only in the the near-equator zone $\{t :\ |\cos(t)| \le \epsilon,\ |t - t^*| \le \epsilon\varepsilon^{-1}\}$. It is easy to estimate its contribution to the correction term: in comparison with the previous subsection we gain factor ε (from factor $(-it)^{-1}$) and we get $C\mu^2\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}} \times \varepsilon = C\mu^2\varepsilon^{\frac{1}{2}}h^{-\frac{1}{2}}$ which is less than the contribution of the near-polar zone. Thus we arrive to

Proposition 1.3.1. *For a pilot-model operator* (1.1) *with* $\tau \asymp 1$, $\varepsilon \le \mu^{-1}$, $\hbar \le \varepsilon$

(i) Correction term $h^{-2}\mathcal{N}_{x,\mathsf{corr}}$ *does not exceed* $C\mu^{\frac{3}{2}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}}$ *and therefore in frames of proposition 1.2.6*

$$\mathsf{R}^{\mathsf{W}}_x := |e^{\mathsf{T}}(x,x,0) - h^{-2}\mathcal{N}^{\mathsf{W}}_x| \le C\mu^{-1}h^{-1} + C\mu^{\frac{3}{2}}\varepsilon^{-\frac{1}{2}}h^{-\frac{1}{2}}. \tag{1.64}$$

(ii) In particular, $\mathsf{R}^{\mathsf{W}}_x \le C\mu^{-1}h^{-1}$ *as* $\varepsilon \ge \mu^5 h$.

1.3.3 Calculating correction term

Situation in the near-equator zone becomes more delicate if we want to estimate an error in r-term stationary phase approximation. Then the contribution of k-th tick does not exceed

$$C\mu\varepsilon h^{-1}(\hbar/|\cos(t_k)|^3)^{r+\frac{1}{2}} \asymp C\mu\varepsilon h^{-1}\bigl(\hbar/(\varepsilon|k-k^*|)^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}$$

which as $r\ge 1$ sums to $C\mu\varepsilon h^{-1}\bigl(\hbar/(\varepsilon|k-k^*|)^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}|k-k^*|$ calculated at the least possible value of $|k-k^*|$; so we get $C\mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}$ as we sum for $|k-k^*|\ge 1$, i.e. as $\varepsilon\ge\hbar^{\frac{2}{3}}$.

On the other hand, we get $C\mu\varepsilon h^{-1}\times\hbar^{\frac{2}{3}}\varepsilon^{-1}=C\mu^{\frac{5}{3}}h^{-\frac{1}{3}}$ as we sum for $|k-k^*|\ge\hbar^{\frac{2}{3}}\varepsilon^{-1}$, i.e. as $\varepsilon\le\hbar^{\frac{2}{3}}$. This takes care of zone (1.45); analysis of zones covered by (1.44) is easy and its contribution to the error is lesser.

Finally, the contribution to the error of zone (1.46) does not exceed $C\mu\varepsilon h^{-1}\max(\varepsilon,\varepsilon^{-1}\hbar,\hbar^{\frac{1}{3}})$ where in comparison with the analysis of the previous subsection we acquired factor ε.

Therefore, contribution of the near-equator zone to the error does not exceed

$$C\mu^{-1}h^{-1}+\left\{\begin{aligned}&C\mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}+C\mu^{\frac{4}{3}}\varepsilon h^{-\frac{2}{3}}\qquad &&\text{as}\ \ \varepsilon\ge\hbar^{\frac{2}{3}},\\ &C\mu^{\frac{5}{3}}h^{-\frac{1}{3}} &&\text{as}\ \ \varepsilon\le\hbar^{\frac{2}{3}}\end{aligned}\right.\tag{1.65}$$

and we arrive to

Proposition 1.3.2. *For a pilot-model operator* (1.1) *with* $\tau\asymp 1$, $\varepsilon\le\mu^{-1}$, $\hbar\le\varepsilon$ *correction term* $h^{-2}\mathcal{N}_{x,\mathsf{corr}}$ *is delivered with* r*-term stationary phase approximation* $h^{-2}\mathcal{N}_{x,\mathsf{corr}(r)}$ *with an error not exceeding*

$$C\mu h^{-1}\bigl(\hbar/\varepsilon\bigr)^{r+\frac{1}{2}}+C\mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}+C\mu^{\frac{4}{3}}\varepsilon h^{-\frac{2}{3}}+C\mu^{-1}h^{-1}\qquad \textit{as}\ \ \varepsilon\ge\hbar^{\frac{2}{3}}\tag{1.66}$$

and

$$C\mu h^{-1}\bigl(\hbar/\varepsilon\bigr)^{r+\frac{1}{2}}+\mu^{\frac{5}{3}}h^{-\frac{1}{3}}+C\mu^{-1}h^{-1}\qquad \textit{as}\ \ \varepsilon\le\hbar^{\frac{2}{3}}.\tag{1.67}$$

1.3.4 Micro-averaging

In the current framework (including assumption $\hbar\le\varepsilon$) micro-averaging does not improve estimate of the correction term or the error as $|\cos(t)|\ge\epsilon_0$ in it unless $\gamma\ge h\varepsilon^{-1}$ (in comparison with the remainder estimate which was improved as $\gamma\ge h$). Then, as $\gamma\ge h\varepsilon^{-1}$ in estimates (1.66), (1.67) the first term gains factor $(h/\varepsilon\gamma)^l$, the last term is preserved and all other terms gain factor $(h/\gamma)^l\le\varepsilon^l$ which effectively kills them and we arrive to estimates (1.68), (1.71) below.

As $h \le \gamma \le h\varepsilon^{-1}$ the first and the last terms do not improve but all further terms gain factor $(h/\gamma)^l$ and we arrive to estimates (1.69) and (1.70) below. Finally, for $\gamma \le \mu h$ micro-averaging brings no improvement.

Proposition 1.3.3. *(i) For a pilot-model operator* (1.1) *with* $\tau \asymp 1$, $\varepsilon \le \mu^{-1}$, $\varepsilon \ge \hbar$,

$$\begin{aligned} \mathsf{R}^{\mathsf{W}}_{x(r),\gamma} := \gamma^{-2} | \int \Bigl(e^{\mathsf{T}}(x,x,0) - h^{-2}\bigl(\mathcal{N}^{\mathsf{W}}_x + \mathcal{N}_{x,\mathsf{corr}(r)}\bigr)\Bigr)\psi_\gamma\, dx| \le \\ C\mu h^{-1}\bigl(\hbar/\varepsilon\bigr)^{r+\frac{1}{2}}\bigl(h/\varepsilon\gamma\bigr)^l + C\mu^{-1}h^{-1} \qquad \textit{as}\ \ \gamma \ge h/\varepsilon, \end{aligned} \tag{1.68}$$

$$\begin{aligned} \mathsf{R}^{\mathsf{W}}_{x(r),\gamma} \le C\mu h^{-1}\bigl(\hbar/\varepsilon\bigr)^{r+\frac{1}{2}} + C\mu\varepsilon\Bigl(h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}} + \mu^{\frac{1}{3}}h^{-\frac{2}{3}}\Bigr)\bigl(h/\gamma\bigr)^l \\ + C\mu^{-1}h^{-1} \qquad \textit{as}\ \ \varepsilon \ge \hbar^{\frac{2}{3}},\ \ h \le \gamma \le h/\varepsilon \end{aligned} \tag{1.69}$$

and

$$\begin{aligned} \mathsf{R}^{\mathsf{W}}_{x(r),\gamma} \le C\mu h^{-1}\bigl(\hbar/\varepsilon\bigr)^{r+\frac{1}{2}} + C\mu^{\frac{5}{3}}h^{-\frac{1}{3}}\bigl(h/\gamma\bigr)^l + C\mu^{-1}h^{-1} \\ \textit{as}\ \ \varepsilon \le \hbar^{\frac{2}{3}},\ \ h \le \gamma \le h/\varepsilon, \end{aligned} \tag{1.70}$$

(ii) For a pilot-model operator (1.1) *with* $\tau \asymp 1$, $\varepsilon \le \mu^{-1}$, $\varepsilon \le \hbar$

$$\begin{aligned} \mathsf{R}^{\mathsf{W}}_{x,\gamma} := \gamma^{-2} | \int \Bigl(e^{\mathsf{T}}(x,x,0) - h^{-2}\mathcal{N}^{\mathsf{W}}_x\Bigr)\psi_\gamma\, dx| \le \\ \mu h^{-1}\bigl(h/\varepsilon\gamma\bigr)^l + C\mu^{-1}h^{-1} \qquad \textit{as}\ \ \gamma \ge h/\varepsilon. \end{aligned} \tag{1.71}$$

1.4 Strong and superstrong magnetic field

1.4.1 Tauberian estimate

Consider strong $\mu h \asymp 1$ and superstrong $\mu h \gtrsim 1$ magnetic field for a Schrödinger-Pauli pilot-model operator

$$A := h^2D_1^2 + (hD_2 - \mu x_1)^2 + 2\alpha x_1 - \mathfrak{z}\mu h \tag{1.72}$$

which alternatively means that we assume that $|\tau - \mathfrak{z}\mu h| \le C$ in the framework of the standard pilot-model (1.1). Formulae (1.15)–(1.16) imply that

Proposition 1.4.1. *For a pilot-model operator* (1.1) *with* $\mu h \gtrsim 1$ *in* $\mathbb{R}^2$ *as* $|x| \le C_0$, $|y| \le C_0$

(i) $e(x, y, \tau) \equiv 0 \mod O(\mu^{-\infty})$ *as* $\tau \le \mu h - \epsilon_0$ *(the lower spectral gap);*

(ii) $e(x, y, \tau) \equiv e(x, y, \tau')$ *and* $\partial_\tau e(x, y, \tau) \equiv 0 \mod O(\mu^{-\infty})$ *as*

$$(2n-1) + \epsilon_0 \le \tau' < \tau \le (2n+1)\mu h - \epsilon_0 \text{ (other spectral gaps);}$$

(iii) $e(x, x, \tau) - e(x, x, \tau') \asymp \mu h^{-1}$ *as* $\tau' < (2n-1)\mu h - \epsilon_0$ *and* $\tau > (2n-1)\mu h + \epsilon_0$*;*

(iv) As $\varepsilon > 0$ *the following estimates hold*

$$|\partial_\tau e(x, y, \tau)| \le C\mu^{\frac{1}{2}}\varepsilon^{-1}h^{-\frac{3}{2}} \tag{1.73}$$

and

$$|F_{t\to\hbar^{-1}\tau} U(x, y, t)| \le C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}} \tag{1.74}$$

and these estimate are sharp as $x = y$ *and* τ *is close to Landau levels* $(2n+1)\mu h$*.*

Further, as before $T^* = \epsilon_0\mu\varepsilon^{-1}\ell$. Then we immediately arrive to

Corollary 1.4.2. *For a self-adjoint operator in domain* X, $B(0, \ell) \subset X \subset \mathbb{R}^2$, $\ell \ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$, *coinciding in* $B(0, \ell)$ *with the pilot-model operator* (1.72) *with* $\tau \asymp 1$, $\varepsilon < 0$, $\mu \ge h^{-1}$

(i) Statements (i)-(iv) of proposition 1.4.1 remain true;

(ii) Formula (1.15) *holds modulo* $O(\mu^{\frac{1}{2}}h^{-\frac{1}{2}}\ell^{-1})$*;*

(iii) Moreover, formula (1.15) *holds modulo* $O(\mu^{-\infty})$

as $(2n-1) + \epsilon_0 \le \tau \le (2n+1)\mu h - \epsilon_0$*;*

1.4.2 Micro-averaging

Let us consider micro-averaging. First let us estimate Fourier transform where as before we rescaled $t \mapsto \mu t$; formulae (1.15)–(1.16) imply immediately

Proposition 1.4.3. *For a pilot-model operator* (1.72) *in* $\mathbb{R}^2$ *with* $\mu h \ge 1$ *and* $\mathfrak{z} = (2n+1)$

$$\gamma^{-2}|F_{t\to h^{-1}\tau}\bar{\chi}_T(t)\Gamma(U\psi_\gamma)| \le C\mu^2 + C\mu\varepsilon^{-1}R^{\mathsf{T}}(\gamma) \tag{1.75}$$

with

$$R^{\mathsf{T}}(\gamma)=\left\{\begin{aligned} &\mu^{\frac12}h^{-\frac12} &&\textit{as}\quad \gamma\le \mu^{-\frac12}h^{\frac12},\\ &\gamma^{-1} &&\textit{as}\quad \gamma\ge \mu^{-\frac12}h^{\frac12}.\end{aligned}\right. \tag{1.76}$$

Corollary 1.4.4. *In frames of corollary 1.4.2*

$$\gamma^{-2}|\Gamma\bigl(e(.,.,\tau)-e^{\mathsf{T}}(.,.,\tau)\bigr)\psi_\gamma|\le C\bigl(\mu\varepsilon+R^{\mathsf{T}}(\gamma)\bigr)\ell^{-1}. \tag{1.77}$$

1.5 Weyl and magnetic Weyl approximation

We can try to use a host of the different approximations but restrict ourselves now to Weyl approximation, which makes sense only as $\mu h\le 1$. Recall that

$$|e^{\mathsf{T}}(x,x,\tau)-h^{-2}\mathcal{N}_x^{\mathsf{W}}(\tau)|\le CR^{\mathsf{W}}, \tag{1.78}$$

$$\gamma^{-2}|\int\bigl(e^{\mathsf{T}}(x,x,\tau)-h^{-2}\mathcal{N}_x^{\mathsf{W}}(\tau)\bigr)\psi_\gamma\,dx|\le CR^{\mathsf{W}}(\gamma) \tag{1.79}$$

with

$$R^{\mathsf{W}}=\left\{\begin{aligned} &\mu^{\frac32}\varepsilon^{-\frac12}h^{-\frac12} &&\text{as}\quad \varepsilon\ge\hbar,\\ &\mu h^{-1} &&\text{as}\quad \varepsilon\le\hbar,\end{aligned}\right. \tag{1.80}$$

$$R^{\mathsf{W}}(\gamma)=\left\{\begin{aligned} &R^{\mathsf{W}} &&\text{as}\quad \gamma\le h/\varepsilon,\\ &R^{\mathsf{W}}(h/\varepsilon\gamma)^{l} &&\text{as}\quad \gamma\ge h/\varepsilon.\end{aligned}\right. \tag{1.81}$$

On the other hand, let us apply instead the magnetic Weyl approximation. Obviously, without micro-averaging $\mathcal{N}_x^{\mathsf{MW}}(\tau)$ cannot produce uniform with respect to τ remainder estimate better than μh^{-1} due to its own jumps. Meanwhile with an averaging (i.e. a micro-averaging with $\gamma=1$) $\mathcal{N}_x^{\mathsf{MW}}(\tau)$ provides a better remainder estimate $O(\mu^{-1}h^{-1})$ (see Chapter 13; for a pilot-model we need only assume that $\varepsilon\ge\mu^{-l}$). So, as $\mu h\le 1$ the magnetic Weyl approximation may provide a better remainder estimate as γ is not too small enough but for small γ Weyl approximation is better. Let us investigate this.

Proposition 1.5.1. *(i) For a pilot-model operator* (1.1) *as* $\mu h\lesssim 1$

$$\mathsf{R}^{\mathsf{MW}}(\gamma):=\gamma^{-2}|\int\bigl(e^{\mathsf{T}}(x,x,0)-h^{-2}\int\mathcal{N}^{\mathsf{MW}}\bigr)\psi_\gamma\,dx|\le C\mu^{-1}h^{-1}+CR^{\mathsf{MW}}(\gamma) \tag{1.82}$$

with

$$R^{\mathsf{MW}}(\gamma)=\begin{cases}\mu^{-1}h^{-1}\gamma^{-2}(h/\varepsilon\gamma)^{l} & \textit{as}\ \ \gamma\ge \max(\mu^{-1},h/\varepsilon),\\ \mu^{-1}h^{-1}\gamma^{-2} & \textit{as}\ \ \mu^{-1}\le\gamma\le h/\varepsilon,\\ \mu h^{-1}(h/\varepsilon\gamma)^{l} & \textit{as}\ \ h/\varepsilon\le\gamma\le\mu^{-1},\\ \mu h^{-1} & \textit{as}\ \ \gamma\le\min(\mu^{-1},h/\varepsilon);\end{cases}\tag{1.83}$$

(ii) For a pilot-model operator (1.72) *as* $\mu h\gtrsim 1$ *and* $\mathfrak{z}=2n+1$ *estimate*

$$\mathsf{R}^{\mathsf{MW}}(\gamma)\le C+CR^{\mathsf{MW}}(\gamma)\tag{1.84}$$

holds with

$$R^{\mathsf{MW}}(\gamma)=\begin{cases}\gamma^{-2} & \textit{as}\ \ \gamma\ge\mu^{-\frac{1}{2}}h^{\frac{1}{2}},\\ \mu h^{-1} & \textit{as}\ \ \gamma\le\mu^{-\frac{1}{2}}h^{\frac{1}{2}}.\end{cases}\tag{1.85}$$

Proof. We need to investigate only case $\mu^{-1}\le\gamma\le h\varepsilon^{-1}$ as in all other cases we can easily pass from results for $\mathsf{R}^{\mathsf{W}}(\gamma)$.

Note that $h^{-2}\mathcal{N}_y^{\mathsf{MW}}(\tau)$ delivers $e(y,y,\tau)$ for a pilot-model operator $\bar{A}:=\bar{A}_y=\bar{A}_y(x,D_x)$ defined by (1.1) or (1.72) with αx_1 frozen at the point y (with respect to which we integrate later); so we consider our pilot-model operator as a perturbation of this one.

Let us plug into Tauberian formula the Schwartz kernel of

$$e^{i\hbar^{-1}tA}=e^{i\hbar^{-1}t\bar{A}}+i\hbar^{-1}\int_0^t e^{i\hbar^{-1}t'\bar{A}}\bigl(A-\bar{A}\bigr)e^{i\hbar^{-1}(t-t')A}\,dt'\tag{1.86}$$

or its iteration for $\pm t>0$

$$(1.86)_m\quad e^{i\hbar^{-1}tA}=e^{i\hbar^{-1}t\bar{A}_x}+\sum_{1\le j\le m-1}\times$$

$$(i\hbar^{-1})^j\int\limits_{\substack{\{\pm t_1>0,\pm t_2>0,\dots,\pm t_j>0,\\ t_1+t_2+\dots+t_j=t\}}} e^{i\hbar^{-1}t_1\bar{A}}\bigl(A-\bar{A}\bigr)e^{i\hbar^{-1}t_2\bar{A}}\bigl(A-\bar{A}\bigr)\cdots\bigl(A-\bar{A}\bigr)e^{i\hbar^{-1}t_j\bar{A}}\,dt_1dt_2\cdots dt_j+$$

$$(i\hbar^{-1})^m\int\limits_{\substack{\{\pm t_1>0,\pm t_2>0,\dots,\pm t_m>0,\\ t_1+t_2+\dots+t_j=t\}}} e^{i\hbar^{-1}t_1\bar{A}}\bigl(A-\bar{A}\bigr)e^{i\hbar^{-1}t_2\bar{A}}\bigl(A-\bar{A}\bigr)\cdots\bigl(A-\bar{A}\bigr)e^{i\hbar^{-1}t_mA}\,dt_1dt_2\cdots dt_m.$$

So let us plug $(1.86)_m$ into our expression with $\chi_T(t)$ instead of $\bar{\chi}_T(t)$. Then our standard methods imply that j-th term ($j = 0, \dots, m$) of what we got does not exceed

$$C\mu h^{-1}\bigl(T\|A - \bar{A}_x\|\hbar^{-1}\bigr)^j \times \bigl(1 + \frac{T}{\bar{T}_2}\bigr)^{-l}\bigl(1 + \frac{T}{\bar{T}_1}\bigr)^{-l} \tag{1.87}$$

with $\bar{T}_1 = C_0\mu\varepsilon^{-1}\bar{\gamma}$ where $\bar{\gamma} := \max(\mu^{-1}, \mu^{-\frac{1}{2}}h^{\frac{1}{2}})$ and $\bar{T}_2 = h/\varepsilon\gamma$; we leave easy details to the reader.

Then we can replace the norm by the "effective norm" which is a norm restricted to $\mu\bar{\gamma}$-vicinity of x (in rescaled coordinates) and equal to $C\alpha\bar{\gamma}$.

Note that $\bar{T}_2 \le \bar{T}_1$ as $\gamma \ge \bar{\gamma}$; then summation with respect to T returns

$$C\mu h^{-1}\bigl(\bar{T}_2\|A - \bar{A}_x\|\hbar^{-1}\bigr)^j \asymp C\mu h^{-1}\bigl(\bar{\gamma}/\gamma\bigr)^j$$

which provides required estimate (1.83) or (1.84) for all terms with $j \ge 2$.

Finally note that while $\bar{A}_y(x-y, D_x)$ is even with respect to $(x-y, D_x)$ (we can always achieve it by the gauge transformation), perturbation $(A - \bar{A}_y)(x-y, D_x)$ is odd and therefore only terms with even j survive as we plug $x = y$. This takes care of term with $j = 1$. □

Remark 1.5.2. (i) For $\mu h \le 1$ magnetic Weyl formula provides better approximation as $\mu^{-1} \le \gamma \le h/\varepsilon$; otherwise Weyl formula provides either better or equally bad (μh^{-1}) approximation.

(ii) For $\mu h \ge 1$ magnetic Weyl formula provides approximation rather than just main term estimate as $\gamma \gg \mu^{-\frac{1}{2}}h^{\frac{1}{2}}$.

1.6 Geometric interpretation

Remark 1.6.1. (i) As we mentioned t_k are length of the classical loops at point 0 (or x if we consider $e(x,x,\tau)$; then $t_k = t_k(x)$). While picture of the fixed trajectory and a point x on it (Figure 2a) is more geometrically appealing, the correct picture is of the fixed point and different trajectories passing through it (Figure 2b); due to assumption $\alpha \ne 0$ "radii" of these trajectories differ by $O(\mu^{-1})$.

Therefore, when we talk here about *loop* we mean a trajectory looping at the fixed point x rather the self-intersections of the fixed trajectory.



(a) Intuitive (b) Correct picture

Figure 2: On the left more geometrically appealing picture, when x moves along given trajectory; on the right the correct one when we consider different trajectories passing through a fixed point.

Similarly, when we talk about *pole* or *equator* we mean not x near a pole or an equator of the fixed trajectory but rather trajectory such that x is near its pole or equator.

So, all further remarks should be interpreted correctly.

(ii) How to interpret our results from heuristic uncertainty principle? Obviously these results mean that as $\varepsilon \gg \hbar$ then the majority of trajectories on Figure 2b do not return to the original point $x = 0$ after $k = 1$ tick[5] while otherwise this happens only after k ticks with $k\varepsilon \gg \hbar$. As the spatial shift is εk (after rescaling) this means that the thickness of the trajectory is $\asymp \hbar$ after rescaling (and therefore $\asymp h$ before rescaling).

It also means that if $|\sin(\theta)| \asymp 1$ where θ is the polar angle (between direction of the trajectory at 0 and the drift direction $(0,1)$) then we can take an interval in θ of the magnitude 1, so θ is a dual variable to x_2. Conversely, with the exception of the interval of the length $\hbar/\varepsilon|k|$ around θ_k (where θ_k corresponds to the classical trajectory returning to 0 after k ticks) trajectories do not return at 0; we can rewrite this condition as

$$|\sin(\theta - \theta_k)| \gg \hbar/(|k|\varepsilon). \tag{1.88}$$

---

[5] In the sense that the difference between two points is observable.

This characteristic length matches to the stationary phase estimates we derived rigorously.


(a) (b) (c)

Figure 3: Looping near pole from microlocal point of view

(iii) Near equator situation changes drastically: spacing between self-intersections is $\asymp \varepsilon|\cos(\theta)|^{-1}$ as $|\cos(\theta)|\ge \epsilon_0\varepsilon^{\frac{1}{2}}$ and there are exceptional $0,1,2$ self-intersections with $|\cos(\theta)|\le \epsilon_0\varepsilon^{\frac{1}{2}}$. Uncertainty principle may prevent us to know how many of them are no matter how "not large" μ is.


(a) (b) (c) (d)

Figure 4: Self-intersections near equator. The first winding shown by a solid line, the closest to it near equator by a dashed one.

Consequent self-intersections are distinguishable as $|\cos(\theta)|\ge \hbar^{\frac{1}{3}}$ according to our calculations.

Problem 1.6.2. *We know that estimates (1.73), (1.74) are sharp as* $\mu h \gtrsim 1$*. Are these estimates sharp as* $\mu h \ll 1$*?*

2 Pointwise asymptotics: general 2D-operators

Let us recreate the approximation similar to the one of the previous section for general operators assuming that $\hbar \le \epsilon$ i.e.

$$\mu \le \epsilon h^{-1}. \tag{2.1}$$

We are interested in $|t| \le C_0\mu^{-1}\varepsilon^{-1}$ for the original (not rescaled) operator as for $C_0\mu^{-1}\varepsilon^{-1} \le |t| \le \mu^{-1}T^* := \epsilon_0\varepsilon^{-1}\ell$ dynamics leaves $B(0, c_0\mu^{-1})$ but remains in $B(0,\ell)$. Here

(2.2) $\varepsilon = \alpha^{-1}\mu^{-1}$ as we assume that $|\nabla V/F| \asymp \alpha$ in $B(0,\ell)$ with

$$C_1\mu^{-1} \le \alpha \le 1 \qquad (\implies C_0\mu^{-2} \le \varepsilon \le \mu^{-1}) \tag{2.3}$$

and $\ell \ge c_0\mu^{-1}$ with sufficiently large constants C_1, c_0.

2.1 Classical dynamics

First we consider classical dynamics starting from point $\mathsf{x} = 0$. So far we use not-rescaled x, t. Let us freeze g^{jk} and F there and replace V_j and V by their linear germs:

$$\begin{aligned} \bar{g}^{jk} = g^{jk}(0), \quad \bar{V}(x) = V(0) + \langle \nabla V(0), x\rangle,\\ \bar{V}_j(x) = V_j(0) + \langle \nabla V_j(0), x\rangle \quad (\implies \bar{F} = F(0)), \end{aligned} \tag{2.4}$$

denote corresponding Hamiltonian by $\bar{a}(x,\xi)$ and all the object associated with it will have bar. We consider original our dynamic system as a perturbation.

Without any loss of the generality we can assume that

$$g^{jk} = \omega^2\delta_{jk} \qquad (\implies a(x,\xi) = \omega^2(p_1^2 + p_2^2) + V), \tag{2.5}$$

$$\omega(0) = F(0) = 1, \quad (\nabla\omega)(0) = 0, \tag{2.6}$$

$$V(0) = 0, \quad (\nabla V/F)(0) = (-\alpha, 0) \tag{2.7}$$

as we can achieve it by an appropriate change of variables (see footnote 9)) and

$$V_1(0) = 0, \quad V_2 = 0 \tag{2.8}$$

as we can achieve it by the gauge transformation.

Proposition 2.1.1. *Under conditions* (2.1)–(2.8) *as* $|t| \le c_0\mu^{-1}\varepsilon^{-1}$

(i) For drift flows $z_t = \Phi_t(0)$, $\bar{z}_t = \bar{\Phi}_t(0)$

$$z_t = \bar{z}_t + O\bigl(\mu^{-1}\varepsilon t^2\bigr); \tag{2.9}$$

(ii) If $F = 1$ *then for Hamiltonian flows* $(x_t, \xi_t) = \Psi_t(0, \eta)$, $(\bar{x}_t, \bar{\xi}_t) = \bar{\Psi}_t(0, \eta)$ *with magnetic parameters* μ *and* $\bar{\mu} = \mu(1 + O(\mu^{-2}))$ *defined by* (2.18) *on the energy level* $\tau \le c$

$$(x_t, \mu^{-1}\xi_t) = (\bar{x}_t, \mu^{-1}\bar{\xi}_t) + O\bigl(\mu^{-2}t\bigr). \tag{2.10}$$

Remark 2.1.2. Assumption (2.1) means that in $B(0, \mu^{-1})$ the constant part of ∇V dominates over its variable part and it implies that as

$$\mu^{-1}\varepsilon t^2 \le c_0\mu^{-2}|t| \le \kappa\varepsilon|t| \qquad \text{as} \quad |t| \le c_0\mu^{-1}\varepsilon^{-1} \tag{2.11}$$

with

$$\kappa = c_0\mu^{-2}\varepsilon^{-1} \le C_0^{-1}c_0 \tag{2.12}$$

and thus the perturbation of the drift is respectively small. In particular drift line deviates from the straight one by no more than $c_0\kappa$.

Also *equator* is defined as a point where x_t intersect its its first winding the last time deviates from $(\mu^{-1}, 0)$ by no more than $c_0\mu^{-1}\kappa$.

Proof of proposition 2.1.1. (i) Proof of assertion (i) is trivial and left to the reader. Just recall that the drift flow is defined by (13.6.5) with an extra factor μ^{-1} in the right-hand expression and in our assumptions $\bar{\Phi}_t$: $(x_1, x_2) \mapsto (x_1 + \varepsilon t, x_2)$ after rescaling.

(ii) Recall that $p_j = \xi_j - \mu V_j(x)$, $\{p_j, x_k\} = \delta_{jk}$ and in virtue of assumption $F = 1$

$$\{p_1, p_2\} = \mu\omega^{-2}. \tag{2.13}$$

According to subsection 13.2.1 one can correct

$$y_1 := x_1 - \{p_1, p_2\}^{-1} p_2 = x_1 - \mu^{-1}\omega^2 p_2,\\ y_2 := x_2 + \{p_1, p_2\}^{-1} p_1 = x_2 + \mu^{-1}\omega^2 p_1 \tag{2.14}$$

modulo $O(\mu^{-2})$ so that corrected expressions satisfy drift equation modulo $O(\mu^{-2})$. More precisely, in the current setup

$$\{a, y_1\} = \mu^{-1}\omega^2\Bigl(\{p_2, \omega^2\}(p_1^2 - p_2^2) - 2\{p_1, \omega^2\}p_1 p_2 + \{p_2, V\}\Bigr)$$

and for $y_1' = y_1 + \mu^{-2}\beta_1 p_1 p_2 + \mu^{-2}\beta_2(p_1^2 - p_2^2)$ with

$$\beta_1 = -\frac{1}{2}\omega^2\{p_2, \omega^2\}, \qquad \beta_2 = \frac{1}{2}\omega^2\{p_1, \omega^2\}$$

we have

$$\{a, y_1'\} \equiv -\mu^{-1}\omega^2\{p_2, V\}$$

modulo terms which are $O(\mu^{-2})$ and also homogeneous polynomials of degrees 1 or 3 with respect to (p_1, p_2) and thus these terms could be corrected by adding to y_1' terms which are $O(\mu^{-3})$ and also homogeneous polynomials of degrees 2 or 4; then we arrive to

$$\{a, y_k''\} \equiv (-1)^k \mu^{-1}\omega^2\{p_{3-k}, V\} \mod O(\mu^{-3}) \qquad k = 1, 2$$

($k = 2$ is considered in the same way as $k = 1$). Note that the right-hand expressions are calculated at point $x = x_t$ rather than $y'' = y_t''$; so we rewrite them as

$$\{a, y_k''\} \equiv\\ (-1)^k\mu^{-1}\bigl(\omega^2\{p_{3-k}, V\}\bigr)(y'') + \sum_{j=1,2}\mu^{-2}(-1)^{j+k-1}(\partial_{3-j}\omega^2\partial_{3-k}V)(x)p_j\\ \mod O(\mu^{-3}) \qquad k = 1, 2$$

and we can correct y_k'' by adding to y_k'' terms which are $O(\mu^{-3})$ and also homogeneous polynomials of degrees 2 with respect to (p_1, p_2).

Therefore

$$y_t''' \equiv z_t \mod O(\mu^{-3}t) \tag{2.15}$$

where $z_t = \Phi_t(0)$.

Meanwhile (2.13) implies

$$\frac{d}{dt}p_j = 2(-1)^j\mu p_{3-j} - (\omega^2)_{x_j}(p_1^2+p_2^2) - V_{x_j} = \\ 2(-1)^j\mu p_{3-j} - 2\omega_{x_j}\omega^{-1}W + W_{x_j}, \qquad j=1,2 \tag{2.16}$$

where we replaced $p_1^2+p_2^2$ by $\omega^{-2}W$, $W=\tau-V$ (no error as τ is an energy level). Let us plug $x_1(t)\equiv z_1(t)+\mu^{-1}\omega^2(z_t)p_2$, $x_2(t)\equiv z_2(t)-\mu^{-1}\omega^2(z_t)p_1$ modulo $O(\mu^{-2})$ in the right-hand expressions; we get

$$\frac{d}{dt}\begin{pmatrix}p_1\\p_2\end{pmatrix} = J\begin{pmatrix}p_1\\p_2\end{pmatrix} + K \tag{2.17}$$

with

$$J := 2\mu\begin{pmatrix} -\mu^{-2}\omega^2(\omega_{x_1}\omega^{-1}W+W_{x_1})_{x_2} & -\bigl(1-\mu^{-2}\omega^2(\omega_{x_1}\omega^{-1}W+W_{x_1})_{x_1}\bigr)\\ \bigl(1-\mu^{-2}\omega^2(\omega_{x_2}\omega^{-1}W+W_{x_2})_{x_2}\bigr) & \mu^{-2}\omega^2(\omega_{x_2}\omega^{-1}W+W_{x_2})_{x_1}\end{pmatrix},$$

and

$$K := \begin{pmatrix} -2(\omega_{x_1}\omega^{-1}W+W_{x_1})\\ -2\omega_{x_2}\omega^{-1}W+W_{x_2}\end{pmatrix}.$$

and here we can calculate elements of J at any point of $B(0,\mu^{-1})$ we choose (and we choose 0) while elements of K are calculated at z_t. Then J becomes a constant coefficient matrix and since $\nabla\omega(0)=0$, $\omega(0)=1$

$$J = 2\mu\begin{pmatrix} -\mu^{-2}(\omega_{x_1x_2}W+W_{x_1x_2}) & -\bigl(1-\mu^{-2}(\omega_{x_1x_1}W+W_{x_1x_1})\bigr)\\ \bigl(1-\mu^{-2}(\omega_{x_2x_2}W+W_{x_2x_2})\bigr) & \mu^{-2}(\omega_{x_2x_1}W+W_{x_2x_1})\end{pmatrix}$$

and its eigenvalues are $\pm 2\bar\mu i$ with

$$\bar\mu = \\ \mu\Bigl((1-\mu^{-2}(\omega_{x_1x_1}W+W_{x_1x_1}))(1-\mu^{-2}(\omega_{x_2x_2}W+W_{x_2x_2})-\mu^{-4}(\omega_{x_1x_2}W+W_{x_1x_2})^2\Bigr)^{\frac12} \\ = \mu+\mu^{-1}(\Delta(\omega W))+O(\mu^{-3}). \tag{2.18}$$

Then $J=Q^{-1}\bar J Q$ with $\bar J = \bar\mu\begin{pmatrix}0 & -1\\ 1 & 0\end{pmatrix}$ and $Q = I + O(\mu^{-2})$.

Due to (2.17)

$$p(t) \equiv e^{tJ}p(0) + \int_0^t e^{(t-t')J}K(z_{t'})\,dt' = e^{tJ}p(0) + \int_0^t e^{(t-t')J}J^{-1}\frac{d}{dt'}K(z_{t'})\,dt' - J^{-1}K(z_t) + e^{tJ}J^{-1}K(z_0). \tag{2.19}$$

The second term in the right-hand expression is $O(\mu^{-1}\varepsilon t)$ as $J^{-1}=O(\mu^{-1})$ and $\frac{d}{dt'}K(z_{t'})=O(\varepsilon)$. Note that with $O(\mu^{-2}t)$ error one can replace in the two last terms J by $\bar{J}$ calculated for a pilot-model. Finally, with an error $O(\mu^{-1}\varepsilon t)$ one can replace $K(z_t)$ by $K(z_0)$; however $K(z_0)$ and $\bar{K}(z_0)$ coincide as $\nabla\omega(0)=0$.

Replacing J by $\bar{J}$ we reduced evolution of p to those of the pilot-model albeit with $\bar{\mu}$ instead of μ. Therefore $\mu^{-1}p_t \equiv \mu^{-1}\bar{p}_t \mod O(\mu^{-2}t)$. Replacing μ by $\bar{\mu}$ does not affect drift (in frames of the indicated precision).

But then x_t could be found from z_t and p_t and their drift also is described by a pilot-model in frames of the same error and then it is true for $\mu^{-1}\xi_t$ as well. □

Consider now the general case, i.e. F different from 1 and variable. Then differential equations describing $(x(t),\xi(t))$ for $a(x,\xi)$ coincide with equations for $F^{-1}a(x,\xi)$ with $F\mapsto 1$ and $\tau-V\mapsto F^{-1}(\tau-V)$ but with the "time" θ satisfying

$$\frac{d\theta}{dt} = F(x_t) \equiv F(z_t) + (\omega^2F^{-1})(z_t)\bigl(F_{x_1}(z_t)p_2 - F_{x_2}(z_t)p_1\bigr) \mod O(\mu^{-2})$$

where $z_t=\Phi_t(0)$. We can correct θ by $O(\mu^{-1})$ eliminating linear terms in the right hand expression and therefore

$$\theta \equiv \int F(z_t)\,dt \equiv F(0)t + \frac{1}{2}\bigl(\frac{d}{dt}F(z_t)\bigr)(0)t^2 \mod O(\mu^{-1}+\mu^{-2}|t|) \tag{2.20}$$

with the second term in the right-hand expression $O(\varepsilon\mu^{-1}t^2)$.

From now on we use rescaling $x\mapsto \mu x$, $t\mapsto \bar{\mu}t$, $\theta\mapsto\mu\theta$. Our analysis implies immediately

Proposition 2.1.3. *Let conditions* (2.1)–(2.8) *be fulfilled*[6]. *Then (after rescaling $x \to \bar{\mu}x$, $t \mapsto \bar{\mu}t$) as $|t| \le c_0\varepsilon^{-1}$*

$$|D^\beta \Psi_t| \le C_\beta \qquad \forall\beta. \tag{2.21}$$

2.2 Semiclassical approximation to $U(x,y,t)$

Proposition 2.2.1. *Let conditions* (2.1)–(2.8) *be fulfilled*[6]. *Then*

(i) Uniformly with respect to $|t| \le c\varepsilon^{-1}$ $e^{i\hbar^{-1}tA}$ is an $\hbar$-Fourier integral operator corresponding to Hamiltonian flow Ψ_t;

(ii) As $|\sin(2\theta)| \ge \epsilon$ with θ defined by (2.20)

$$U(x,y,t) \equiv (4\pi\hbar)^{-1} i(\sin(\theta))^{-1} e^{i\hbar^{-1}\phi(x,y,t)} \sum_m b_m(x,y,t)\hbar^m \tag{2.22}$$

with ϕ defined by (2.25)–(2.29) *below and satisfying (with all derivatives)*

$$\phi = \bar{\phi}(\theta) + O(\mu^{-2}\varepsilon^{-1}), \tag{2.23}$$

$$b_m = \delta_{0m} + O(\mu^{-2}\varepsilon^{-1}) \tag{2.24}$$

with $\bar{\phi}$ defined by (1.10).

Proof. Both assertions of proposition are standard as $|t| \le T = \mathsf{const}$ but we need to extend them for larger t. Further, ϕ is an *action*[7]

$$\phi = -\int_0^t L\bigl(x(t'), \dot{x}(t')\bigr)\, dt' \tag{2.25}$$

with the Lagrangian

$$L(x,\dot{x}) = \bigl(\sum_k \dot{x}_k\xi_k - a(x,\xi)\bigr)|_{\xi_k=(2.29)} = \frac{1}{4}\sum_{j,k} g_{jk}(x)\dot{x}_j\dot{x}_k + \frac{1}{2}\sum_j \dot{x}_j V_j(x) - V(x). \tag{2.26}$$

---

[6] Before rescaling.

[7] Sign "–" is due to considering of propagator $e^{i\hbar^{-1}tA}$ rather than $e^{-i\hbar^{-1}tA}$.

which is the Legendre transformation of the Hamiltonian

$$a(x,\xi)=\sum_{j,k} g^{jk}(x)\bigl(\xi_j-V_j(x)\bigr)\bigl(\xi_k-V_k(x)\bigr)+V(x) \tag{2.27}$$

$$\dot{x}_j=2\sum_k g^{jk}(x)\bigl(\xi_k-V_k(x)\bigr), \tag{2.28}$$

$$\xi_k=\frac{1}{2}\sum\sum_j g_{jk}\dot{x}_j+V_k(x), \tag{2.29}$$

$\dot{f}:=df/dt'$ and we are talking about trajectories from y as $t'=0$ to x as $t'=t$.

To extend assertions (i), (ii) to t: $|t|\le T=c\varepsilon^{-1}$ note that

$$e^{\pi t\hbar^{-1}A}=-e^{i\mu^{-1}\hbar^{-1}B} \tag{2.30}$$

with $\hbar$-pseudo-differential operator B; $B=0$ if A is a pilot-model operator with $\alpha=0$. Therefore as usual

$$e^{\pi t\hbar^{-1}A}=e^{ik\hbar^{-1}A}e^{i(t-\pi k)\hbar^{-1}A}=(-1)^k e^{ik\mu^{-1}\hbar^{-1}B}e^{i(t-\pi k)\hbar^{-1}A} \tag{2.31}$$

with $k=\lfloor t/\pi\rfloor$ and the right-hand expression is $\hbar$-Fourier integral operator as $|k|\le c\varepsilon^{-1}$. Assertion (i) is proven.

Note that the corresponding canonical manifold is well-projected to (x,y) space ($\mathbb{R}^4$) as $|\sin(2\theta)|\ge\epsilon$ and compare it with the canonical manifold for the pilot-model operator; this proves (ii) as either $F=1$ or $|t|\le\epsilon_1\varepsilon$ with sufficiently small $\epsilon_1=\epsilon_1(\epsilon)$.

But then one can represent $e^{i\hbar^{-1}tA}$ as $e^{i\hbar^{-1}t_1A}e^{i\hbar^{-1}t_2A}\cdots e^{i\hbar^{-1}t_nA}$ with $n\le c_1$ and $|t_j|\le\epsilon_1\mu$, $|\sin(2t_j)|\ge\frac{1}{2}\epsilon$ which implies (ii) in the general case. $\square$

So far we exclude both vicinities of $\sin(\theta)=0$ which matches $x=y$ (poles) and $\cos(\theta)=0$ which matches x and y being antipodal points of the trajectory (equator) but we need to approach both of them. Actually exclusion of the latter was no more than a precaution but poles require a modification:

Proposition 2.2.2. *Let conditions* (2.1)–(2.8) *be fulfilled*[6]*. Then*

(i) Decomposition (2.22) *remains valid as* $|\cos(\theta)|\le\epsilon$*;*

(ii) Decomposition (2.22) *remains valid as*

$$C\max\bigl(\hbar, \mu^{-1}\varepsilon|t|\bigr) \le |\sin(\theta)| \le \epsilon \tag{2.32}$$

albeit with an error not exceeding

$$C\hbar^{-1}|\sin(\theta)|^{-1}(\hbar/|\sin(\theta)|)^{l} \tag{2.33}$$

and with b_m *such that*

$$|D^{\beta}(\phi-\bar{\phi})| \le C_{\beta}\varepsilon|\sin(\theta)|^{-|\beta|} \qquad \forall \beta, \tag{2.34}$$

$$|D^{\beta}(b_m-\delta_{m0})| \le C_{m\beta}\varepsilon|\sin(\theta)|^{-m-|\beta|} \qquad \forall \beta, m. \tag{2.35}$$

Proof. As

$$U(x,y,t)=\mu^{-2}\int U(x,z,t')U(z,y,-t'')\,dz$$

for $t=t'-t''$ (where factor μ^{-d} is due to rescaling U as a function) we need to consider this oscillatory integral. If we consider oscillatory integral with the propagator for the pilot-model operator we note that the standard stationary phase method applies with an effective semiclassical parameter $\hbar$ as $(\cot(t')-\cot(t''))=-\sin(t)\sec(t')\sec(t'')$ disjoint from 0 i.e. also as $|\cos(t)|\le\epsilon$.

On the other hand, an effective semiclassical parameter is $\hbar/|\sin(t)|$ near poles.

In virtue of proposition 2.2.1 both these claims remain true for a general operator as well albeit with t replaced by θ. ☐

Remark 2.2.3. So far we need only $|\nabla V/F|\lesssim \mu\varepsilon$ rather than $|\nabla V/F|\asymp \mu\varepsilon$ [6)].

2.3 Semiclassical approximation to $e(x,x,\tau)$

Therefore in zone $\{|\sin(\theta)|\ge C\hbar\}$ all arguments of the pilot-model theory work (with obvious modifications) under non-degeneracy assumptions

$$V-\tau\le-\epsilon_0, \tag{2.36}$$

$$|\nabla(\tau-V)F^{-1}|\asymp\varepsilon \tag{2.37}$$

we conclude that

(2.38) Contribution of zone $\{|\sin(\theta)| \geq C\hbar\}$ to $F_{t\to h^{-1}\tau}(1-\bar{\chi}_1(t))\Gamma_x U$ does not exceed $C\mu h^{-1} + C\mu^{\frac{3}{2}}\varepsilon^{-1}h^{-\frac{1}{2}}$.

Consider now zone $\{|\sin(\theta)| \leq \epsilon|\}$. Both operators A and B (defined by (2.30) are ξ-microhyperbolic and unless factors they are coming with (namely $(\theta - \pi k)$ (with the closest πk) and εt respectively are of the same magnitude we can use ξ-microhyperbolicity to prove

(2.39) Contribution of zone $\{|t| \asymp T \geq \varepsilon^{-1}\max(\mu\hbar, C|\sin(\theta)|)\}$ to $F_{t\to h^{-1}\tau}\Gamma_x U$ does not exceed

$$C\mu^2\hbar^{-2} \times T \times \varepsilon T \times (\hbar/\varepsilon T)^{l+1} = C\mu^2 h^{-1} T (\hbar/\varepsilon T)^l$$

(with arbitrarily large l) while its contribution to expression (1.59) does not exceed the same expression albeit without factor T i.e.

$$C\mu^2 h^{-1}(\hbar/\varepsilon T)^l,$$

where factor μ^2 is due to rescaling, $T\times\varepsilon T$ is the measure of the zone $\{\theta : |\theta| \asymp T,\ |\sin(\theta)| \leq \varepsilon T\}$ and other factors are standard; recall that B comes with the factor ε and therefore effective semiclassical parameter is $\hbar/\varepsilon$. We leave easy details to the reader.

After summation with respect to T we conclude that

(2.40) As $\varepsilon \geq C\hbar$ contributions of zone $\{|t| \geq 1,\ |\sin(\theta)| \leq c\hbar\}$ to both $F_{t\to h^{-1}\tau}\Gamma_x U$ and expression (1.59) do not exceed $C\mu h^{-1}(\hbar/\varepsilon)^l$

and

(2.41) As $\varepsilon \leq C_0\hbar \leq 1$ contribution of zone $\{|t| \geq C_0\hbar/\varepsilon,\ |\sin(\theta)| \leq c\hbar\}$ to $F_{t\to h^{-1}\tau}\Gamma_x U$ does not exceed $C\mu^2\varepsilon^{-1}$ while its contribution to expression (1.59) does not exceed $C\mu h^{-1}$.

So, as $\varepsilon \geq C\hbar$ we are done but as $\varepsilon \leq C\hbar$ we need to consider zone $\{1 \leq |t| \leq c\hbar/\varepsilon,\ |\sin(\theta)| \leq c\hbar\}$ and the same arguments imply

(2.42) As $\varepsilon \leq C_0\hbar \leq 1$ contribution of zone $\{1 \leq |t| \leq c\hbar/\varepsilon,\ |\sin(\theta)| \leq c\hbar\}$ to $F_{t\to h^{-1}\tau}\Gamma_x U$ does not exceed

$$C\mu^2\hbar^{-2} \times \varepsilon^{-1}\hbar \times \hbar = C\mu^2\varepsilon^{-1}$$

where factor μ^2 is due to rescaling, $\varepsilon^{-1}\hbar \times \hbar$ is the measure of the zone $\{\theta : |\theta| \leq \varepsilon^{-1}\hbar,\ |\sin(\theta)| \leq \hbar\}$ and factor $\hbar^{-2}$ is the standard one.

Combining with the results for zone $\{|\sin(\theta)| \ge c\hbar\}$ we arrive to

Proposition 2.3.1. *For magnetic Schrödinger operator in domain* X, $B(0,\ell) \subset X \subset \mathbb{R}^2$ *under standard smoothness assumptions and non-degeneracy assumptions* (0.6) *and* (2.36)–(2.37), (2.3) *after rescaling* $x \mapsto \mu x$, $t \mapsto \mu t$ *(as* $\mu \le h^{-1}$*) estimates* (1.49)–(1.50) *of proposition 1.2.4 and estimates* (1.55)–(1.57) *of proposition 1.2.7 hold as* $T \le c\varepsilon^{-1}$.

Further, estimates (1.78)–(1.81) *hold as well.*

Now we apply the standard Tauberian arguments. We are interested mainly in the case of $\alpha \asymp 1$; then automatically $\ell \asymp 1$ and $T^* \asymp \mu^2$ inside of domain. However in more general case $\ell \asymp \alpha \asymp \mu\varepsilon$ automatically and $T^* \asymp \ell/\varepsilon\mu\mu^2 \asymp \mu^2$ again and we arrive to

Proposition 2.3.2. *For magnetic Schrödinger operator in domain* X, $B(0,\ell) \subset X \subset \mathbb{R}^2$ *under standard smoothness assumptions and non-degeneracy assumptions* (0.6) *and* (2.36)–(2.37), (2.3) *Tauberian estimates* (1.51) *of proposition 1.2.6 and estimates* (1.57)–(1.58) *of corollary 1.2.8 hold with the Tauberian expression* (1.52) *and* $\ell \asymp \alpha$.

2.4 Stationary phase calculations

Let apply the stationary phase method to the Tauberian expression. To do so we need to remember that assumptions $|t| \ll \varepsilon^{-1}$, $\varphi' = 0$ yield $\varphi'' \asymp |\sin(\theta)|^{-1}$ and all other estimates hold; therefore the stationary phase construction is available there. Also this construction works as long as $|t| \le c\varepsilon^{-1}$ and $|\varphi''| \ge \epsilon$ i.e. $|t - t^*| \ge \epsilon\varepsilon^{-1}$ where $T^* = \varepsilon^{-1}$.

Moreover, $|\varphi''| \ge \epsilon_0$ as $|t - t^*| \le \epsilon\varepsilon^{-1}$ and $\varphi' = 0$ and therefore all our estimates work here, in the near equator zone as well where

Definition 2.4.1. *Equator just moves to the point where* $|\varphi''|$ *is minimal for* $\varphi' = 0$. *Let us in the near equator zone redefine* θ *in such way that equator is* $\cos(\theta) = 0$.

Let us introduce

Definition 2.4.2. *Consider* r*-term representation* (2.22) *(i.e. with summation over* $m < r$*) and plug it into Tauberian expression, excepting* $|t| \le \epsilon_0$ *and calculate it by the stationary phase method with* r *terms again. Let us call the result* r-term semiclassical approximation *and denote it by* $h^{-2}\mathcal{N}_{x,\mathsf{corr}(r)}$.

Then its main term is delivered by modified (1.60)–(1.62)

$$\sum_{k\in\mathfrak{Z}} \frac{1}{4\pi\sqrt{\pi}}\mu\hbar^{-\frac{1}{2}}(t_k\sin(\theta_k))^{-1}|\varphi''(t_k)|^{-\frac{1}{2}}b_0(t_k)\times \\ \exp\Bigl(\frac{i\pi}{4}\operatorname{sign}\varphi''(t_k)+i\hbar^{-1}\varphi(t_k)\Bigr) \tag{2.43}$$

with

$$\varphi''(t_k)\sim 2\tau\cot(\theta_k)\qquad \text{as}\ \ |\cos(\theta_k)|\ge C\varepsilon^{\frac{1}{2}} \tag{2.44}$$

and

$$\mathfrak{Z}:=\bigl\{k\ne 0:\ |\cos(\theta_k)|\ge C\max\bigl(\varepsilon^{\frac{1}{2}},\hbar^{\frac{1}{3}}\bigr)\bigr\} \tag{2.45}$$

where t_k is a time of k-th return to x (along k-th loop), $\theta_k=\theta(t_k)$ and $\varphi(t_k)$ is the corresponding action.

Then repeating arguments of the previous section we arrive to

Proposition 2.4.3. *For magnetic Schrödinger operator in domain X, $B(0,1)\subset X\subset\mathbb{R}^2$ under standard smoothness assumptions and non-degeneracy assumptions* (0.6) *and* (2.36)*–*(2.37)*,* (2.3)

(i) As $\varepsilon\ge\hbar$ estimate (1.64) *of proposition 1.3.1 holds;*

(ii) Estimates (1.66)*–*(1.67) *of proposition 1.3.2, estimates* (1.68)*–*(1.71) *of proposition 1.3.3 and estimates* (1.82)*–*(1.85) *of proposition 1.5.1 hold.*

2.5 Approximation by a pilot-model operator

2.5.1 Weak magnetic field case

We cannot do better in this framework; however using the pilot-model operator as an approximation we could improve these results. Without any loss of the generality we can assume that (2.5)–(2.7) are fulfilled.

Assume temporarily that $F=1$. Then one can prove easily that

$$t_k=\theta(t_k)\equiv\bar{t}_k,\qquad \phi(t_k)\equiv\bar{\phi}(\bar{t}_k)\quad \operatorname{mod}\ \mu^{-2}|k| \tag{2.46}$$

where bar denotes objects related to the pilot-model operator. Then $\hbar^{-1}\bigl(\phi(t_k)-\bar{\phi}(\bar{t}_k)\bigr)=O(\mu^{-2}\hbar^{-1}|k|)$ and it is less than 1 as

$$|k|\le \tilde{k}:=\mu^3 h. \tag{2.47}$$

Therefore it makes sense to apply this approach only as $\tilde{k}\ge 1$ i.e.

$$\mu\ge h^{-\frac{1}{3}}. \tag{2.48}$$

Under this assumption let us compare r-th terms for our operator and the pilot model as $|k|\le\tilde{k}$.

One can see easily that their difference does not exceed

$$C\sum_k \mu h^{-1}(\hbar/\varepsilon|k|)^{r+\frac{1}{2}}|k|^{-1}\times\mu^{-3}h^{-1}|k| \tag{2.49}$$

which as $r\ge 1$ does not exceed $C\mu^{-2}h^{-2}(\hbar/\varepsilon|k|)^{r+\frac{1}{2}}|k|$ calculated for the smallest possible $|k|$ which is 1 provided $\hbar\le\varepsilon$. Therefore we get $C\mu^{-2}h^{-2}(\hbar/\varepsilon)^{r+\frac{1}{2}}$. On the other hand in (2.49) one should replace the last factor by 1 as $|k|\ge\tilde{k}$.

Therefore, if we start from l-term approximation and apply estimate (1.66), (1.67) as $\varepsilon\ge\hbar^{\frac{2}{3}}$, $\hbar\le\varepsilon\le\hbar^{\frac{2}{3}}$ respectively and pass from l-term to r-term approximation we get estimate

$$\begin{aligned}
&\mathsf{R}^{\mathsf{W}\prime\prime}_{x(r)\gamma}:=\\
&\qquad|e^{\mathsf{T}}(x,x,\tau)-h^{-2}\mathcal{N}_{x,\mathsf{corr}(r)}-\bar{e}^{\mathsf{T}}(x,x,\tau)+h^{-2}\bar{\mathcal{N}}_{x,\mathsf{corr}(r)}|\le\\
&\qquad\qquad C\mu^{-1}h^{-1}+C\mu^{-2}h^{-2}(\hbar/\varepsilon)^{r+\frac{1}{2}}+C\mu h^{-1}(\hbar/\varepsilon)^{l+\frac{1}{2}}+\\
&\qquad\qquad\qquad\begin{cases}
C\mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}}+C\mu^{\frac{4}{3}}\varepsilon h^{-\frac{2}{3}} & \text{as }\ \varepsilon\ge\hbar^{\frac{2}{3}}\\
C\mu^{\frac{5}{3}}h^{-\frac{1}{3}} & \text{as }\ \hbar\le\varepsilon\le\hbar^{\frac{2}{3}}
\end{cases}
\end{aligned} \tag{2.50}$$

where terms $-h^{-2}\mathcal{N}^{\mathsf{W}}_x$ and $h^{-2}\bar{\mathcal{N}}^{\mathsf{W}}_x$ obviously cancel one another.

2.5.2 Successive approximations

Now we need to get rid off the third term in the right-hand expression of (2.50) without condition $\varepsilon\gg\hbar$ which allows to eliminate it. To do this we need more sophisticated arguments. Namely, let us apply the successive approximations.

Assuming that the original operator is perturbed by $O(\mu^{-2})$ we arrive to the heuristic conclusion that

(2.51) Contribution of $|t| \asymp T$ to $R^{\mathsf{W}''}$ does not exceed the previous estimate of the contribution to $\mathsf{R}^{\mathsf{W}'}$ multiplied by $\mu^{-2}T\hbar^{-1}$;

this statement needs justification. We can achieve a better estimate by taking more terms in the successive approximation but it leads to a rather overcomplicated formula; so we take just one term.

Then, in frames of (2.51) we have three cases:

(a) $\varepsilon \ge \hbar$; then all what successive approximation does is to eliminate the third term in the right-hand expression of (2.50).

(b) $\mu^{-3}h^{-1} \le \varepsilon \le \hbar$; then $1 \ge \bar{k} \le \varepsilon^{-1}$ and we need to adjust our arguments treating separately $|k| \le \bar{k}$ and $|k| \ge \bar{k}$ and recalculate the second term in the right-hand expression of (2.50); however equator zone is not affected.

(c) $\varepsilon \le \min(\hbar, \mu^{-3}h^{-1})$. In this case in addition we need to recalculate the last term in the right-hand expression of (2.50).

Let us reconsider the second term in the right-hand expression of (2.50). First, expression (2.49) with summation over $k : |k| \ge \bar{k}$ returns $C\mu^{-2}h^{-2}(\hbar/\varepsilon|k|)^{r+\frac{1}{2}}|k|$ calculated as $k = \bar{k}$; so we get $C\mu^{-2}h^{-2}\bar{k} = C\mu^{-1}h^{-1}\varepsilon^{-1}$.

Further, as $|k| \le \bar{k}$ we need to replace $(\hbar/\varepsilon|k|)^{r+\frac{1}{2}}$ by 1; so we sum $C\mu h^{-1}|k|^{-1} \times \mu^{-3}h^{-1}|k|$ which returns $C\mu^{-2}h^{-2}\bar{k}$ again. Therefore[8] we conclude that

(2.52) As $\hbar \ge \varepsilon$, $r \ge 1$ one should replace the second term in (2.50) by $C\mu^{-1}h^{-1}\varepsilon^{-1}$.

Furthermore, as $\varepsilon \le \min(\hbar, \mu^{-3}h^{-1})$ one should multiply the last term in (2.50) i.e. $C\mu^{\frac{5}{3}}h^{-\frac{1}{3}}$ by $\mu^{-3}\varepsilon^{-1}h^{-1}$ resulting in $C\mu^{-\frac{4}{3}}\varepsilon^{-1}h^{-\frac{4}{3}}$. Therefore[8] we conclude that

---

[8] Pending (2.51).

(2.53) As $\varepsilon \le \min(\hbar, \mu^{-3}h^{-1})$, $r \ge 1$ one should replace the second term in (2.50) by $C\mu^{-\frac{1}{3}}\varepsilon^{-1}h^{-\frac{4}{3}}$.

Therefore we (heuristically, the proof follows) arrive to

Proposition 2.5.1. *Consider two magnetic Schrödinger operators A and $\bar{A}$ satisfying the standard smoothness assumptions, non-degeneracy assumptions* (0.6) *and* (2.36)–(2.37), (2.3) *and such that*

$$g^{jk} = \bar{g}^{jk} + O(\mu^{-2}), \qquad V = \bar{V} + O(\mu^{-2}), \qquad V_j = \bar{V}_j + O(\mu^{-3}). \tag{2.54}$$

Then for $r \ge 1$

$$\begin{aligned} \mathsf{R}^{\mathsf{W}\prime\prime}_{x(r)} \le C\mu^{-1}h^{-1} + C &\begin{cases} \mu^{-2}h^{-2}(\hbar/\varepsilon)^{r+\frac{1}{2}} & \text{as } \ \varepsilon \ge \hbar \\ C\mu^{-1}h^{-1}\varepsilon^{-1} & \text{as } \ \varepsilon \le \hbar \end{cases} \quad + \\ &\begin{cases} C\mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}} + C\mu^{\frac{4}{3}}\varepsilon h^{-\frac{2}{3}} & \text{as } \ \varepsilon \ge \hbar^{\frac{2}{3}} \\ C\mu^{\frac{5}{3}}h^{-\frac{1}{3}} & \text{as } \ \min(\hbar, \mu^{-3}h^{-1}) \le \varepsilon \le \hbar^{\frac{2}{3}} \\ C\mu^{-\frac{4}{3}}\varepsilon^{-1}h^{-\frac{4}{3}} & \text{as } \ \varepsilon \le \min(\hbar, \mu^{-3}h^{-1}) \end{cases} \end{aligned} \tag{2.55}$$

and

$$\begin{aligned} &\mathsf{R}^{\mathsf{W}\prime\prime}_{x(r)\gamma} := \\ &\gamma^{-2}\Bigl|\int \Bigl(e^{\mathsf{T}}(x,x,\tau) - h^{-2}\mathcal{N}_{x,\mathsf{corr}(r)} - \bar{e}^{\mathsf{T}}(x,x,\tau) + h^{-2}\bar{\mathcal{N}}_{x,\mathsf{corr}(r)}\Bigr)\psi_\gamma\, dx\Bigr| \le \\ &C\mu^{-1}h^{-1} + C\begin{cases} \mu^{-2}h^{-2}(\hbar/\varepsilon)^{r+\frac{1}{2}}(\hbar/\varepsilon\gamma)^{l} & \text{as } \ \gamma\varepsilon \ge \hbar \\ \mu^{-2}h^{-2}(\hbar/\varepsilon)^{r+\frac{1}{2}} & \text{as } \ \varepsilon \ge \hbar \\ \mu^{-1}h^{-1}\varepsilon^{-1} & \text{as } \ \varepsilon \le \hbar \end{cases} \quad + \\ &C\min\bigl(1, (h/\gamma)^{l}\bigr)\times\begin{cases} \mu\varepsilon h^{-1}\bigl(\hbar/\varepsilon^{\frac{3}{2}}\bigr)^{r+\frac{1}{2}} + C\mu^{\frac{4}{3}}\varepsilon h^{-\frac{2}{3}} & \text{as } \ \varepsilon \ge \hbar^{\frac{2}{3}} \\ \mu^{\frac{5}{3}}h^{-\frac{1}{3}} & \text{as } \ \min(\hbar, \mu^{-3}h^{-1}) \le \varepsilon \le \hbar^{\frac{2}{3}} \\ \mu^{-\frac{1}{3}}\varepsilon^{-1}h^{-\frac{4}{3}} & \text{as } \ \varepsilon \le \min(\hbar, \mu^{-3}h^{-1}) \end{cases} \end{aligned} \tag{2.56}$$

while for $r = 0$

$$\begin{aligned} \mathsf{R}^{\mathsf{W}\prime\prime}_{x(0)} := |e^{\mathsf{T}}(x,x,\tau) - \bar{e}^{\mathsf{T}}(x,x,\tau)| \le C\mu^{-1}h^{-1} + \\ C\begin{cases} h^{-1}\varepsilon^{-\frac{1}{2}} & \text{as } \ \mu^3 h \le \varepsilon^{-1}, \\ \mu^{-\frac{3}{2}}h^{-\frac{3}{2}}\varepsilon^{-1} & \text{as } \ \mu^3 h \ge \varepsilon^{-1} \end{cases} \end{aligned} \tag{2.57}$$

and

$$\mathsf{R}^{\mathsf{W}\prime\prime}_{x(0)\gamma} := \gamma^{-2}\bigl|\int \Bigl(e^{\mathsf{T}}(x,x,\tau) - \bar{e}^{\mathsf{T}}(x,x,\tau)\Bigr)\psi_\gamma\, dx\bigr| \le C\mu^{-1}h^{-1} + \\ C\begin{cases} \mu^{-\frac{3}{2}}h^{-1}\varepsilon^{-1}\gamma^{-\frac{1}{2}} & \textit{as}\ \ \mu^{-3}\varepsilon^{-1}\le\gamma\le\min(h\varepsilon^{-1},\mu^{-1}),\\ \mu^{-2}h^{-1}\varepsilon^{-1}\gamma^{-1} & \textit{as}\ \ \mu^{-1}\le\gamma\le h\varepsilon^{-1},\\ \mu h^{-1}\min\bigl(1,(\mu h/\varepsilon)^{\frac{1}{2}}\bigr)(h/\varepsilon\gamma)^{l} & \textit{as}\ \ \gamma\ge h\varepsilon^{-1}. \end{cases} \tag{2.58}$$

2.5.3 Justification

To justify (2.51) and thus to prove proposition 2.5.1 we need

Proposition 2.5.2. *Consider two magnetic Schrödinger operators A and $\bar{A}$ satisfying* (2.54)*. Then as $|t|\le \epsilon\mu^3 h$ and $|\sin(t)|\ge\max\bigl(\epsilon\varepsilon|t|, C\hbar\bigr)$*

$$U(x,y,t)-\bar{U}(x,y,t)\equiv \\ (4\pi\hbar)^{-1}i(\sin(\bar{\theta}(t)))^{-1}e^{i\hbar^{-1}\bar{\phi}(x,y,t)}\sum_m b'_m(x,y,t)\hbar^m \\ \mod O\Bigl(\hbar^{-2}\mu^{-3}|t||\sin(\theta(t))|^{-1}\times(\hbar/|\sin(t)|)^{l}\Bigr) \tag{2.59}$$

with

$$|D^\beta b'_m|\le C_{m\beta}\mu^{-3}|t|\hbar^{-1}|\sin(\theta(t))|^{-m-|\beta|}. \tag{2.60}$$

Proof. Obviously in our assumptions

$$U(x,y,t)\equiv(4\pi\hbar)^{-1}i(\sin(\bar{\theta}(t)))^{-1}e^{i\hbar^{-1}\bar{\phi}(x,y,t)}\sum_m b''_m(x,y,t)\hbar^m \\ \mod O\Bigl(\hbar^{-1}|\sin(\bar{\theta}(t))|^{-1}\times(\hbar/|\sin(\bar{\theta}(t))|)^{l}\Bigr)$$

with b''_m satisfying

$$|D^\alpha b''_m|\le C_{m\alpha}|\sin(\bar{\theta}(t))|^{-m-|\alpha|}$$

and plugging it into the right-hand expression of

$$e^{i\hbar^{-1}tA}-e^{i\hbar^{-1}t\bar{A}} = i\hbar^{-1}\int_0^t e^{i\hbar^{-1}t'A}(A-\bar{A})e^{i\hbar^{-1}(t-t')\bar{A}}\,dt' \tag{2.61}$$

we arrive to (2.59). Here to cover zones where either $|\sin(t')| \le \epsilon$ or $|\sin(t-t')| \le \epsilon$ we just pass to the standard representation through non-singular oscillatory integrals (with one or two extra variable) and then apply the stationary phase method. We leave easy but tedious details to the reader. $\square$

Corollary 2.5.3. *In frames of proposition 2.5.2 equality* (2.59) *holds with* $U(x,y,t)$ *and* $\bar{U}(x,y,t)$ *replaced respectively by*

$$(4\pi\hbar)^{-1} i(\sin(\theta(t)))^{-1} e^{i\hbar^{-1}\phi(x,y,t)} \sum_m b_m(x,y,t)\hbar^m \tag{2.62}$$

and

$$(4\pi\hbar)^{-1} i(\sin(\bar{\theta}(t)))^{-1} e^{i\hbar^{-1}\bar{\phi}(x,y,t)} \sum_m \bar{b}_m(x,y,t)\hbar^m. \tag{2.63}$$

2.5.4 Reduction to the pilot-model operator

From now we are interested *only* in the estimates without micro-averaging.

The problem however is that $\bar{A}$ at this moment is a more general operator than the pilot-model: it contains linear with respect to x terms in the original (before rescaling) g^{jk} and V, and also it contains quadratic terms in $V_2 = x_1 + \frac{1}{2}\beta_{11}x_1^2 + \beta_{12}x_1x_2 + \frac{1}{2}\beta_{22}x_2^2 + ...$ all of which generate $O(\mu^{-1})$ perturbation.

However one select a conformal coordinate system in such way that $\nabla\omega(0) = 0$ [9]. Then if we know that $F = F(0) + O(|x|^2)$ we conclude that $\beta_{11} = \beta_{12} = 0$ and by a gauge transformation we can make $\beta_{22} = 0$.

Further, we can assume that $\tau = 0$ and then make $V(0) = -1$ by division $V(0) \mapsto -1$, $h \mapsto h(-V(0))^{-\frac{1}{2}}$, $\mu \mapsto \mu(-V(0))^{-\frac{1}{2}}$ before rescaling. Rescaling μ (and changing coordinate orientation if needed we can make $F(0) = 1$.

Furthermore, we can assume that $F = 1$. Really, as we can consider energy level $\tau = 0$ we starting from equation $-\hbar D_t u = Au$, go to

$$F^{-1}Au = -\hbar F^{-1}D_t u = -hD_t + \hbar F^{-1}(F-1)D_t u \tag{2.64}$$

---

[9] Really, changing conformal system to another conformal system is done with $y_1 = f$, $\nabla y_2 = (-f_{x_2}, f_{x_1})$ which can be satisfied iff $\Delta f = 0$; then ω is replaced by $\omega|\nabla f|$ and if $f = ax_1 + bx_1x_2 + \frac{1}{2}c(x_1^2 - x_2^2)$ then $|\nabla f| = a + bx_2 + cx_1 + O(|x|^2)$ and selecting a, b, c we can make $\omega(0) = 1$, $\nabla\omega(0) = 0$.

and on an interval $|\tau|\le \hbar h^{-\delta}T^{-1}$ (which is completely microlocally admissible as $|t|\asymp T$ the last term is $O(\mu^{-1}\hbar h^{-\delta}T^{-1})$ and multiplying by $T\hbar^{-1}$ we get $O(\mu^{-1}h^{-\delta})$ which allows us to use the successive approximation method.

We leave to the reader to use our standard methods and solve a rather easy

Problem 2.5.4. Prove that effectively in estimates one can make $\delta=0$.

Note that $\mu^{-1}\le \mu^{-3}h^{-1}$ as $\mu\le h^{-\frac{1}{2}}$ and therefore this new error (F is replaced by $F(0)$) does not exceed the right-hand expressions of (2.55) and (2.57) as $r\ge 1$ and $r=0$ respectively.

Meanwhile our new error does not exceed $C\mu h^{-1}\times \mu^{-1}=Ch^{-1}$ anyway and as $\mu\ge h^{-\frac{1}{2}}$ ($\implies \varepsilon\le \hbar$) it is less than $\mu^{\frac{5}{3}}h^{-\frac{1}{3}}$ leave alone $\varepsilon^{-\frac{1}{2}}h^{-1}$ which are parts of right-hand expressions of (2.55) and (2.57) respectively. Thus we arrive to

(2.65) Estimates (2.55) and (2.57) hold with $\bar{A}$ which is our original pilot-model operator.

2.6 Superstrong magnetic field

Finally, let $\mu\ge h^{-1}$. Then as only matter is what is inside of the circle of the radius $C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ (we can get rid off $h^{-\delta}$ factor in estimates) we get in successive approximations $C\mu h^{-1}\times \mu^{-1}h\times \mu^{-\frac{1}{2}}h^{\frac{1}{2}}\varepsilon^{-1}\times h^{-1}$ where $\mu^{-1}h$ is the size of the perturbation, $\mu^{-\frac{1}{2}}h^{\frac{1}{2}}\varepsilon^{-1}$ is T (without rescaling); so we get $C\mu^{-\frac{1}{2}}h^{-\frac{1}{2}}\varepsilon^{-1}$.

We leave to the reader a rather easy

Problem 2.6.1. Prove that effectively in estimates one can make $\delta=0$ as well in $\mu^{-\frac{1}{2}+\delta}h^{\frac{1}{2}}$ radius and $\mu^{-\frac{1}{2}+\delta}h^{\frac{1}{2}}\varepsilon^{-1}$ time.

So we arrive to

Proposition 2.6.2. *Under assumptions* (2.36), (2.37), $F=1$ *and* (2.54) *estimate*

$$|e^{\mathsf{T}}(0,0,\tau)-\bar{e}^{\mathsf{T}}(0,0,\tau)|\le C\mu^{-\frac{1}{2}}h^{-\frac{1}{2}}\varepsilon^{-1}\qquad \textit{as}\quad \mu\ge h^{-1}. \tag{2.66}$$

We leave to the reader yet another easy

Problem 2.6.3. Write correction to τ and may be α as $F\ne \mathsf{const}$ and $g^{jk}\ne \mathsf{const}$.

2.7 Main theorem

Now we can write our main theorem of these two sections:

Theorem 2.7.1. *For magnetic Schrödinger operator in domain $X\subset\mathbb{R}^2$ such that $B(0,1)\subset X$ and satisfying conditions* (0.5)–(0.7) *the following estimates hold for $x\in B(0,\frac{1}{2})$*

(i) As $1\le\mu\le h^{-\frac{1}{2}}$

$$|e(x,x,0)-h^{-2}\mathcal{N}_x^{\mathsf{W}}(0)|\le C\mu^{-1}h^{-1}+C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}+C\mu^2h^{-\frac{1}{2}} \tag{2.67}$$

and

$$\begin{multline*}\mathsf{R}^{\mathsf{W}}_{x(r)}:=|e(x,x,0)-h^{-2}\bigl(\mathcal{N}_x^{\mathsf{W}}(0)+\mathcal{N}^{\mathsf{W}}_{x,\mathsf{corr}(r)}(0)\bigr)|\le\\ C\mu^{-1}h^{-1}+C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}+C\mu h^{-1}\bigl(\mu^2h\bigr)^{r+\frac{1}{2}}+\\ C\begin{cases}\Bigl(h^{-1}\bigl(h\mu^{\frac{5}{2}}\bigr)^{r+\frac{1}{2}}+\mu^{\frac{1}{3}}h^{-\frac{2}{3}}\Bigr) & \textit{as}\quad \mu\le h^{-\frac{2}{5}},\\ \mu^{\frac{5}{3}}h^{-\frac{1}{3}} & \textit{as}\quad \mu\ge h^{-\frac{2}{5}}.\end{cases}\end{multline*} \tag{2.68}$$

(ii) As $h^{-\frac{1}{2}}\le\mu\le h^{-\frac{1}{2}}$

$$\begin{multline*}\mathsf{R}^{\mathsf{W}\prime\prime}_{x(r)}:=\\ |\Bigl(e(x,x,\tau)-h^{-2}\mathcal{N}_{x,\mathsf{corr}(r)}-\bar{\mathsf{e}}_x(x,x,\tau)+h^{-2}\bar{\mathcal{N}}_{x,\mathsf{corr}(r)}\Bigr)|\le\\ C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}+C\begin{cases}\mu^{-2}h^{-2}(\mu^2h)^{r+\frac{1}{2}} & \textit{as}\ \ \mu\le h^{-\frac{1}{2}}\\ h^{-1} & \textit{as}\ \ \mu\le h^{-\frac{1}{2}},\end{cases}\quad+\\ C\begin{cases}h^{-1}\bigl(\mu^{\frac{5}{2}}h\bigr)^{r+\frac{1}{2}}+\mu^{\frac{1}{3}}h^{-\frac{2}{3}} & \textit{as}\ \ \mu\le h^{-\frac{2}{5}}\\ \mu^{\frac{5}{3}}h^{-\frac{1}{3}} & \textit{as}\ \ h^{-\frac{2}{5}}\le\mu\le h^{-\frac{1}{2}},\\ \mu^{-\frac{1}{3}}h^{-\frac{4}{3}} & \textit{as}\ \ \mu\ge h^{-\frac{1}{2}}\end{cases}\end{multline*} \tag{2.69}$$

while for $r=0$

$$\begin{multline*}\mathsf{R}^{\mathsf{W}\prime\prime}_{x(0)}:=|e(x,x,\tau)-\bar{\mathsf{e}}_x(x,x,\tau)|\le C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}+\\ C\begin{cases}h^{-1}\mu^{\frac{1}{2}} & \textit{as}\ \ \mu\le h^{-\frac{1}{2}},\\ \mu^{-\frac{1}{2}}h^{-\frac{3}{2}} & \textit{as}\ \ \textit{as}\ \ \mu\le h^{-\frac{1}{2}};\end{cases}\end{multline*} \tag{2.70}$$

where here and in (iii) $\bar{\mathsf{e}}_x$ *is constructed for a pilot-model in* x*;*

(iii) For magnetic Schrödinger-Pauli operator with $\mu \ge h^{-1}$

$$|e(x,x,\tau) - \bar{\mathsf{e}}_x(x,x,\tau)| \le C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}. \tag{2.71}$$

2.8 Problems

Finally, let us formulate a series of the problems with probably simple parts (i) and really difficult parts (ii):

Problem 2.8.1. *As* $\mu \le h^{-1}$ *drop condition (0.5) in the pointwise asymptotics:*

(i) Use the simple rescaling technique $\mu \mapsto \mu\rho\frac{1}{2}\mu$, $h \mapsto h\rho^{-\frac{3}{2}}$ *with the scaling function* $\rho(x) = \max(\epsilon_0|V(x)|, \mu h, \mu^{-1})$ *reducing operator to the similar one either satisfying condition (0.5)* *or* *with* $\mu h \asymp 1$ *or* *with* $\mu = 1$ *and considered in subsection 5.2.1; in two latter cases condition (0.5) is not needed;*

(ii) Using propagation, improve what follows from the simple rescaling technique.

Problem 2.8.2. *Derive asymptotics without condition (0.6) replaced by the non-degeneracy assumption (14.1.1) i.e.* $|F| + |\nabla F| \asymp 1$*:*

(i) Use the simple rescaling technique $\mu \mapsto \mu\rho^2$, $h \mapsto h\rho^{-1}$ *with the scaling function* $\rho(x) = \max(\epsilon_0|F|, \mu^{-\frac{1}{2}})$*;*

(ii) Improve what follows from the simple rescaling technique.

Problem 2.8.3. *Derive asymptotics under non-degeneracy assumption (0.8) rather than (0.7). The results of this section provide them at point* x *where* $|\nabla V/F| \gg \bar{\gamma} := \max(\mu^{-1}, \mu^{-\frac{1}{2}}h^{\frac{1}{2}})$.

(i) Can one improve those results using that under condition $(0.8)^+$ *and* $(0.8)^-$ *(when (0.8) holds but* $(0.8)^+$ *does not) the drift dynamic is elliptic elliptic and hyperbolic respectively?*

(ii) Can one improve a trivial estimate $O(\mu h^{-1})$ *at point* x *where* $|\nabla V/F| \lesssim \bar{\gamma}$*? Will be results affected by the difference between cases of* $(0.8)^+$ *and* $(0.8)^-$*?*

3 Dirac energy: $2\mathsf{D}$-estimates

In this and the next sections we consider asymptotics of expression I defined by (0.2).

3.1 Tauberian formula

Let us consider first contribution of zone $\{|x-y|\ge C\gamma\}$.

Proposition 3.1.1. *Under conditions* (0.5)–(0.7) *the contribution of zone* $\{|x-y|\ge C\gamma\}$*, to the remainder is* $O(\mu^{-1}h^{-1}\gamma^{-\kappa})$ *while the main part is given by the Tauberian approximation* I^{T}*, i.e. by the same expression* (0.1) *with* $e(x,y,0)$ *replaced by its standard implicit Tauberian approximation with* $T\asymp\epsilon\mu$ (0.11).

Proof. Recall that $e^{ih^{-1}tA}$ is the propagator of A and $U(x,y,t)$ is its Schwartz' kernel.

Consider expression (0.1) with $\omega(x,y)$ replaced by $\omega_\gamma(x,y)$ which is a cut-off of $\omega(x,y)$ in the zone $\{|x-y|\asymp\gamma\}$ and with the original functions ψ_1,ψ_2 replaced by 1. Let us replace *one* copy of $e(x,y,\tau)$ by $e(x,y,\tau,\tau')=\bigl(e(x,y,\tau)-e(x,y,\tau')\bigr)$ with $\tau'\le\tau$ and the second copy by $e(x,y,\tau'')$ and denote the resulting expression by $\mathsf{I}_\gamma(\tau,\tau',\tau'')$.

Now let us use decomposition

$$\omega_\gamma(x,y)=\gamma^{-d-\kappa}\int\psi_{1,\gamma}(x,z)\psi_{2,\gamma}(y,z)\,dz \tag{3.1}$$

with some $\psi_1,\psi_2\in\mathscr{C}_0^\infty$; as before subscript γ means rescaling.

Then $\mathsf{I}_\gamma(\tau,\tau',\tau'')$ does not exceed

$$\sum_j C\gamma^{-\kappa}\|\varphi_jE(\tau,\tau')\varphi_j\|_1 \tag{3.2}$$

where $E(\tau,\tau')=E(\tau)-E(\tau')$, φ_j are real-valued γ-admissible functions supported in $C_0\gamma$-vicinities of z_j and balls $B(z_j,2C_0\gamma)$ cover domain X with the multiplicity not exceeding C_0. Here we used that $\|E(\tau'')\|=1$. Since $E(\tau,\tau')$ is a positive operator and $\varphi_j=\varphi_j^*$, one can replace the trace norm by the trace itself and get

$$\sum_j C\gamma^{-\kappa}\operatorname{Tr}\varphi_jE(\tau,\tau')\varphi_j=C\gamma^{-\kappa}\operatorname{Tr}E(\tau,\tau')\bar{\psi} \tag{3.3}$$

with $\bar{\psi}=\sum_j \varphi_j^2$.

Further, we know from the standard theory of Chapter 13 that under conditions (0.5)–(0.7)

$$\|E(\tau,\tau')\bar{\psi}\|_1 \le Ch^{-2}\bigl(|\tau-\tau'|+CT^{-1}h\bigr)\qquad \forall \tau,\tau'\in[-\epsilon,\epsilon],\ \ T=\epsilon\mu \tag{3.4}$$

and therefore

$$|\mathsf{I}_\gamma(\tau,\tau',\tau'')|\le C\gamma^{-\kappa}h^{-2}\bigl(|\tau-\tau'|+CT^{-1}h\bigr) \tag{3.5}$$

in the same framework and therefore due to the standard Tauberian arguments we conclude that the contribution of zone $\{|x-y|\asymp\gamma\}$ to the Tauberian remainder estimate does not exceed $C\mu^{-1}h^{-1}\gamma^{-\kappa}$ which implies the statement immediately. □

However we need to consider also zone $\{|x-y|\le C\gamma\}$, complementary to one above. Assume that

$$\Omega_\kappa(z)=\sum_j D_{z_j}\Omega_{\kappa-1,j}+\Omega_{\kappa-1,0} \tag{3.6}$$

with the first subscript at Ω showing the degree of the singularity. Then

$$\omega_\kappa(x,x-y)\psi_\gamma(x-y)=\sum_j D_{x_j}\bigl(\omega_{\kappa-1,j}\psi_\gamma\bigr)+\omega_\kappa\psi'_\gamma+\omega_{\kappa-1}\psi''_\gamma \tag{3.7}$$

where $\psi_\gamma=\psi((x-y)\gamma^{-1})$ with ψ supported in $B(0,1)$ and equal 1 in $B(0,\frac{1}{2})$ while ψ'_γ is defined similarly with ψ' supported in $B(0,1)\setminus B(0,\frac{1}{2})$ and the last term gains 1 in the regularity.

After integration by parts expression $\mathsf{I}_{\kappa,\gamma}$, defined by (0.10) with Ω replaced by $\Omega\psi_\gamma$, becomes

$$-h^{-1}\sum_j\iint \omega_{\kappa-1,j}(x,y)(hD_{x_j})\bigl(e(x,y,\tau)\cdot e(y,x,\tau)\bigr)\,dxdy \tag{3.8}$$

plus two other terms: the term defined by (0.1) with the kernel $\Omega'_{\kappa,j}$ of the same singularity κ, albeit without factor h^{-1} and supported in the zone $\{|x-y|\ge\frac{1}{2}\gamma\}$, and the term defined by (0.1) with the kernel $\Omega'_{\kappa-1,j}$, also without factor h^{-1} and of singularity $(\kappa-1)$.

The former term could be considered as before yielding to the same remainder estimate $O(\mu^{-1}h^{1-d}\gamma^{-\kappa})$. To the latter term we can apply the same trick again and again raising power (and these terms are treated in the same manner (but simpler) as we deal below with (3.8).

So, one needs to consider (3.8) and thus, denoting the second copy of $e(y,x,\tau)$ by $f(y,x,\tau)$ and without using that they are equal

$$\begin{aligned}(hD_{x_j})\bigl(e(x,y,\tau)\cdot f(y,x,\tau)\bigr) &=\\ \bigl(hD_{x_j}e(x,y,\tau)\bigr)f(y,x,\tau) \ -\ e(x,y,\tau)\bigl(f(y,x,\tau)\,{}^t\!(hD_{x_j})\bigr) &=\\ \bigl(P_{j,x}e(x,y,\tau)\bigr)f(y,x,\tau) \ -\ e(x,y,\tau)\bigl(f(y,x,\tau)\,{}^tP_{j,x}\bigr).&\end{aligned}\tag{3.9}$$

Recall that $P_j=hD_j-\mu V_j(x)$ and ${}^tP_j=-hD_j-\mu V_j(x)$ is the dual operator. Also recall that if $e(x,y,\tau)$ and $f(y,x,\tau)$ are Schwartz kernels of $E(\tau)$ and $F(\tau)$, then $P_{j,x}e(x,y,\tau)$ and $f(y,x,\tau)\,{}^tP_{j,x}$ are those of $P_jE(\tau)$ and $F(\tau)P_j$.

Therefore we are interested in the expressions of the type

$$h^{-1}\iint \omega_{\kappa-1}(x,y)e(x,y,\tau)f(x,y,\tau)\psi_\gamma\, dxdy.\tag{3.10}$$

If $\kappa\le 1$ then replacing $e(x,y,\tau)$ and $f(y,x,\tau)$ by their standard Tauberian expressions one gets an error not exceeding $Ch^{-1}\times\mu^{-1}h^{-2}\gamma^{1-\kappa}$ because $\|P_jE(\tau)\|\le C_0$, $\|P_jF(\tau)\|\le C_0$ where $F(\tau)$ is an operator with the Schwartz kernel $f(x,y,\tau)$ and also because

$$\begin{aligned}\sum_j \|\varphi_jP_jE(\tau,\tau')\varphi_j\|_1 \le \sum_j \|\varphi_jP_jE(\tau,\tau')\|_2\cdot\|E(\tau,\tau')\varphi_j\|_2 \le\\ \sum_j \|\varphi_jP_jE(\tau,\tau')\|_2^2+\sum_j\|E(\tau,\tau')\varphi_j\|_2^2 =\\ \sum_j \operatorname{Tr}\varphi_jP_jE(\tau,\tau')P_j^*\varphi_j+\sum_j\operatorname{Tr}\varphi_jE(\tau,\tau')\varphi_j\\ \le Ch^{-2}\bigl(|\tau-\tau'|+C\mu^{-1}h\bigr)\qquad \forall \tau,\tau'\in[-\epsilon,\epsilon]\end{aligned}\tag{3.11}$$

which also easily follows from Chapter 13.

So, in this case one gets remainder estimate $O\bigl(\mu^{-1}h^{-1}\gamma^{-\kappa}+\mu^{-1}h^{-2}\gamma^{1-\kappa}\bigr)$ which is optimized to $O(\mu^{-1}h^{-1-\kappa})$ as $\gamma\asymp h$.

On the other hand, as $1<\kappa<2$ one can apply the same trick again since we did not use the fact that $e(.,.,.)$ and $f(.,.,.)$ coincide; then we arrive to the same estimates with P_j replaced by P_jP_k or even by $P^J := P_{j_1}P_{j_2}\cdots P_{j_l}$:

$$\operatorname{Tr} P^J E(\tau,\tau')(P^J)^* \le Ch^{-2}(|\tau-\tau'| + \mu^{-1}h) \qquad \forall \tau,\tau' \in [-\epsilon,\epsilon]. \tag{3.12}$$

Finally,

Remark 3.1.2. note that as $\kappa \ne 1$, decomposition (3.6) is always possible. Further, as $\kappa = 1$ this decomposition is possible as well provided one adds term $\varkappa(x)|x-y|^{-1}$ with an appropriate coefficient. On the other hand, if $\kappa = 1$ and $\omega(x,y) = \varkappa(x)|x-y|^{-1}$ then this decomposition is also possible but with $\omega_{0,j}(x,y) = \varkappa(x)(x_j-y_j)|x-y|^{-1}\log|x-y|$.

So we arrive to

Proposition 3.1.3. *Let conditions* (0.5)–(0.7) *be fulfilled. Then*

(i) As $0 < \kappa < 2$ and *either* *$\kappa \ne 1$* *or* *$\kappa = 1$ and $\omega(x,y)$ is replaced by $\omega(x,y) - \varkappa(\frac{1}{2}(x+y))|x-y|^{-1}$ with an appropriate smooth coefficient $\varkappa(x)$, with the error $O(\mu^{-1}h^{-1-\kappa})$ one can replace $e(x,y,\tau)$ by its standard Tauberian expression* (0.11) *in the formula* (0.11) *for* I.

(ii) As $\kappa = 1$ and $\omega = \varkappa(\frac{1}{2}(x+y))|x-y|^{-1}$, with the error $O(\mu^{-1}h^{-1-\kappa}|\log h|)$ one can replace $e(x,y,\tau)$ by its standard Tauberian expression (0.11) *in the formula* (0.11) *for* I.

Remark 3.1.4. (i) The arguments above show that in an appropriate sense one can consider arbitrary $\kappa \in \mathbb{R}$ and even in $\mathbb{C}$.

(ii) One needs only (3.12) rather than (0.5)–(0.7), and (3.12) holds as (0.7) is replaced by a weaker non-degeneracy condition $(0.8)^+$;

(iii) Furthermore (3.12) with an extra factor $(1+\mu h|\log h|)$ in the right-hand expression holds under condition (0.8).

Problem 3.1.5. *Can one prove the similar result for I_m defined by (1.0.6) V. Ivrii [Ivr3] with $m \ge 3$?*

Thus we arrive to

Proposition 3.1.6. *Let conditions* (0.5) *and* (0.6) *be fulfilled.*

(i) Further, let *either* *condition* $(0.8)^+$ *be fulfilled* *or* *condition* (0.8) *be fulfilled and $\mu \le h^{-1}|\log h|^{-1}$. Then* (0.11) *and statements (i), (ii) of proposition 3.1.3 hold.*

(ii) On the other hand, let condition (0.8) *be fulfilled and* $h^{-1}|\log h|^{-1}\le \mu\le h^{-1}$*. Then* (0.11) *and statements (i), (ii) of proposition 3.1.3 hold with an extra factor* $(1+\mu h|\log h|)$ *in the right-hand expressions.*

Remark 3.1.7. Under certain assumptions (see [IRO4]) this result could be generalized for $d\ge 4$. However in calculations we use that $d=2$.

3.2 Superstrong magnetic field

Consider Schrödinger-Pauli operator as $\mu h\ge \epsilon_0$. Then we arrive immediately

Proposition 3.2.1. *Let* $\mu h\ge\epsilon_0$*. Then*

(i) Contribution of $\{|x-y|\ge\gamma\}$ *to* I *does not exceed* $C\mu h^{-1}\gamma^{-\kappa}$*;*

(ii) Further,

$$|\mathsf{I}|\le C\mu^{1+\frac{1}{2}\kappa}h^{-1-\frac{1}{2}\kappa}. \tag{3.13}$$

(iii) Furthermore, $\mathsf{I}=O(\mu^{-\infty})$ *provided*

$$V+\mu hF\ge\epsilon\qquad \forall x\in B(0,1); \tag{3.14}$$

in particular it is the case as $\mathfrak{z}<1$ *and* $\mu h\ge C_0$*.*

Proof. Statement (i) trivially follows from the fact that $\mathscr{L}^2$ norm of $e(.,,,\tau)$ does not exceed $C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}$.

Meanwhile contribution of $\{(x,y): x\in B(0,1), y\in B(0,1), |x-y|\le\gamma\}$ does not exceed $C\mu^2h^{-2}\gamma^{2-\kappa}$. Adding to (i) we conclude that the sum reaches its minimal $C\mu^{1+\frac{1}{2}\kappa}h^{-1-\frac{1}{2}\kappa}$ value as $\gamma=\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$.

Finally, statement (iii) is due to the fact that $e(x,y,0)=O(\mu^{-\infty})$ as condition (3.14) is fulfilled in one of points x,y; see subsection 13.5.1. ☐

Proposition 3.2.2. *Let conditions* (0.1)–(0.5) *be fulfilled. Let* $\mu h\ge\epsilon$ *and one of the nondegeneracy conditions*

$$|\frac{V}{F}+(2m+1)\mu h|+|\nabla\frac{V}{F}|\ge\epsilon \tag{3.15}$$

or

$$|\frac{V}{F}+(2m+1)\mu h|+|\nabla\frac{V}{F}|\le\epsilon\implies \operatorname{det}\operatorname{Hess}\frac{V}{F}\ge\epsilon \tag*{(3.17)$^+$}$$

be fulfilled.

(i) As $\gamma \ge C_0\mu^{-\frac{1}{2}}h^{-\frac{1}{2}}$ *contribution of* $\{|x-y|\ge\gamma\}$ *to* I *does not exceed* $C\gamma^{-2-\kappa}$*;*

(ii) Estimate

$$|\mathsf{I}-\mathsf{I}^{\mathsf{T}}| \le C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa} \tag{3.16}$$

holds as $\kappa\ne 1$*; as* $\kappa=1$ *this estimate holds with an extra factor* $|\log\mu|$ *in its right-hand expression;*

(iii) If instead of (3.15) *or* $(3.17)^+$ *we assume that*

$$|\frac{V}{F}+(2m+1)\mu h|+|\nabla\frac{V}{F}|\le\epsilon \implies |\det \operatorname{Hess}\frac{V}{F}\ge\epsilon| \tag{3.17}$$

then (i), (ii) hold with an extra factor $|\log\mu|$ *in the right-hand expressions;*

(iv) $\mathsf{I}-\mathsf{I}^{\mathsf{T}}=O(\mu^{-\infty})$ *under ellipticity condition* $|V+(2m+1)\mu hF|\ge\epsilon$*. In particular, it happens as* $1<\mathfrak{z}\notin 2\mathbb{Z}+1$ *and* $\mu h\ge C_0$*.*

Proof. To prove statement (i) recall that the drift speed does not exceed $C\mu^{-1}$ and therefore Hilbert-Schmidt norm of $\psi E(\tau)\psi'$ does not exceed C as ψ,ψ' are $\mathscr{C}_0^\infty$-functions with $\operatorname{dist}(\operatorname{supp}\psi,\operatorname{supp}\psi')\asymp 1$.

Really, it is true for a Hilbert-Schmidt norm of $\bigl(E(\tau)-E(\tau')\bigr)\psi$ with $|\tau-\tau'|\le\mu^{-1}h$ and then by Tauberian theorem it is true for a Hilbert-Schmidt norm of $\bigl(E(\tau)-E^{\mathsf{T}}(\tau)\bigr)\psi'$ with E^{T} operator with the Schwartz kernel e^{T} with time $T\asymp\mu$. However $\psi E^{\mathsf{T}}\psi'$ is negligible as $T\le\epsilon\mu$ due to propagation results.

Then rescaling $x\mapsto x/\gamma$, $y\mapsto y/\gamma$, $h\mapsto h/\gamma$, $\mu\mapsto\mu\gamma$ we conclude that for γ-admissible ψ,ψ' with $\operatorname{dist}(\operatorname{supp}\psi,\operatorname{supp}\psi')\asymp 1$ Hilbert-Schmidt norm of $\psi E(\tau)\psi'$ does not exceed C and therefore the contribution of

$$K(z)=\{(x,y): |x-z|\le 2\gamma, |y-z|\le 2\gamma, |x-y|\ge\gamma\}$$

does not exceed $\gamma^{-\kappa}$; therefore taking partition we conclude that the contribution of zone $\{(x,y)\in B(0,1)\times B(0,1), |x-y|\ge\gamma\}$ does not exceed $C\gamma^{-2-\kappa}$.

Statement (iv) follows from the fact $e-e^{\mathsf{T}}=O(\mu^{-\infty})$ under ellipticity condition.

To prove statement (ii) we apply the same arguments as in proposition 3.1.3. While (3.1)–(3.3), (3.6)–(3.10) remain true, (3.4), (3.5) should be modified: factor h^{-2} must be replaced by μh^{-1}.

We claim that

(3.18) As $0<\kappa<1$ (3.11) holds with factor h^{-2} replaced by $\mu^{\frac{3}{2}}h^{-\frac{1}{2}}$.

Really, the same arguments are applied as before albeit now $P_jE(\tau)$ is bounded by $C_0(\mu h)^{\frac{1}{2}}$ rather than by C_0.

Then we arrive to the remainder estimate $O\bigl(\gamma^{-\kappa}+\mu^{\frac{1}{2}}h^{-\frac{3}{2}}\gamma^{1-\kappa}\bigr)$ and optimizing by $\gamma=\bar{\gamma}:=C\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ we arrive to the remainder estimate $O(\bar{\gamma}^{-\kappa})$.

As $\kappa=1$ we get the same estimate albeit with a logarithmic factor.

To tackle the case $1<\kappa<2$ one should replace (3.12) by

$$\operatorname{Tr} P^JE(\tau,\tau')(P^J)^*\le C\mu h^{-1}(|\tau-\tau'|+\mu^{-1}h)(\mu h)^{|J|}\qquad \forall \tau,\tau'\in[-\epsilon,\epsilon] \tag{3.12$'$}$$

as $\mu h\ge 1$; one can prove it easily as $\|P^JE(\tau)\|\le C(1+\mu h)^{\frac{1}{2}|J|}$.

Then as $1<\kappa<2$ we arrive to the remainder estimate $O(\gamma^{-\kappa}+\mu h^{-1}\gamma^{2-\kappa})$ and optimizing by $\gamma=\bar{\gamma}$ we again arrive to the remainder estimate $O(\bar{\gamma}^{-\kappa})$.

Finally, the above arguments imply (iii). □

4 Dirac energy: 2D-calculations

4.1 Pilot Model

4.1.1 Transformations

Consider pilot-model operator (1.1). Let us rescale as before[10]. Then as $U(x,y,t)$ is defined by (1.9)–(1.10), and $e(x,y,\tau)$ is given by (1.13), and † means a complex conjugation we get

---

[10] We need to add factor $\mu^{-4+\kappa}=\mu^{-2}\times\mu^{-2}\times\mu^{-\kappa}$ where two factors μ^{-2} are coming from dx and dy and μ^{κ} from ω.

$$
\begin{aligned}
(4.1)\quad \mathsf{I} := (2\pi)^{-2}\mu^{-4+\kappa} \iint\iint \omega(\mu^{-1}x, \mu^{-1}y)(t't'')^{-1}\times \\
e^{-i\hbar^{-1}(t'-t'')\tau} U(x,y,t')U^{\dagger}(x,y,t'')\, dt'dt''\, dxdy = \\
\frac{1}{4}(2\pi)^{-4}\hbar^{-2}\mu^{\kappa} \iint\iint \omega(x,y)\csc(t')\csc(t'')(t't'')^{-1}\times \\
\exp\Bigl(i\hbar^{-1}\bigl(\bar\phi(x,y,t') - \bar\phi(x,y,t'') - (t'-t'')\tau\bigr)\Bigr)\, dt'dt''\, dxdy
\end{aligned}
$$

with

$$
\begin{aligned}
(4.2)\quad \bar\phi(x,y,t') - \bar\phi(x,y,t'') - (t'-t'')\tau = \\
-\frac{1}{4}\bigl(\cot(t') - \cot(t'')\bigr)(x_1-y_1)^2 \\
-\frac{1}{4}\cot(t')(x_2-y_2+2t'\varepsilon)^2 + \frac{1}{4}\cot(t'')(x_2-y_2+2t''\varepsilon)^2 \\
+(x_1+y_1+2\varepsilon)(t'-t'')\varepsilon - (t'-t'')(\tau+\varepsilon^2).
\end{aligned}
$$

Recall that in the original coordinates $\omega(x,y) = \Omega\bigl(\frac{1}{2}(x+y); x-y\bigr)$ where $\Omega(.,.)$ is uniformly smooth with respect to the first variable and positively homogeneous of degree $-\kappa$ with respect to the second one as $|x-y|\le 1$. Therefore without any loss of the generality one can assume that

$$
\omega(x,y) = \Omega\bigl(\frac{1}{2}(x+y); x-y\bigr)\psi(x-y) \tag{4.3}
$$

where now Ω is positively homogeneous of degree $-\kappa$ and $\psi\in\mathscr{C}_0^\infty\bigl(B(0,1)\bigr)$ equal 1 in $B(0,\frac{1}{2})$; the difference would be of the same nature but with κ replaced by 0.

We can replace variables x, y with new variables $\mathsf{x} := \frac{1}{2}(x+y)$ and $z := (x-y)$ and rescale. Note that (4.2) depends on x in a very specific way: it does not depend on x_2 at all and it is linear with respect to x_1; thus

(4.1) becomes[11)]

$$\frac{1}{4}(2\pi)^{-3}\hbar^{-2}\mu^{\kappa+2}\iint\iint \hat{\Omega}\bigl(2h^{-1}\varepsilon(t'-t''),z\bigr)\csc(t')\csc(t'')(t't'')^{-1}\times\\
\exp\Bigl(i\hbar^{-1}\Bigl(-\frac{1}{4}\bigl(\cot(t')-\cot(t'')\bigr)z_1^2-\frac{1}{4}\cot(t')(z_2+2t'\alpha\mu^{-1})^2+\\
\frac{1}{4}\cot(t'')(z_2+2t''\varepsilon)^2-(t'-t'')(\tau-\varepsilon^2)\Bigr)\Bigr)\,dt'dt''\,\psi(\mu^{-1}z)\,dz_1dz_2$$

where $\hat{\Omega}(\lambda;z)=F_{x_1\to\lambda}\Omega$ is a partial Fourier transform. Rewriting the third line in (4.2) as

$$-\frac{1}{4}\bigl(\cot(t)-\cot(t')\bigr)(x_2-y_2+(t+t')\varepsilon)^2\\
-\frac{1}{4}\bigl(\cot(t)-\cot(t')\bigr)(t-t')^2\varepsilon^2-\\
\frac{1}{2}\bigl(\cot(t)+\cot(t')\bigr)\bigl(x_2-y_2+(t+t')\varepsilon\bigr)(t-t')\varepsilon$$

and replacing $t'=t+s$, $t''=t-s$ we arrive to

$$\mathsf{I}=\tag{4.4}\\
\frac{1}{2}(2\pi)^{-3}\hbar^{-2}\mu^{\kappa+2}\iint\iint\hat{\Omega}(4h^{-1}\varepsilon s,z)\csc(t+s)\csc(t-s)(t+s)^{-1}(t-s)^{-1}\times\\
\exp\Bigl(i\hbar^{-1}\Bigl[-\frac{1}{4}\bigl(\cot(t+s)-\cot(t-s)\bigr)\bigl(z_1^2+(z_2+2t\varepsilon)^2+4s^2\varepsilon^2\bigr)\\
-\bigl(\cot(t+s)+\cot(t-s)\bigr)s\varepsilon-2s(\tau-\varepsilon^2)\Bigr]\Bigr)\,dtds\,\psi(\mu^{-1}z)dz_1dz_2.$$

Remark 4.1.1. (i) For a sake of simplicity we assume (without any loss of the generalization) that $\alpha>0$.

(ii) If $\alpha\ll 1$ in the above analysis applied to the general operators we need to take α-admissible with respect to x function Ω_α instead of Ω; thus up to the shift we replace $\Omega(\mathsf{x},\cdot)$ by $\Omega(\mathsf{x}\alpha^{-1},\cdot)$ and thus $\hat{\Omega}(2h^{-1}\varepsilon s,\cdot)$ by $\alpha^2\hat{\Omega}(2h^{-1}\varepsilon\alpha s,\cdot)$. In this case we will use notation I_α instead of I.

In the same time we need to consider separately $|z|\le\mu\alpha$ and $|z|\ge\mu\alpha$. For a while we will not assume that $\varepsilon\asymp\mu^{-1}\alpha$.

[11)] In these calculations we skip x_2 as an argument of Ω and integration by $d\mu^{-1}\mathsf{x}_2$

4.1.2 Case $\alpha^2 \gg \mu h$

Assume first that

$$\alpha^2 \ge \mu h^{1-\delta} \tag{4.5}$$

where for a sake of simplicity we assume that $\alpha > 0$ (and therefore $\varepsilon > 0$).

Remark 4.1.2. (i) In the virtue of the factor $\hat{\Omega}(2\hbar^{-1}\alpha^2 s, \cdot)$ under assumption (4.5) we need to consider only $|s| \le h^{\delta}$ and therefore we can consider separately $|t'| \le \epsilon_0, |t''| \le \epsilon_0$ and $|t'| \ge \epsilon_0, |t''| \ge \epsilon_0$;

(ii) Note that due to section 6.3 contribution of zone $\{|t'| \le \epsilon_0, |t''| \le \epsilon_0\}$ defined by integral expressions (4.1) or (4.4) with an extra factor $\bar{\chi}_{\epsilon_0}(t')$ or $\bar{\chi}_{\epsilon_0}(t'')$ or $\bar{\chi}_{\epsilon_0}(t)$ differs from the same expression for non-magnetic Schrödinger operator by $O(\mu h^{-1-\kappa} \times \hbar^{\kappa}) = O(\mu^{\kappa+1} h^{-1})$ as $\kappa \ne 1$ and by $O(\mu^2 h^{-1} |\log \mu|)$ as $\kappa = 1$.

(iii) Furthermore, if we remove from this expression for a non-magnetic Schrödinger operator cut-off $\{|t'| \ge \epsilon_0\}$ then the error would not exceed the same expression as well.

Let us consider contribution of zone $\{|t'| \ge \epsilon_0, |t''| \ge \epsilon_0\}$ defined by an integral expressions (4.1) or (4.4) with an extra factor $\bigl(1 - \bar{\chi}_{\epsilon_0}(t')\bigr)$. Due to remark 4.1.2(i) we need to consider only t', t'' belonging to the same tick.

Let us consider first zone

$$\{|s| \ge \hbar/\alpha^2, |\sin(t)| \ge C|s|\}. \tag{4.6}$$

Then $\cot(t') - \cot(t'') \asymp -\sin^{-2}(t)s$ and integration by parts with respect to z delivers one of the factors

$$|s|^{-1}\hbar\bigl(|z_1|^2 + |z_2| \cdot |z_2 + \varepsilon t|\bigr)^{-1} |\sin(t)|^2, \tag{4.7}$$

$$|s|^{-1}\hbar\bigl(|z_1|^2 + |z_2 + \varepsilon t|^2\bigr)^{-1} |\sin(t)|^2. \tag{4.8}$$

Thus integrating by parts many times in the zone where both of these factors are less than 1 we acquire factors

$$\Bigl(1 + |s|\hbar^{-1}\bigl(|z_1|^2 + |z_2| \cdot |z_2 + \varepsilon t|\bigr) |\sin(t)|^{-2}\Bigr)^{-l}, \tag{4.9}$$

$$\Bigl(1 + |s|\hbar^{-1}\bigl(|z_1|^2 + |z_2 + \varepsilon t|^2\bigr) |\sin(t)|^{-2}\Bigr)^{-l} \tag{4.10}$$

respectively. Multiplying by $|z|^{-\kappa}$ and integrating we get after multiplication by $|\sin(t)|^{-2}$

$$\asymp |\sin(t)|^{-\kappa}|s|^{-1+\frac{1}{2}\kappa}\hbar^{1-\frac{1}{2}\kappa} \tag{4.11}$$

and integrating by t over one tick intersected with $\{t:\ |\sin(t)|\ge |s|\}$ we get

$$C\hbar^{1-\frac{1}{2}\kappa}\begin{cases} |s|^{-1+\frac{1}{2}\kappa} & \kappa<1,\\ |s|^{-\frac{1}{2}\kappa} & \kappa>1,\\ |s|^{-\frac{1}{2}}(1+|\log|s||) & \kappa=1.\end{cases} \tag{4.12}$$

This expression (4.12) must be either integrated by with respect to s: $|s|\le \hbar/\alpha^2$ or multiplied by $(\hbar/\alpha^2)^l|s|^{-l}$ due to factor $\hat{\Omega}$ and integrated over $|s|\ge \hbar/\alpha^2$, resulting in both cases in the same answer which is the value of (4.12) $\times |s|$ calculated as $s=\hbar/\alpha^2$ i.e.

$$C\hbar^{1-\frac{1}{2}\kappa}\begin{cases} (\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1.\end{cases} \tag{4.13}$$

In addition to zone (4.6) we need to consider zone

$$\{|\sin(t')|\asymp |s|,\ |\sin(t'')|\le |s|\}; \tag{4.14}$$

its tween $\{|\sin(t')|\asymp |s|,\ |\sin(t'')|\le |s|\}$ is considered in the same way.

In zone (4.14) $|\cot(t')-\cot(t'')|\asymp |\sin(t'')|^{-1}$ and in this case (4.7), (4.8) are replaced by

$$\hbar\bigl(|z_1|^2+|z_2|\cdot|z_2+\varepsilon t|\bigr)^{-1}|\sin(t'')|, \tag*{(4.7)$'$}$$

$$\hbar\bigl(|z_1|^2+|z_2+\varepsilon t|^2\bigr)^{-1}|\sin(t'')| \tag*{(4.8)$'$}$$

and (4.9), (4.10) by

$$\Bigl(1+\hbar^{-1}\bigl(|z_1|^2+|z_2|\cdot|z_2+\varepsilon t|\bigr)|\sin(t'')|^{-1}\Bigr)^{-l}, \tag*{(4.9)$'$}$$

$$\Bigl(1+\hbar^{-1}\bigl(|z_1|^2+|z_2+\varepsilon t|^2\bigr)|\sin(t'')|^{-1}\Bigr)^{-l}. \tag*{(4.10)$'$}$$

Then, multiplying by $|z|^{-\kappa}$ and integrating we get after multiplication by $|\sin(t'')|^{-1}|\sin(t')|^{-1}$

$$\hbar^{1-\frac{1}{2}\kappa}|\sin(t'')|^{-\frac{1}{2}\kappa}|s|^{-1}; \tag{4.11$'$}$$

then integrating by $|t''|$ over one tick but intersected with $\{|\sin(t'')|\le |s|\}$ we get

$$\hbar^{1-\frac{1}{2}\kappa}|s|^{-\frac{1}{2}\kappa}. \tag{4.12$'$}$$

Finally, either integrating over $|s|\le \hbar/\alpha^2$ or multiplying by $|s|^{-l}(\hbar/\alpha^2)^l$ and integrating over $|s|\ge \hbar/\alpha^2$ we get in both cases the same answer $\hbar^{1-\frac{1}{2}\kappa}(\hbar\alpha^2)^{1-\frac{1}{2}\kappa}$ not exceeding (4.13).

Therefore the total contribution of zones (4.6) and (4.14) is given by expression (4.13). Then multiplying by $|k|^{-2}\mu^\kappa h^{-2}\alpha^2$ we get after summation with respect to $k: |k|\ge 1$ the value as $k=1$ i.e. (4.13) × $\mu^\kappa h^{-2}\alpha^2$.

Therefore we arrive to

Proposition 4.1.3. *For the pilot-model operator with* $1\le\mu\le h^{-1}$ *and* $\alpha\ge\epsilon_0\mu^{-1}$ *under additional assumption* (4.5)

$$|\mathsf{I}^{\mathsf{T}}_\alpha-\mathsf{I}^{\mathsf{T}}_\alpha{}'|\le CR^{\mathsf{W}}(\alpha):=\\ C\left\{\begin{aligned} &\mu^{1+\kappa}h^{-1}\alpha^{2-\kappa} && 0<\kappa<1,\\ &\mu^2h^{-\kappa}\alpha^\kappa && 1<\kappa<2,\\ &\mu^2h^{-1}\alpha(1+|\log(\mu h/\alpha^2)|) && \kappa=1\end{aligned}\right. \tag{4.15}$$

where $\mathsf{I}^{\mathsf{T}}_\alpha{}'$ *is a Tauberian expression for* I *albeit with* $T=\epsilon_0\mu^{-1}$.

Now we need to explore the difference between $\mathsf{I}^{\mathsf{T}}_\alpha{}'$ and $\mathcal{I}^{\mathsf{W}}_\alpha$ defined by (0.9).

Proposition 4.1.4. *For the pilot-model operator with* $1\le\mu\le h^{-1}$ *and* $\alpha\ge\epsilon_0\mu^{-1}$

$$|\mathsf{I}^{\mathsf{T}}_\alpha{}'-\mathcal{I}^{\mathsf{W}}_\alpha|\le C\mu^2h^{-\kappa}\alpha^2. \tag{4.16}$$

Proof. Repeating arguments of subsection 6.3.4 one can prove easily that

$$|\mathsf{I}^{\mathsf{T}}_\alpha{}'-\sum_{l:0\le l\le L-1,j=0,1}\mathcal{I}^{\mathsf{W}}_{(l)j,\alpha}\mu^l|\le C(\mu h)^Lh^{-2-\kappa}\alpha^2+C\mu^{\kappa+1}h^{-1}\alpha^2 \tag{4.17}$$

where $\mathcal{I}^{\mathsf{W}}_{(l)j,\mu^{-1}}$ is defined by (0.9) albeit with $\Omega(\frac{1}{2}(x+y), x-y)$ multiplied by a homogeneous polynomial of degree $(l+j)$ with respect to $(x-y)$ and for $l=j=0$ this polynomial is 1 and with integrals taken $\underline{\text{only}}$ over zone $\{(x,) : |x-y| \lesssim \mu^{-1}\}$.

Since without any loss of the generality we can assume that Ω is even with respect to the second argument, all terms with odd $(l+j)$ vanish and since replacing μ by $-\mu$ we should arrive to the same result, all terms with odd l vanish and therefore

(4.18) In (4.17) all terms vanish except those with $j=0$ and even l.

Picking $L=2$ we arrive then to (4.16). □

Combining propositions 4.1.3 and 4.1.4 and noting that the right-hand expression of (4.16) does not exceed $CR^{\mathsf{W}}(\alpha)$ we arrive to

Corollary 4.1.5. *For the pilot-model operator with* $1 \le \mu \le h^{-1}$ *and* $\alpha \ge \epsilon_0\mu^{-1}$ *under additional assumption* (4.5)

$$|\mathsf{I}^{\mathsf{T}}_{\alpha} - \mathcal{I}^{\mathsf{W}}_{\alpha}| \le CR^{\mathsf{W}}(\alpha). \tag{4.19}$$

4.1.3 Improvement

In a certain case (under assumption (4.22) below) we can improve the results of the previous subsubsection 4.1.2. To do this we note that (an easy proof is left to the reader)

Remark 4.1.6. Let $\zeta \le \varepsilon|t|$ with

$$\zeta := (\hbar/|s|)^{\frac{1}{2}}|\sin(t)|. \tag{4.20}$$

Then the sum of (4.9) and (4.10), multiplied by $|z|^{-\kappa}\,dz$ and integrated, does not exceed

$$C\begin{cases} \zeta^2(\varepsilon|t|)^{-\kappa} & 0<\kappa<1,\\ \zeta^{4-2\kappa}(\varepsilon|t|)^{\kappa-2} & 1<\kappa<2,\\ \zeta^2(\varepsilon|t|)^{-1}(1+|\log(\zeta/\varepsilon|t|)) & \kappa=1. \end{cases} \tag{4.21}$$

We need to multiply by $|\sin(t)|^{-2}\,dt$ and integrate. Then as $0<\kappa<1$ main contribution came from $\sin(t)\asymp 1$ and improvement is possible iff then $\zeta\le\varepsilon$ (as $|k|=1$ provided the large part of contribution). But as $s\asymp\hbar/\alpha^2$, $\sin(t)\asymp 1$ (which was the main contributor in the previous subsubsection) $\zeta\asymp\alpha$ while $\varepsilon=\mu^{-1}\alpha$.

So, improvement is possible only for $1<\kappa<2$. In this case main contribution came from $\sin(t)\asymp s$ and improvement is possible if then $\zeta\asymp(\hbar|s|)^{\frac{1}{2}}\le\mu^{-1}\alpha$. Setting $|s|=\hbar/\alpha^2$ we have $\zeta\asymp\hbar/\alpha$ and therefore improvement is possible if

$$\alpha^2\ge\mu^2 h. \tag{4.22}$$

Consider zone (4.6) first. Then multiplication by $|\sin(t)|^{-2}\,dt$ and integration results instead of (4.12) in

$$C\begin{cases}\hbar^{\frac{1}{2}}|s|^{-\frac{1}{2}}\varepsilon^{1-\kappa} & 1<\kappa<\dfrac{3}{2},\\ \hbar^{2-\kappa}|s|^{1-\kappa}\varepsilon^{\kappa-2} & \dfrac{3}{2}<\kappa<2,\\ \hbar^{\frac{1}{2}}|s|^{-\frac{1}{2}}\varepsilon^{-\frac{1}{2}}(1+|\log(\hbar|s|/\varepsilon^2)) & \kappa=\dfrac{3}{2}\end{cases} \tag{4.23}$$

where we set $|k|=1$. Finally, either integrating over $|s|\le\hbar/\alpha^2$ or multiplying by $|s|^{-l}(\hbar/\alpha^2)^l$ and integrating over $|s|\ge\hbar/\alpha^2$ we get instead of (4.13)

$$C\begin{cases}\mu^{\kappa}h\alpha^{-\kappa} & 1<\kappa<\dfrac{3}{2},\\ \mu^{6-3\kappa}h^{4-2\kappa}\alpha^{3\kappa-6} & \dfrac{3}{2}<\kappa<2,\\ \mu^{\frac{3}{2}}h\alpha^{-\frac{3}{2}}(1+|\log(\mu^2h/\alpha^2)|) & \kappa=\dfrac{3}{2}.\end{cases} \tag{4.24}$$

Consider zone (4.14); we apply remark 4.1.6 albeit with

$$\zeta:=(\hbar|\sin(t'')|)^{\frac{1}{2}}. \tag*{(4.20)$'$}$$

Again multiplying by $|\sin(t')|^{-1}|\sin(t'')|^{-1}$ and integrating by $|t''|$ over one tick but intersected with $\{|\sin(t'')|\le|s|\}$ and setting $|\sin(t')|=|s|$, and finally either integrating over $|s|\le\hbar/\alpha^2$ or multiplying by $|s|^{-l}(\hbar/\alpha^2)^l$ and integrating over $|s|\ge \hbar/\alpha^2$, we arrive to the same answer, not exceeding (4.24).

Finally, multiplying by $|k|^{-2}\mu^\kappa h^{-2}\alpha^2$ we get after summation with respect to $k:|k|\ge 1$ the following improvement to proposition 4.1.3

Proposition 4.1.7. *For the pilot-model operator with* $1\le\mu\le h^{-1}$ *and* $\alpha\ge\max(\mu^{-1},\mu h^{\frac12},\mu^{\frac12}h^{\frac12-\delta})$

$$|\mathsf{I}^{\mathsf{T}}_\alpha-\mathsf{I}^{\mathsf{T}\prime}_\alpha|\le CR^{\mathsf{W}}_1(\alpha):= C\begin{cases}\mu^{2\kappa}h^{-1}\alpha^{2-\kappa} & 1<\kappa<\dfrac{3}{2},\\ \mu^{6-2\kappa}h^{2-2\kappa}\alpha^{3\kappa-4} & \dfrac{3}{2}<\kappa<2,\\ \mu^3h^{-1}\alpha^{\frac12}(1+|\log(\mu^2h/\alpha^2)|) & \kappa=\dfrac{3}{2}.\end{cases}\tag{4.25}$$

Remark 4.1.8. Right-hand expression of (4.25) in comparison with (4.15) has factor

$$C\begin{cases}(\mu^2h/\alpha^2)^{\kappa-1} & 1<\kappa<\dfrac{3}{2},\\ (\mu^2h/\alpha^2)^{2-\kappa} & \dfrac{3}{2}<\kappa<2,\\ (\mu^2h/\alpha^2)^{\frac12}(1+|\log(\mu^2h)|) & \kappa=\dfrac{3}{2}.\end{cases}\tag{4.26}$$

While we cannot directly apply estimate (4.16) now as $\mu^2h^{-\kappa}$ may be greater than $R^{\mathsf{W}}_1(\alpha)$, we can apply (4.17)–(4.18) with $L=4$ resulting in

Proposition 4.1.9. *For the pilot-model operator with* $1\le\mu\le h^{-1}$ *and* $\alpha\ge\max(\mu^{-1},\mu h^{\frac12},\mu^{\frac12}h^{\frac12-\delta})$

$$|\mathsf{I}^{\mathsf{T}}_\alpha-\mathcal{I}^{\mathsf{W}}_\alpha-\mathcal{I}^{\mathsf{W}}_{\mathsf{corr}}|\le CR^{\mathsf{W}}_1(\alpha)\tag{4.27}$$

with $\mathcal{I}^{\mathsf{W}}_{\mathsf{corr}}:=\mathcal{I}^{\mathsf{W}}_{(2)0,\alpha}$ *but taken* *only* *over zone* $\{|x-y|\le\mu^{-1}\}$.

4.1.4 Case $\alpha^2\not\gg\mu h$

Assume now that $\alpha\ge\mu^{-1}$ but (4.5) fails. Then in contrast to the previous we will need to compute contributions of pair of ticks (k',k'') with $k'\ne k''$. Then if t',t'' belong to k'-th and k''-th ticks respectively we denote $r=k'-k''$ and $s=t'-t''-2\pi r$.

Contribution of $k'=k''\ne 0$. Consider case $k'-k''=r=0$ first. As $\alpha^2\ge \hbar$ applying the same arguments as before we get the same answer (4.13) as before. Meanwhile, as $\alpha^2\le\hbar$ we need to integrate (4.12) over $\{s:|s|\le 1\}$ and we arrive to $\hbar^{1-\frac{1}{2}\kappa}$ for all $\kappa: 0<\kappa<2$.

Multiplication by $\mu^\kappa h^{-2}|k|^{-2}\alpha^2$ and summation with respect to k results in the right hand expression of (4.15) as $\alpha^2\ge\hbar$ and

$$C\mu^\kappa h^{-2}\hbar^{1-\frac{1}{2}\kappa}\alpha^2 = C\mu^{\frac{1}{2}\kappa+1}h^{-1-\frac{1}{2}\kappa}\alpha^2 \tag{4.28}$$

as $\alpha^2\le\hbar$.

Contribution of $0\ne k'\ne k''\ne 0$. Consider now case $k'-k''=r\ne 0$. As $\alpha^2\ge\hbar$ expression (4.13) acquires factor $\bigl(|k'-k''|\alpha^2/\hbar\bigr)^{-l}$ and multiplying by $\mu^\kappa h^{-2}\alpha^2|k'|^{-1}|k''|^{-1}$ we get after summation with respect to $k'\ne 0$, $k''\ne 0$ the same right-hand expression of (4.15) [12].

Meanwhile, as $\alpha^2\le\hbar$ instead of (4.13) we get

$$C\hbar^{1-\frac{1}{2}\kappa}\bigl(1+|k'-k''|\alpha^2/\hbar\bigr)^{-l}.$$

Multiplication by $\mu^\kappa h^{-2}\alpha^2|k'|^{-1}|k''|^{-1}$ and summation with with respect to k', k'' returns

$$C\mu^{1+\frac{1}{2}\kappa}h^{-1-\frac{1}{2}\kappa}\alpha^2\bigl(1+(\log(\hbar/\alpha^2))_+\bigr)^2. \tag{4.29}$$

Contribution of $0=k'\ne k''$. Consider now $k'=0, k''\ne 0$; its tween case $k'\ne 0, k''=0$ is addressed in the same way. To do this we need to modify our expression fo I: namely, we do not divide by t' and then take Fourier transform $F_{t'\to\hbar^{-1}\tau}$; instead we take Fourier transform $F_{t'\to\hbar^{-1}\tau'}$, then integrate by τ' to τ and divide by $\hbar$; modifying this way (4.1) we arrive instead of (4.4) to

$$\begin{multline}\mathsf{I}_\alpha = \\ \frac{1}{2}(2\pi)^{-3}\hbar^{-3}\mu^{\kappa+2}\alpha^2\int\iint\iint \hat{\Omega}(4\hbar^{-1}\alpha^2 s, z)\times \\ \csc(t+s)\csc(t-s)(t+s)^{-1}(t-s)^{-1}\times \\ \exp\Bigl(i\hbar^{-1}\Bigl[-\frac{1}{4}\bigl(\cot(t+s)-\cot(t-s)\bigr)\bigl(z_1^2+(z_2+2t\varepsilon)^2+4s^2\varepsilon^2\bigr) \\ -\bigl(\cot(t+s)+\cot(t-s)\bigr)s\varepsilon-2s(\tau-\varepsilon^2)+(t+s)(\tau-\tau')\Bigr]\Bigr)\,dtds\,\psi(\mu^{-1}z)dz_1dz_2\,d\tau'.\end{multline} \tag{4.30}$$

---

[12] Actually, with an extra factor $(\hbar/\alpha^2)^l$ but we do not need it here.

Then repeating our above arguments we conclude that

(4.31) As $\alpha^2 \ge \hbar$ the resulting term (instead of (4.13)) does not exceed (4.13) $\times (\hbar/\alpha^2|k''|)^l \times \hbar^{-1}$ and as $\alpha^2 \le \hbar$ the resulting term does not exceed $C\hbar^{1-\frac{1}{2}\kappa} \times \bigl(1+\alpha^2|k''|\hbar^{-1}\bigr)^{-l} \times \hbar^{-1}$

and we can rewrite the result in both cases as

$$C\hbar^{-\frac{1}{2}\kappa}\bigl(1+\alpha^2|k''|\hbar^{-1}\bigr)^{-l}.$$

Multiplying by $\mu^\kappa h^{-2}|k''|^{-1}\alpha^2$ we get after summation with respect to $k''\ne 0$

$$C\mu^{\frac{1}{2}\kappa}h^{-2-\frac{1}{2}\kappa}\alpha^2(\hbar/\alpha^2)^l \qquad \text{as}\quad \alpha^2\ge h, \tag{4.32}$$

$$C\mu^{\frac{1}{2}\kappa}h^{-2-\frac{1}{2}\kappa}\alpha^2\bigl(1+(\log(\hbar/\alpha^2))_+\bigr) \qquad \text{as}\quad \alpha^2\le \hbar. \tag{4.33}$$

So we have proven

Proposition 4.1.10. *For the pilot-model operator with* $1\le\mu\le h^{-1}$, $\alpha\ge\mu^{-1}$

$$|\mathsf{I}^{\mathsf{T}}_{(\alpha)}-\mathcal{I}^{\mathsf{W}}_{(\alpha)}| \le (4.15)+(4.32) \qquad \textit{as}\quad \alpha^2\ge\hbar, \tag{4.34}$$

$$|\mathsf{I}^{\mathsf{T}}_{(\alpha)}-\mathcal{I}^{\mathsf{W}}_{(\alpha)}| \le (4.28)+(4.29)+(4.33) \qquad \textit{as}\quad \alpha^2\le\hbar. \tag{4.35}$$

4.2 General operators

Consider now general operators satisfying (0.5)–(0.6) and either (0.7) or (0.8) or $(0.8)^+$, $\tau=0$.

First, using proposition 3.1.1, remark 3.1.4(ii),(ii) and propagation results we conclude that

Proposition 4.2.1. *Let conditions* (0.5)–(0.6) *be fulfilled. Then*

(i) Contribution of zone $\{|x-y|\ge C_0\mu^{-1}\}$ *to* I^{T} *does not exceed* $C\mu^{\kappa+1}h^{-1}$ *and*

(ii) Under condition (0.7) *contribution of time* $\{|t|\ge C_0\}$ *(before rescaling) to* I^{T} *does not exceed* $C\mu^{\kappa+1}h^{-1}$.

Note that $O(\mu^{\kappa+1}h^{-1})$ is smaller than the remainder estimate $R^{\mathsf{W}}(1)$ or $R^{\mathsf{W}}_1(1)$ and as long as we are interested in Weyl approximation $\mathcal{I}^{\mathsf{W}}$ (even with correction term) we should be completely happy with it.

Under condition (0.7) we are done, immediately arriving to statement (i) and related part of (iii) of theorem 4.2.3 below.

Under condition (0.8) let us introduce

$$\alpha(x)=\epsilon_0|\nabla VF^{-1}|+\frac{1}{2}\bar{\alpha},\qquad \bar{\alpha}=C_0\max\bigl(\mu^{-1},(\mu h)^{\frac{1}{2}-\delta}\bigr)\tag{4.36}$$

and consider covering of $B(0,1)$ by α-admissible elements. In virtue of proposition 4.2.1(i) we need to consider only pairs (x,y) belonging to the same element. Due to rescaling, contribution of one such element to $|I^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed $C\mu h^{-1-\kappa}\alpha^2$ and therefore

(4.37) Under condition (0.8) contribution of zone

$$\{(x,y):|x-y|\le C_0\mu^{-1},\ |\nabla VF^{-1}|\le\alpha\}$$

to $|I^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed $C\mu h^{-1-\kappa}\alpha^2$.

Therefore we need to consider only zone where $|\nabla VF^{-1}(x)|\ge\alpha$ and $|x-y|\le C_0\mu^{-1}$. We are going to prove that

Proposition 4.2.2. *Let conditions* (0.5)–(0.6) *be fulfilled. Further, let* $\mu\le h^{-1}$. *Then*

(i) For each z *with* $\alpha(z)\ge\bar{\alpha}$ *contribution of zone*

$$\{(x,y)\in B(z,\alpha(z)),\ |x-y|\le C_0\mu^{-1}|\}$$

to $|\mathsf{I}^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ *does not exceed* $R^{\mathsf{W}}(\alpha)$ *which is defined as the right-hand expression of* (4.15) *or* (4.25).

(ii) Further, as $\alpha^2\ge\mu^2h$, $1<\kappa<2$ *this contribution does not exceed* $R^{\mathsf{W}}_1(\alpha)$ *which is defined as the right-hand expression of* (4.25).

Note that both $R^{\mathsf{W}}(\alpha)$ and $R^{\mathsf{W}}_1(\alpha)$ contain α in the positive powers and thus after integration with respect $\alpha^{-1}\,d\alpha$ we get their values as $\alpha=1$.

Therefore we arrive to statement (ii) and related part of (iii) of theorem 4.2.3:

Theorem 4.2.3. *Let conditions* (0.5)–(0.6) *be fulfilled. Then as* $\mu \le h^{-1}$

(i) Under condition (0.7)

$$|\mathsf{I} - \mathcal{I}^{\mathsf{W}}| \le C\mu^{-1}h^{-1-\kappa} + R^{\mathsf{W}} \tag{4.38}$$

with

$$R^{\mathsf{W}} := C\begin{cases} \mu^{\kappa+1}h^{-1} & \text{as } \kappa < 1,\\ \mu^2 h^{-\kappa} & \text{as } \kappa > 1,\\ \mu^2 h^{-1}|\log(\mu h)| & \text{as } \kappa = 1.\end{cases} \tag{4.39}$$

(ii) Under condition (0.8) [13]

$$|\mathsf{I} - \mathcal{I}^{\mathsf{W}}| \le C\mu^{-1}h^{-1-\kappa} + R^{\mathsf{W}} + C\mu^2 h^{-\kappa}(\mu h)^{-\delta}; \tag{4.40}$$

(iii) As $\mu^2 h \le 1$ *and* $1 < \kappa < 2$ *in these estimates one can replace* $\mathcal{I}^{\mathsf{W}}$ *by* $\mathcal{I}^{\mathsf{W}} + \mathcal{I}^{\mathsf{W}}_{\mathsf{corr}}$ *and* R^{W} *by*

$$R^{\mathsf{W}}_1 := \begin{cases} \mu^{2\kappa}h^{-1} & 1 < \kappa < \dfrac{3}{2},\\ \mu^{6-2\kappa}h^{2-2\kappa} & \dfrac{3}{2} < \kappa < 2,\\ \mu^3 h^{-1}(1 + |\log(\mu^2 h)|) & \kappa = \dfrac{3}{2}.\end{cases} \tag{4.41}$$

Remark 4.2.4. (i) Actually as $\mu^2 h \le 1$, $\kappa \ge \frac{3}{2}$ this correction term does not exceed $C\mu^{-1}h^{-1-\kappa}$.

(ii) Consider other terms in the estimates. Then (4.28), (4.29), (4.33) contain α in the positive powers and (4.32) contains α in the negative power and thus after integration with respect $\alpha^{-1}\,d\alpha$ we get their values as $\alpha = (\mu h)^{\frac{1}{2}}$ i.e. (4.28), (4.29) become $C\mu^{\frac{1}{2}\kappa+2}h^{-\frac{1}{2}\kappa}$ which is less than (4.39) and (4.41) while (4.32), (4.33) become $C\mu^{\frac{1}{2}\kappa+1}h^{-1-\frac{1}{2}\kappa}$ which is larger than (4.39). Therefore selecting $\alpha \ge (\mu h)^{\frac{1}{2}-\delta}$ saves us from all these problems for the price of $\delta \ne 0$.

Even as $h^{-1+\delta} \le \mu \le h^{-1}$ it works as $C\mu^{\frac{1}{2}\kappa+1}h^{-1-\frac{1}{2}\kappa}$ acquires factor $(\mu h)^l$ which makes it subordinate to (4.39).

Problem 4.2.5. *Prove above results with* $\delta = 0$.

---

[13] If $(0.8)^+$ fails the first term should acquire factor $(1 + \mu h|\log h|)$ but it is important only as μ is close to h^{-1} but then the other terms dominate.

Proof of proposition 4.2.2. First of all applying proposition 2.2.2 and the same arguments as in the analysis of section 2 we arrive to

(4.42) Contribution of zone $\{|\sin(\theta(t'))|\ge C_0\hbar,\ |\sin(\theta(t''))|\ge C_0\hbar\}$ to the error $|\mathsf{I}^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed expression (4.15) or (4.41) (in its frames).

Therefore we need to explore contributions of three remaining zones

$$\{|\sin(\theta(t'))|\le 2C_0\hbar,\ |\sin(\theta(t''))|\le 2C_0\hbar\} \tag{4.43}$$

and

$$\{|\sin(\theta(t'))|\ge 2C_0\hbar,\ |\sin(\theta(t''))|\le C_0\hbar\} \tag{4.44}$$

(the same analysis would cover its tween zone $\{|\sin(\theta(t'))|\ge 2C_0\hbar,\ |\sin(\theta(t''))|\le C_0\hbar\}$).

Note first that the standard propagation arguments imply that we get a factor $C(|z_1|+|z_2|)^{-l}(|\sin(\theta(t'))|+\varepsilon|t'|)^l$ and the similar factor with t' replaced by t''.

Zone (4.43) *as* $\varepsilon\ge C_1\hbar$. Then in the classical propagation shift $\varepsilon|k'|$ is always greater than $\hbar$.

Obviously contribution of zone (4.43), intersected with (k',k'') windings $\times\{(x,y):|x-y|\ge\rho\}$ to the Tauberian expression does not exceed

$$C\mu^2h^2|k'|^{-1}|k''|^{-1}\rho^{-\kappa}\sup_t\iint|\tilde{U}(x,y,t)|^2\,dxdy\le C\mu^2h^{2-d}\rho^{-\kappa}\alpha^d \tag{4.45}$$

where factor $\hbar^2$ is due to integration with respect to t',t'' and $\tilde{U}$ means cut-off of U to the energy level $|\tau|\le c$.

Plugging (as $d=2$)

$$\rho=\varepsilon\mu^{-1}\max(|k'|,|k''|), \tag{4.46}$$

$\varepsilon=\mu^{-1}\alpha$ we get

$$C\mu^{2\kappa+2}|k'|^{-1}|k''|^{-1}\alpha^{2-\kappa}\bigl(\max(|k'|,|k''|)\bigr)^{-\kappa}. \tag{4.47}$$

Summation by k',k'' returns $C\mu^{2\kappa+2}\alpha^{2-\kappa}$ which is less than the right-hand expression of (4.15) or (as $\alpha^2\ge\mu^2h$, $1<\kappa<2$) the right-hand expression of (4.25).

Meanwhile, estimating $\tilde{U}(x,y,t)$ by h^{-d} and integrating by $|x-y|\le \rho$ and then integrating by t',t'' over $|\theta(t')-\pi k'|\le \hbar$, $|\theta(t'')-\pi k''|\le \hbar$ we get

$$Ch^{2-2d}\mu^2\rho^{d-\kappa}|k'|^{-1}|k''|^{-1}\bigl(h/\rho\bigr)^l\alpha^d \tag{4.48}$$

where the last factor is due to propagation results; plugging $d=2$, (4.46) we get

$$C\mu^2h^{-\kappa}|k'|^{-1}|k''|^{-1}\alpha^2\bigl(\hbar/\varepsilon\max(|k'|,|k''|)\bigr)^l; \tag{4.49}$$

after summation we get $C\mu^2h^{-\kappa}\bigl(\hbar/\varepsilon\bigr)^l\alpha^2$ which is again less than the right-hand expression of (4.15) or (as $\alpha^2\ge\mu^2h$, $1<\kappa<2$) the right-hand expression of (4.25).

Zone (4.43) *as* $\varepsilon\le C_1\hbar$*.* Taking sum of (4.47) over $k',k'':\max(|k'|,|k''|)\ge \hbar/\varepsilon$ we get

$$C\mu^{2\kappa+2}\alpha^{2-\kappa}(1+|\log\mu^2h/\alpha|)\times(\mu^2h/\alpha)^{-\kappa}=C\mu^2h^{-\kappa}\alpha^2(1+|\log\mu^2h/\alpha|)$$

while taking sum of (4.49) we get the right-hand expression as well. Obviously, it is less than the right-hand expression of (4.15).

Thus we need to consider only $k',k'':\max(|k'|,|k''|)\le\hbar/\varepsilon$.

Using the same arguments as before, we conclude that the contribution of zone (4.43), intersected with (k',k'') windings $\times\{(x,y):|x-y|\ge\hbar\}$ to the Tauberian expression does not exceed $C\mu^2h^{-\kappa}|k'|^{-1}|k''|^{-1}$; similarly contribution of zone (4.43), intersected with (k',k'') windings $\times\{(x,y):|x-y|\le\hbar\}$ to the Tauberian expression does not exceed $C\mu^2h^{-\kappa}|k'|^{-1}|k''|^{-1}\alpha^2$ as well.

After summation we get $C\mu^2h^{-\kappa}(1+|\log\mu^2h/\alpha|)^2$ which again is less than the right-hand expression of (4.15).

Zone (4.44) *as* $\varepsilon\ge C_1\hbar$*.* Now shift with respect to $(x-y)$ is $\mu^{-1}\max\bigl(\varepsilon|t|,|\varepsilon t'\mp\sin(t')|\bigr)$; then repeating analysis in zone (4.43) we arrive to (4.47) for $1<\kappa<2$ and to (4.49) for $0<\kappa<2$; as $0<\kappa\le 1$ one should replace (4.47) by

$$C\mu^{2\kappa+2}\hbar|k'|^{-1}|k''|^{-1} \tag{4.50}$$

with a logarithmic factor as $\kappa=1$ and summation with respect to k',k'' results in $C\mu^{3\kappa+2}h(1+|\log h|)^2$ which again is less than $R^{\mathsf{W}}(\alpha)$.

Zone (4.44) *as* $\varepsilon\le C_1\hbar$*.* With the above adjustment arguments of zone (4.43) work here as well. □

4.3 Perturbations

4.3.1 General scheme

Contribution of $k'=k''\ne 0$. Consider first case $k'=k''=k$. Assume first that $\alpha^2\ge \hbar$.

Recall that k-th tick' contribution to I^T was not exceeding expression (4.13) multiplied by $\mu^\kappa h^{-2}\alpha^2|k|^{-2}$ i.e.

$$C\mu^{\kappa+2}\alpha^2|k|^{-2}\hbar^{-1-\frac{1}{2}\kappa}\begin{cases}(\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1\end{cases}\tag{4.51}$$

(as $|\sin(t')|\le\hbar$ or $|\sin(t'')|\le\hbar$ one should use proposition 4.2.2). Then we can suspect that the successive approximations bringing factor $\sigma T/\hbar$ as U is replaced by U^0 simultaneously brings the same factor when we replace U by $\bar{U}$ in I; so contribution of k-th tick to the error is estimated by the previous expression multiplied by $C\sigma|k|/\hbar$ i.e.

$$C\sigma\mu^{\kappa+2}\alpha^2|k|^{-1}\hbar^{-2-\frac{1}{2}\kappa}\begin{cases}(\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1.\end{cases}\tag{4.52}$$

and summation over $k:|k|\le\bar{k}:=\hbar/\sigma$ results in

$$C\sigma\mu^{\kappa+2}\alpha^2|\log\hbar/\sigma|\hbar^{-2-\frac{1}{2}\kappa}\begin{cases}(\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1.\end{cases}\tag{4.53}$$

Remark 4.3.1. (i) Here σ is the magnitude of perturbation and as we approximate by pilot-model it is $O(\mu^{-2})$ but if we are trying to use the magnetic Weyl approximation it is $O(\varepsilon)$.

(ii) Expression (4.53) provides us with the answer as $\alpha^2\ge\hbar$; as $\alpha^2\le\hbar$ we need to recall that if we are interested only in zone $\{|x-y|\le C_0\mu^{-1}\}$ before rescaling then $|z|\le C_0$ and we need therefore there reset $\hbar/\alpha^2$ to 1 and also $|s|$ to 1 resulting in

$$C\sigma\mu^{\kappa+2}\alpha^2|\log\hbar/\sigma|\hbar^{-2-\frac{1}{2}\kappa}\tag{4.54}$$

(iii) In these settings we need to assume that $T \le C_0\varepsilon^{-1}$ i.e. $\bar{k} \le C_0\varepsilon^{-1}$ i.e.

$$\mu h \le \sigma/\varepsilon. \tag{4.55}$$

This inequality always holds as $\sigma=\varepsilon$ but it may fail as $\sigma=\mu^{-2}$; in this case we either reset $\bar{k}$ to $C\varepsilon^{-1}$ or reevaluate σ and analyze separately in both cases (see in the next subsubsection).

(iv) Note, as $\mu h \le \sigma/\varepsilon$ summation (4.51) for $k:|k|\ge \bar{k}$ brings the same expression albeit without logarithmic factor; otherwise if we reset $\bar{k}=C\varepsilon^{-1}$ this sum returns $C\mu^\kappa h^{-1}$.

(iv) However for the pilot-model approximation we reexamine zone $\{|x-y|\ge C_0\mu^{-1}\}$ and improve the remainder estimate of (iv)

Contribution of $0\ne k'\ne k''\ne 0$. Next we need to consider $0\ne k'\ne k''\ne 0$. Recall that as $\alpha^2\ge\hbar$ the contribution of such pair to I^T does not exceed

$$C\mu^{\kappa+2}\alpha^2|k'|^{-1}|k''|^{-1}(\hbar/\alpha^2|k'-k''|)^l\hbar^{-1-\frac{1}{2}\kappa}\times \begin{cases} (\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1; \end{cases} \tag{4.56}$$

multiplying by $C\sigma|k'|/\hbar$ (as perturbation goes to the first "factor") we get

$$C\sigma\mu^{\kappa+2}\alpha^2|k''|^{-1}(\hbar/\alpha^2|k'-k''|)^l\hbar^{-2-\frac{1}{2}\kappa}\times \begin{cases} (\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1. \end{cases} \tag{4.57}$$

Then summation with respect to k' of $(\hbar/\alpha^2|k'-k''|)^l$ returns $(\hbar/\alpha^2)^l$ as $\hbar\le\alpha^2$ and summation with respect to k'' then returns (4.53) with an extra factor $(\hbar/\alpha^2)^l$ and therefore is less than (4.53).

On the other hand, as $\hbar\ge\alpha^2$ summation with respect to k' of 1 returns $\hbar/\alpha^2$ and as we recall that big left brace expression should be replaced by 1 we arrive after summation with respect to k'' to

$$C\sigma\mu^{\kappa+2}\hbar^{-1-\frac{1}{2}\kappa}(1+(\log\hbar/\sigma)) \tag{4.58}$$

which is (4.54) with an extra factor $(\hbar/\alpha^2)$ and therefore is greater than (4.53).

Contribution of $0 = k'' \ne k'$. If $k' \ne 0$, $k'' = 0$ we recall that $|k''|^{-1}$ should be replaced by $\hbar^{-1}$ in (4.56)–(4.57) and perturbation brings factor $\sigma|k'|/\hbar$ so we get instead of (4.57)

$$C\sigma\mu^{\kappa+2}\alpha^2(\hbar/\alpha^2|k'|)^l\hbar^{-3-\frac{1}{2}\kappa}\begin{cases}(\hbar/\alpha^2)^{\frac{1}{2}\kappa} & \kappa<1,\\ (\hbar/\alpha^2)^{1-\frac{1}{2}\kappa} & \kappa>1,\\ (\hbar/\alpha^2)^{\frac{1}{2}}(1+|\log(\hbar/\alpha^2)|) & \kappa=1.\end{cases}$$

Then summation with respect to k' returns

$$C\sigma\mu^{\kappa+2}\alpha^2(\hbar/\alpha^2)^l\hbar^{-3-\frac{1}{2}\kappa} \tag{4.59}$$

as $\hbar \le \alpha^2$ and

$$C\sigma\mu^{\kappa+2}\hbar^{-2-\frac{1}{2}\kappa} \tag{4.60}$$

as $\hbar \ge \alpha^2$; (4.60) is even greater than (4.58).

Contribution of $0 = k'$. With perturbation factors are no more of the same rights and we need to consider this case as well.

(i) If $k' = 0$, $k'' \ne 0$ we recall that $|k'|^{-1}$ should be replaced by $\hbar^{-1}$ in (4.56)–(4.57) but perturbation brings factor σ so together we get factor $\sigma\hbar^{-1}$ which falls into into case $k' = 1$, $k'' \ne 0$.

(ii) Similarly, if $k' = k'' = 0$ we fall into case $k' = 1$, $k'' = 0$.

In total we get (4.53)+(4.59) as $\alpha^2 \ge \hbar$ and (4.60) as $\alpha^2 \le \hbar$. So, we arrive to

Proposition 4.3.2. *Let conditions* (0.5)–(0.6) *be fulfilled in* $B(0,1)$ *and let* $\alpha := \epsilon_0|\nabla V/F(\bar{x})| \ge C_0\mu^{-1}$ *where* $\bar{x} \in B(0,\frac{1}{2})$.

Let $\bar{e}_z(x,y,\tau)$ *be a Schwartz kernel of the spectral projector of* *either* *the magnetic Weyl approximation (then* $\sigma = \alpha\mu^{-1}$*)* *or* *the pilot-model approximation (then* $\sigma = \mu^{-2}$*) at point* z*. Let* $T \ge \min\bigl(\epsilon_0\hbar/\sigma, C_0\mu\alpha^{-1}\bigr) \ge 1$.

Let $\psi, \psi_1 \in \mathscr{C}_0^\infty(B(0,1))$.

(i) Then as $\alpha^2 \ge \hbar$, $\gamma = C_0\mu^{-1}$

$$\begin{gathered}\mathsf{R}_2 = \mathsf{R}_2(\alpha,\gamma) := \\ |\int \Omega_{\alpha,\gamma}\bigl(\frac{1}{2}(x+y), x-y\bigr)\Bigl(|e^{\mathsf{T}}(x,y,0)|^2 - |\bar{e}^{\mathsf{T}}_{\frac{1}{2}(x+y)}(x,y,0)|^2\Bigr)\,dxdy| \le \\ (4.53)+(4.59)\end{gathered} \tag{4.61}$$

where

$$\Omega_{\alpha,\gamma}(x,z) := \psi\bigl(\alpha^{-1}(x-\bar{x})\bigr)\psi_1\bigl(\gamma^{-1}z\bigr);$$

(ii) As $\alpha^2 \le \hbar$, $\gamma = C_0\mu^{-1}$ we get

$$\mathsf{R}_2 \le (4.54) + (4.60). \tag{4.62}$$

(ii) Further, in this inequality one can replace $\bar{\mathsf{e}}^{\mathsf{T}}$ by $\bar{\mathsf{e}}$.

Proof. The easy albeit tedious proof following our standard technique we leave to the reader. □

4.3.2 Improvement

We want to apply arguments of the previous subsubsection "Improvement" to improve results of the previous subsubsection. One can see easily that improvement makes sense only if we can improve $k' = k'' = 1$ and thus as $\alpha^2 \ge \mu^2 h$. This excludes magnetic Weyl approximation.

In this case $R_1^{\mathsf{W}}(\alpha)$ defined by (4.25) acquires factor $\mu^{-3}h^{-1}$:

$$C\begin{cases}\mu^{2\kappa-3}h^{-2}\alpha^{2-\kappa} & 1<\kappa<\dfrac{3}{2},\\ \mu^{3-2\kappa}h^{1-2\kappa}\alpha^{3\kappa-4} & \dfrac{3}{2}<\kappa<2,\\ h^{-2}\alpha^{\frac{1}{2}}(1+|\log(\mu^2h/\alpha^2)|) & \kappa=\dfrac{3}{2}.\end{cases} \tag{4.63}$$

This replaces (4.53) and (4.59) becomes negligible.

So, we arrive to

Proposition 4.3.3. *Let conditions of proposition 4.3.2 be fulfilled, we consider the pilot-model approximation and let $\alpha^2 \ge \mu^2 h$. Then*

(i) $\mathsf{R}_2(\alpha,\gamma)$ does not exceed (4.63).

(ii) Further, in this inequality one can replace $\bar{\mathsf{e}}^{\mathsf{T}}$ by $\bar{\mathsf{e}}$.

Proof. The easy albeit tedious proof following our standard technique we leave to the reader. □

4.3.3 Results. I

Let condition (0.7) be fulfilled.

Magnetic Weyl approximation. If we are looking for the magnetic Weyl approximation, we should assume that $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ and pick up $\sigma = \varepsilon = \mu^{-1}$, $\alpha = 1$ and look what happens with expressions (4.53) and (4.59). The former returns the last term in the estimate (4.64) below and the latter returns the lesser expression $C\mu^{\kappa-1}h^{-2}\hbar^{l}$. So we arrive to estimate (4.64) of theorem 4.3.4 below albeit with an extra factor $\bigl(1 + (\log \mu^2 h)_+\bigr)$ in front of the big left brace.

To get rid off it let us consider two-term successive approximation. Then to estimate a remainder one needs to sum the same expressions as before but with an extra factor which is either $\mu^{-2}h^{-1}|k'|$ or $\mu^{-2}h^{-1}|k''|$.

This really does not matter for $k' = k'' \ne 0$ and we sum $\mu^{-2}h^{-1}$ instead of $|k|^{-1}$ from $|k| = 1$ to $|k| = \mu^2 h$ resulting in 1 rather than logarithm. As $0 \ne k' \ne k'' \ne 0$ we get the same answer albeit with a factor $(\mu h)^l$ as $\alpha \asymp 1$. If $k' = 0$ and/or $k'' = 0$ we replace $|k'|^{-1}$ and/or $|k''|^{-1}$ by $\hbar^{-1}$ but this is more than compensated by $(\mu h)^l$ but effectively $T' = \hbar$ and/or $T'' = \hbar$ so in fact we reset these cases to $k' = 1$ and/or $k'' = 1$.

On the other hand, the second term in the successive approximation refers to perturbation being strictly $\alpha(x_1 - y_1)$ but this leads to 0 as Ω is even with respect to $(x - y)$. Thus we arrive to estimate (4.64) of theorem 4.3.4 below.

Theorem 4.3.4. *Let conditions* (0.5)–(0.7) *be fulfilled. Then as* $h^{-\frac{1}{2}} \le \mu \le h^{-1}$

$$|\mathsf{I} - \mathcal{I}^{\mathsf{MW}}| \le C\mu^{-1}h^{-1-\kappa} + C\begin{cases} \mu^{\kappa-1}h^{-2} & 0 < \kappa < 1,\\ h^{-1-\kappa} & 1 < \kappa < 2,\\ h^{-2}(1 + |\log(\mu h)|) & \kappa = 1.\end{cases} \tag{4.64}$$

Pilot-model approximation. If we are looking for the pilot-model approximation, we should assume that $h^{-\frac{1}{3}} \le \mu \le h^{-1}$ and pick up $\sigma = \mu^{-2}$, $\varepsilon = \mu^{-1}$, $\alpha = 1$. We need to consider variants:

(i) $h^{-\frac{1}{3}}\le \mu\le h^{-\frac{1}{2}}$. As $0<\kappa\le 1$ (4.53) becomes $C\mu^{\kappa-2}h^{-2}(1+|\log \mu^3 h|)$ (with another logarithmic factor as $\kappa=1$ and it is larger than $C\mu^\kappa h^{-1}$ (which is the contribution of $\{|x-y|\ge C_0\mu^{-1}\}$).

As $1<\kappa<2$ we can replace (4.53) by (4.63) which is then $O(\mu^{-1}h^{-1-\kappa})$ and $\mu^\kappa h^{-1}$ is also $O(\mu^{-1}h^{-1-\kappa})$.

Noting that (4.59) is negligible we arrive to statement (i) of theorem 4.3.6 below.

(ii) $h^{-\frac{1}{2}}\le \mu\le h^{-1}$. In this case we note that (4.59) is still less than (4.53) but $\mu^\kappa h^{-1}$ may become leading term. To improve this term let us consider contribution of zone $|x-y|\asymp \gamma\ge C_0\mu^{-1}$ into an error: we need to take $\sigma=\gamma^2$ and $T\asymp \mu^2\gamma$ so we get the estimate of perturbation

$$C\mu h^{-1}T^{-1}\gamma^{-\kappa}\times \sigma T/\hbar \asymp Ch^{-2}\gamma^{2-\kappa} \tag{4.65}$$

recall that unperturbed term does not exceed

$$C\mu h^{-1}T^{-1}\gamma^{-\kappa}\asymp C\mu^{-1}h^{-1}\gamma^{-\kappa-1} \tag{4.66}$$

because the drift propagation does not expand $B(x,\rho)$ with $\rho\ge C_0\mu^{-1}$ but merely shifts it. We sum (4.65) as $\gamma^2 T\le \hbar$ i.e. $\gamma^3\le \mu^{-1}h$ i.e. $\gamma\le\bar{\gamma}=\mu^{-\frac{1}{3}}h^{\frac{1}{3}}=\mu^{-1}(\mu^2 h)^{\frac{1}{3}}$.

Then we get $Ch^{-\frac{4}{3}-\frac{1}{3}\kappa}\mu^{\frac{1}{3}\kappa-\frac{2}{3}}$ and we arrive to statement (ii) of theorem 4.3.6 below.

Remark 4.3.5. Obviously arguments of the magnetic Weyl approximation allowing us to get rid off logarithmic factor still work albeit the second term does not necessarily vanish as the perturbation is quadratic. The logarithmic factor in estimates (4.67) and (4.69) does not seem be worth of trouble to write this second term.

Theorem 4.3.6. *Let conditions* (0.5)–(0.7) *be fulfilled. Then*

(i) As $h^{-\frac{1}{3}}\le \mu\le h^{-\frac{1}{2}}$

$$\begin{aligned}|\mathsf{I}-\bar{\mathsf{I}}|\le C\mu^{-1}h^{-1-\kappa}+&\\ C|\log\mu^3 h|&\begin{cases}\mu^{\kappa-2}h^{-2} & 0<\kappa<1,\\ 0 & 1<\kappa<2,\\ \mu^{-1}h^{-2}(1+|\log h|) & \kappa=1\end{cases}\end{aligned}\tag{4.67}$$

where

$$\bar{I} := \int \Omega_{1,1}\bigl(\frac{1}{2}(x+y), x-y\bigr)|\bar{e}_{\frac{1}{2}(x+y)}(x,y,0)|^2\, dxdy \tag{4.68}$$

(ii) As $h^{-\frac{1}{2}} \le \mu \le h^{-1}$

$$|\mathsf{I}-\bar{\mathsf{I}}| \le C\mu^{-1}h^{-1-\kappa} + Ch^{-\frac{4}{3}-\frac{1}{3}\kappa}\mu^{\frac{1}{3}\kappa-\frac{2}{3}} + \\ C|\log h| \begin{cases} \mu^{\kappa-2}h^{-2} & 0<\kappa<1,\\ \mu^{-1}h^{-1-\kappa} & 1<\kappa<2,\\ \mu^{-1}h^{-2}(1+|\log(\mu h)|) & \kappa=1. \end{cases} \tag{4.69}$$

4.3.4 Results. II

Assume now that condition (0.8) is fulfilled instead of (0.7) and estimate an approximation error again.

Magnetic Weyl approximation Let us consider the magnetic Weyl approximation first. Then we should assume that $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ and pick up $\sigma = \varepsilon = \alpha\mu^{-1}$. Recall that the contribution of $\{(x,y) : |x-y| \ge C_0\mu^{-1}\}$ does not exceed $C\mu^\kappa h^{-1}$ and therefore we need to consider zone

$$\{(x,y) : |x-y| \le C_0\mu^{-1},\ \alpha(x) \ge C_1\mu^{-1}\} \tag{4.70}$$

where also $(1-\epsilon) \le \alpha(y)/\alpha(x) \le (1+\epsilon)$ [14].

Therefore we can use α-admissible partition in zone (4.69) and we need just look what happens with expressions (4.53) and (4.59) after integration with respect to $\alpha^{-1}d\alpha$ as $\alpha \ge (\mu h)^{\frac{1}{2}}$ and (4.60) $\mu^{-1} \le \alpha \le (\mu h)^{\frac{1}{2}}$.

Obviously integral of (4.53) resets to its value as $\alpha = 1$ as under assumption (0.7); integrals of (4.59) and (4.60) reset to their values as $\alpha = (\mu h)^{\frac{1}{2}}$ which are equal and less than what we got already.

Therefore estimate (4.64) still holds[15] and we arrive to statement (i) of theorem 4.3.7 below after we apply the same arguments with two-term approximation as in the previous subsubsection to get rid off the logarithmic factor.

---

[14] And add contribution of the zone $\{(x,y) : \alpha(x) \le \bar{\alpha} := C_1\mu^{-1},\ \alpha(y) \le \bar{\alpha}\}$ which would be $C\mu h^{-1-\kappa} \times \bar{\alpha}^2 = C\mu^{-1}h^{-1-\kappa}$ in virtue (0.8).

[15] Albeit the first term $C\mu^{-1}h^{-1-\kappa}$ acquires factor $|\log h|$ if assumption $(0.8)^+$ fails: see Tauberian estimate.

Pilot-model approximation Let us consider the pilot-model approximation; then again $\sigma = \mu^{-2}$. Then we need to consider two cases.

(a) $h^{-\frac{1}{3}} \le \mu \le h^{-\frac{1}{2}}$. In this case we are completely happy with the zone $\{(x, y) : |x - y| \ge C_0\mu^{-1}\}$.

As $0 < \kappa \le 1$ we need to integrate with respect to $\alpha^{-1}d\alpha$ (4.53) and (4.59) as $\alpha \ge (\mu h)^{\frac{1}{2}}$ and (4.60) as $\alpha \le (\mu h)^{\frac{1}{2}}$. The first integral resets to (4.53) as $\alpha = 1$; integrals of (4.59) and (4.60) reset to their values as $\alpha = (\mu h)^{\frac{1}{2}}$ which are equal (up to a logarithmic factor) and less than what we got already. Thus we arrive to statement (ii) of theorem 4.3.7 below.

As $1 < \kappa < 2$ we need to replace integral of (4.53) by its integral over zone $(\mu h)^{\frac{1}{2}} \le \alpha \le \mu h^{\frac{1}{2}}$ plus integral of (4.63) over zone $\mu h^{\frac{1}{2}} \le \alpha \le 1$. These integrals are reset to their integrands values as $\alpha = (\mu h)^{\frac{1}{2}}$ and $\alpha = 1$ respectively; the latter is larger than the former albeit not necessarily larger than integral of (4.60).

Thus we arrive to statement (iii) of theorem 4.3.7 below.

(b) $h^{-\frac{1}{2}} \le \mu \le h^{-1}$. In this case the contribution of $\{(x, y) : |x-y| \le C_0\mu^{-1}\}$ does not exceed the right-hand expression of (4.69) but we need to reexamine zone $\{(x, y) : |x - y| \ge C_0\mu^{-1}\}$ which we split into two: $\{|x - y| \le \epsilon_0\alpha(x)\}$ and $\{|x - y| \ge \epsilon_0\alpha(x)\}$.

Repeating corresponding arguments of the analysis under assumption (0.7) we conclude that the contribution of the former zone does not exceed what we got then. In the latter zone $T \asymp \mu^2$ and its contribution does not exceed $C\mu^{-1}h^{-1-\kappa}$.

Thus we arrive to statement (iv) of theorem 4.3.7 below.

Theorem 4.3.7. *Let conditions* (0.5)–(0.6) *and* (0.8) *be fulfilled. Then*[15)]

(i) As $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ *estimate* (4.64) *holds;*

(ii) As $h^{-\frac{1}{3}} \le \mu \le h^{-\frac{1}{2}}$, $0 < \kappa \le 1$ *estimate* (4.67) *holds;*

(iii) As $h^{-\frac{1}{3}} \le \mu \le h^{-\frac{1}{2}}$, $1 < \kappa < 2$

$$|\mathsf{I} - \bar{I}| \le C\mu^{-1}h^{-1-\kappa} + C\mu^{\frac{1}{2}\kappa-2}h^{-\frac{1}{2}\kappa-2}(1 + |\log \mu^3 h|); \tag{4.71}$$

(iv) As $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ *estimate* (4.69) *holds.*

4.4 Superstrong magnetic field

Consider under assumption (0.8) the Schrödinger-Pauli operator with $\mu \ge h^{-1}$. Recall that according to propositions 3.2.1(ii), 3.2.2(ii) respectively

$$|\mathsf{I}| \le C\mu^{1+\frac{1}{2}\kappa}h^{-1-\frac{1}{2}\kappa}$$

and

$$|\mathsf{I}-\mathsf{I}^{\mathsf{T}}| \le C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}$$

(as $\kappa \ne 1$; as $\kappa = 1$ the latter estimate holds with an extra factor $|\log \mu|$ in its right-hand expression).

Let us assume that $F=1$ to avoid some unpleasant correction terms as we reduce operator to one with $F=1$. Recall that now $\hbar := \mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ is the magnitude of the cyclotron radius.

Magnetic Weyl approximation. Consider first the magnetic Weyl approximation. Then $\sigma = \alpha\hbar = \alpha\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ and $T = \sigma^{-1}\hbar = \alpha^{-1}\mu^{\frac{3}{2}}h^{\frac{1}{2}}$; then the the shift (unrescaled) due to magnetic drift does not exceed $C\alpha\mu^{-2}T = C\hbar$, which justifies the choice of σ.

Therefore as $\alpha \ge \bar{\alpha} = C_1\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ the contribution of α-ball intersected with $\{(x,y) : |x-y| \le \bar{\gamma} := C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}\}$ to $|\mathsf{I}^{\mathsf{T}} - \mathcal{I}^{\mathsf{MW}}|$ does not exceed $C\mu h^{-1}\sigma \times \mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa} \asymp C\alpha\mu^{\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{1}{2}-\frac{1}{2}\kappa}$. Then summation with respect to all such balls does not exceed $C\mu^{\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{1}{2}-\frac{1}{2}\kappa}$, as in estimate (4.72) below.

Meanwhile the contributions to both I and $\bar{\mathsf{I}}$ of zone $\{(x,y) : |x-y| \ge \bar{\gamma}\}$ does not exceed $C\bar{\gamma}^{-1-\kappa}$ which is also of the same magnitude.

Finally, the contributions to both I and $\bar{\mathsf{I}}$ of zone $\{(x,y) : \alpha(x) \le \bar{\alpha},\ \alpha(y) \le \bar{\alpha}\}$ do not exceed $C\mu^{1+\frac{1}{2}\kappa}h^{-1-\frac{1}{2}\kappa} \times \bar{\alpha}^2 \asymp C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}$ which is less than the right-hand expressions in both (4.72) and (4.73).

Thus we arrive to estimate (4.72) below.

Pilot-model approximation. Meanwhile for the pilot-model approximation $\sigma = \mu^{-1}h$ as long as $T \le T_* := \alpha^{-1}\mu^{\frac{3}{2}}h^{\frac{1}{2}}$; so we conclude that as $\alpha \ge \bar{\alpha}$, the contribution to $(\mathsf{I} - \bar{\mathsf{I}})$ of α-ball intersected with $\{(x,y) : |x-y| \le \bar{\alpha}\}$ does not exceed $C\alpha\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}$. After summation with respect to $\alpha^{-1}d\alpha$ we get $C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}$ which is less than the right-hand expression in (4.73).

Now we need to consider larger T and γ; then $\sigma = \gamma^2$ and $T \asymp \alpha^{-1}\mu^2\gamma$. Then we should define $\bar{\gamma}$ so that $\sigma T \le \hbar$ iff $\gamma \le \bar{\gamma}$. Therefore we pick up $\bar{\gamma} = \mu^{-\frac{1}{3}}h^{\frac{1}{3}}\alpha^{\frac{1}{3}}$ and matching $\bar{T} = \mu^{\frac{4}{3}}h^{\frac{1}{3}}\alpha^{-\frac{2}{3}}$ and $\bar{\gamma} = \mu^{-\frac{2}{3}}h^{\frac{2}{3}}\alpha^{\frac{2}{3}}$.

Using the same arguments as above we conclude that the contribution to $(\mathsf{I}-\bar{\mathsf{I}})$ of the zone $\{(x,y):\bar\alpha\le|x-y|\le\bar\gamma\}$ and contributions to both I and $\bar{I}$ of the zone $\{(x,y):\bar\alpha\ge|x-y|\le\bar\gamma\}$ also do not exceed the right-hand expression in (4.73).

Thus we arrive to estimate (4.73) below.

Theorem 4.4.1. *For Schrödinger-Pauli operator under conditions* (0.5)–(0.6), (0.8), $\mu\ge h^{-1}$ *and* $F=1$

$$|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|\le C\mu^{\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{1}{2}-\frac{1}{2}\kappa} \tag{4.72}$$

and

$$|\mathsf{I}-\bar{\mathsf{I}}|\le C\mu^{\frac{1}{3}+\frac{1}{3}\kappa}h^{-\frac{1}{3}-\frac{1}{3}\kappa}. \tag{4.73}$$

4.5 Problems and remarks

Remark 4.5.1. The main difference between cases $1<\kappa<2$ and $0<\kappa<1$ as $\mu h\ll 1$ is that in the former case the main contribution to the remainder is delivered by (x,y) close to one another ($|x-y|\ll\mu^{-1}$) while in the latter case by (x,y) with $|x-y|\asymp\mu^{-1}$.

We can calculate $\mathcal{I}^{\mathsf{MW}}$ and $\bar{\mathsf{I}}$ plugging corresponding expressions $e_z^{\mathsf{MW}}(x,y,0)$ and $\bar{e}_z(x,y,0)$ with $z=\frac{1}{2}(x+y)$ into I.

Problem 4.5.2. *Find nice expressions for* $\mathcal{I}^{\mathsf{MW}}$ *and* $\bar{\mathsf{I}}$.

Problem 4.5.3. *As* $\mu h\le 1$ *get rid off condition (0.5); according to Chapter 13 we do not need it for estimate* $|\mathsf{I}-\mathsf{I}^{\mathsf{T}}|$ *but we want to get rid of it in estimates for* $|\mathsf{I}-\mathcal{I}^{\mathsf{W}}|$, $|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|$ *and* $|\mathsf{I}-\bar{\mathsf{I}}|$. *To do this*

(i) Assume first that condition

$$|V|+|\nabla V|\asymp 1 \tag{4.74}$$

is fulfilled and consider the scaling function $\alpha(x)=\max\bigl(\epsilon|V(x)|,\mu h, C\mu^{-1}\bigr)$.

(ii) Assume then that condition

$$|V|+|\nabla V|+|\det\operatorname{Hess}V|\asymp 1 \tag{4.75}$$

is fulfilled and consider the scaling function

$$\alpha(x)=\max\bigl(\epsilon(|V(x)|+|\nabla V(x)|^2)^{\frac{1}{2}},(\mu h)^{\frac{1}{2}},C\mu^{-1}\bigr).$$

Problem 4.5.4. *Get rid off condition (0.6) assuming instead that* $|F| + |\nabla F| \ge \epsilon$*.*

(i) Repeating arguments of section 3 and using results of Chapter 14 one can prove easily that $|\mathsf{I} - \mathsf{I}^{\mathsf{T}}| \le C\mu^{-1/2}h^{-1-\kappa}$ *as* $\mu \le h^{-1}$*. Derive similar estimates for Schrödinger and Schrödinger-Pauli operators as* $h^{-1} \le \mu \le h^{-2}$ *and for Schrödinger-Pauli operator as* $\mu \ge h^{-2}$*;*

(ii) Introducing scaling function $\ell(x) = \epsilon|F(x)|$ *in zone* $\{x:\ \ell(x) \ge \mu^{-\frac{1}{2}}\}$ *one can rescale* $\mu \mapsto \mu\ell^2$*,* $h \mapsto h\ell^{-1}$*,* $\mu h^{-1-\kappa} \mapsto \mu^{-1}h^{-1-\kappa}\ell^{-1+\kappa}$*, after integration over* $\ell^{-2}d\ell$ *we get* $C\mu^{-1}h^{-1-\kappa}\ell^{-2+\kappa}$ *calculated as* $\ell = \mu^{-\frac{1}{2}}$ *i.e.* $\mu^{-\frac{1}{2}\kappa}h^{-1-\kappa}$*. This is an estimate for the contribution of the regular zone.*

Prove the same estimate for the contribution of the singular zone as $\mu \le h^{-1}$*. Derive similar estimates as* $h^{-1} \le \mu \le h^{-2}$ *and as* $\mu \ge h^{-2}$*;*

(iii) Is it possible to upgrade the above estimate to $C\mu^{-\frac{1}{2}}h^{-1-\kappa}$ *as* $\kappa < 1$*? Explore also cases* $h^{-1} \le \mu \le h^{-2}$ *and* $\mu \ge h^{-2}$*.*

5 Pointwise asymptotics: 3D-pilot-model

5.1 Pilot-model in $\mathbb{R}^3$: propagator

Consider the *pilot-model operator*

$$A = \bar{A} := h^2D_1^2 + (hD_2 - \mu x_1)^2 + h^2D_3^2 + 2\alpha x_1 + 2\beta x_3 \tag{5.1}$$

which is the sum of 2D-pilot-model (1.1) $\bar{A}_{(2)}$ and 1D Schrödinger operator

$$B := h^2D_3^2 + \beta x_3 \tag{5.2}$$

and therefore propagator of $\bar{A}$ is

$$U(x,y,t) = U_{(2)}(x',y',t)U_{(1)}(x_3,y_3,t) \tag{5.3}$$

where $U_{(2)}(x',y',t)$ is the Schwartz kernel of the propagator $e^{ih^{-1}\bar{A}_{(2)}}$ and therefore *after rescaling* it is defined by (1.1) while $U_{(1)}(x_3,y_3,t)$ is the Schwartz kernel of the propagator $e^{ih^{-1}tB}$ and then one can prove easily that *before rescaling*

(5.4) $U_{(1)}(x_3,y_3,t)=$

$$(2\pi h)^{-1}\int \exp\Bigl(ih^{-1}\bigl((\zeta+\beta t)x_3-(\zeta-\beta t)y_3-2\zeta^2 t-\frac{2}{3}\beta^2t^3\bigr)\Bigr)\,d\zeta = \frac{1}{2}(2\pi ht)^{-\frac12}\exp\Bigl(ih^{-1}\bigl(\beta t(x_3+y_3)+\frac{1}{8}t^{-1}(x_3-y_3)^2-\frac{2}{3}\beta^2t^3\bigr)\Bigr);$$

in particular

$$U_{(1)}(x_3,x_3,t)=\frac{1}{2}(2\pi ht)^{-\frac12}\exp\Bigl(ih^{-1}\bigl(2\beta tx_3-\frac{2}{3}\beta^2t^3\bigr)\Bigr). \tag{5.5}$$

Therefore

(5.6) After standard rescaling $x\mapsto \mu x$, $t\mapsto \mu t$ $U(x,x,t)$ in comparison with $U_{(2)}(x',x',t)$ acquires factor

$$\frac{1}{2}\mu(2\pi\hbar t)^{-\frac12}\exp\Bigl(i\hbar^{-1}\bigl(2\beta tx_3-\frac{2}{3}\mu^{-2}\beta^2t^3\bigr)\Bigr) \tag{5.7}$$

and therefore (1.23) and (1.24) are replaced respectively by

$$\varphi(t):=-t^2\mu^{-2}\alpha^2\cot(t)+t\mu^{-2}\alpha^2+\frac{2}{3}\mu^{-2}\beta^2t^3-t\tau \tag{5.8}$$

and

$$\mu^{-2}\alpha^2(t^2-t\sin(2t))+2\mu^{-2}\beta^2t^2\sin^2(t)=(\tau-\mu^{-2}\alpha^2)\sin^2(t); \tag{5.9}$$

the latter equation could be rewritten as (1.24) with τ replaced by τ'

$$\mu^{-2}\alpha^2(t^2-t\sin(2t))=(\tau'-\mu^{-2}\alpha^2)\sin^2(t) \tag{5.10}$$

and

$$2\mu^{-2}\beta^2t^2=(\tau-\tau'); \tag{5.11}$$

the former equation shows the return times of 2D-movement on the energy level τ' and the former the looping time of 1D-movement associated with B (after rescaling) on the energy level $(\tau-\tau')$.

Remark 5.1.1. We should set $T = \epsilon_0\mu$ (in contrast to 2D-case). Really, unless we want to make assumptions outside of the ball $B(0,\ell)$ with a small constant ℓ we need to take $T \le \epsilon_0$ (before rescaling) and then $T \le \epsilon_0\mu$ (after rescaling). This absolves us from the analysis of the equatorial zone.

Let us compare solutions t^*_k of (5.9) and solutions t_k to (1.24). One can see easily that

(5.12) As $|k| \le \epsilon|\beta|^{-1}\mu$

$$t^*_k = t_k\bigl(1 + O(\mu^{-2}\beta^2k^2)\bigr) \tag{5.13}$$

and

$$\varphi''(t^*_k) = \varphi''_{(2)}(t_k)\bigl(1 + O(\mu^{-2}\beta^2k^2)\bigr) \tag{5.14}$$

with $\varphi_{(2)}$ defined by (1.23). Furthermore other properties of $\varphi_{(2)}$ are fulfilled for φ as well.

5.2 Tauberian estimates

Then as $|k| \le \epsilon\mu^{-1}$ in virtue of (5.6) and calculations of section 1 contribution of k-th tick to $F_{t\to\hbar^{-1}\tau}\Gamma_x U$ does not exceed

$$C\left\{\begin{aligned} &\mu h^{-1} \times \bigl(\mu/h|k|\bigr)^{\frac{1}{2}} &&\text{as}\ \ 1 \le |k| \le \bar{k},\\ &\mu h^{-1}(\mu^2h/\alpha|k|)^{\frac{1}{2}} \times \bigl(\mu/h|k|\bigr)^{\frac{1}{2}} &&\text{as}\ \ \bar{k} \le |k| \le \epsilon\mu^{-1},\end{aligned}\right. \tag{5.15}$$

where as before $\bar{k} = \epsilon\mu^2h\alpha^{-1}$. Recall that without an extra factor $\bigl(\mu/h|k|\bigr)^{\frac{1}{2}}$ we would have estimate for $F_{t\to\hbar^{-1}\tau}\Gamma_{x'}U_{(2)}$.

However, this extra factor is a game changer. As before there are three cases:

(a) $\alpha \ge \mu^2h$; then we set $\bar{k} = 1$ and the summation with respect to $k : 1 \le |k| \le \epsilon\mu\ell$ returns

$$C\mu h^{-1}(\mu^2h/\alpha)^{\frac{1}{2}} \cdot (\mu/h)^{\frac{1}{2}} \log(\mu\ell) \asymp C\mu^{\frac{5}{2}}h^{-1}\alpha^{-\frac{1}{2}} \log(\mu\ell);$$

(b) $\mu h\ell^{-1} \le \alpha \le \mu^2 h$; then $1 \le \bar{k} \le \epsilon\mu\ell$ and summation with respect to $k : \bar{k} \le |k| \le \epsilon\mu\ell$ returns

$$C\mu^{\frac{5}{2}} h^{-1}\alpha^{-\frac{1}{2}} \log(\mu\ell)/\bar{k} \asymp C\mu^{\frac{5}{2}} h^{-1}\alpha^{-\frac{1}{2}}\bigl(1 + (\log(\alpha\ell/\mu h))_+\bigr)$$

while summation with respect to $k : 1 \le k| \le \bar{k}$ returns the same expression albeit without logarithmic factor;

(c) $\alpha \le \mu h\ell^{-1}$; then $\bar{k} \ge \epsilon\mu\ell$ and we reset set $\bar{k} = \epsilon\mu\ell$ and summation with respect to k returns

$$C\mu h^{-1} \times \bigl(\mu/h\bigr)^{\frac{1}{2}}\bar{k}^{\frac{1}{2}} \asymp C\mu^2 h^{-\frac{3}{2}}\ell^{\frac{1}{2}}.$$

So we have proved

Proposition 5.2.1. *(cf. proposition 1.2.4). (i) After rescaling as* $T = \epsilon\mu\ell$, $\ell \ge C_0\mu^{-1}$

$$|F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)\Gamma_x \mathrm{U}| \le C\mu h^{-2} + C\left\{\begin{aligned} &\mu^{\frac{5}{2}} h^{-1}\alpha^{-\frac{1}{2}}\bigl(1 + (\log(\alpha\ell/\mu h))_+\bigr) && \text{as}\quad \alpha\ell \ge \mu h,\\ &\mu^2 h^{-\frac{3}{2}}\ell^{\frac{1}{2}} && \text{as}\quad \alpha\ell \le \mu h;\end{aligned}\right. \tag{5.16}$$

(ii) Therefore as operator A *coincides with the pilot-model operator* (5.1) *in* $B(0,\ell)$

$$|e(0,0,\tau) - e^{\mathsf{T}}(0,0,\tau)| \le Ch^{-2}\ell^{-1} + C\left\{\begin{aligned} &\mu^{\frac{3}{2}} h^{-1}\alpha^{-\frac{1}{2}}\bigl(1 + (\log(\alpha\ell/\mu h))_+\bigr)\ell^{-1} && \text{as}\quad \alpha\ell \ge \mu h,\\ &\mu h^{-\frac{3}{2}}\ell^{-\frac{1}{2}} && \text{as}\quad \alpha\ell \le \mu h;\end{aligned}\right. \tag{5.17}$$

(iii) In particular, as $\alpha \asymp \ell \asymp 1$ *the right-hand expression in* (5.17) *does not exceed* $Ch^{-2} + C\mu^{\frac{3}{2}} h^{-1}\bigl(1 + |\log \mu h|\bigr)$ *which is* $O(h^{-2})$ *as* $\mu \le (h|\log h|)^{-\frac{2}{3}}$.

5.3 Weyl estimates

Even simpler are Weyl estimates: in comparison with 2D-case they acquire factor $\mu^{\frac{1}{2}} h^{-\frac{1}{2}}$ and (1.64) becomes

$$\mathsf{R}^{\mathsf{W}}_x := |e^{\mathsf{T}}(x,x,0) - h^{-3}\mathcal{N}^{\mathsf{W}}_x| \le C\mu^{-\frac{1}{2}} h^{-\frac{3}{2}} + C\mu^{\frac{5}{2}}\alpha^{-\frac{1}{2}} h^{-1} \tag{5.18}$$

and more generally we arrive to statement (i) below

Proposition 5.3.1. *(cf. proposition 1.3.2). For a pilot-model operator* (5.1) *with* $\tau \asymp 1$,

(i) As $\mu^2 h \le \alpha \le 1$ *estimate* (5.18) *holds; moreover if we replace* $\mathcal{N}_x^{\mathsf{W}}$ *by* $\mathcal{N}_x^{\mathsf{W}} + \mathcal{N}_{x,\mathsf{corr}(r)}$ *with the correction term* $h^{-3}\mathcal{N}_{x,\mathsf{corr}(r)}$ *delivered by* r*-term stationary phase approximation then an error does not exceed*

$$C\mu^{-\frac{1}{2}} h^{-\frac{3}{2}} + C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} \bigl(\mu^2 h/\alpha\bigr)^{r+\frac{1}{2}}; \tag{5.19}$$

(ii) As $\alpha \le \mu^2 h$ *estimate holds:*

$$\mathsf{R}_x^{\mathsf{W}} := |e^{\mathsf{T}}(x,x,0) - h^{-3}\mathcal{N}_x^{\mathsf{W}}| \le C\mu^{\frac{3}{2}} h^{-\frac{3}{2}}; \tag{5.20}$$

(iii) In particular, as $\mu \le h^{-\frac{1}{3}}$ *we have* $\mathsf{R}^{\mathsf{W}} = O(h^{-2})$ *without any assumptions to* α, β*:* $|\alpha| \le 1$, $|\beta| \le 1$.

Recall that here we do not need to analyze the equatorial zone and therefore the difference between cases $\alpha \ge \mu\hbar^{\frac{2}{3}}$ and $\alpha \le \mu\hbar^{\frac{2}{3}}$ disappears together with corresponding terms in (1.66) or (1.67).

5.4 Micro-averaging

Micro-averaging adds yet another layer of complexity. Let us consider an *isotropic micro-averaging* with function $\psi_\gamma(x) = \psi(x/\gamma)$ and an *anisotropic micro-averaging* with function $\psi_{\boldsymbol\gamma}(x) = \psi(x'/\gamma, x_3/\gamma_3)$ where $\boldsymbol\gamma = (\gamma, \gamma_3)$ is a scale with respect to (x', x_3).

Then one needs to integrate by parts with respect to x_3 taking in account factor $\exp\bigl(2ih^{-1}\beta t x_3\bigr)$ in (5.5) and rescaling with respect to t which brings factor $(\mu h/|\beta|\gamma_3|k|)^l$ as $(\mu h/|\beta|\gamma_3|k|) \le 1$ in addition to factor $(\mu h/|\alpha|\gamma|k|)^l$ as $(\mu h/|\alpha|\gamma|k|) \le 1$. Thus

(5.21) Micro-averaging brings factor $(\mu h/\nu(\boldsymbol\gamma)|k|)^l$ as $(\mu h/\nu(\boldsymbol\gamma)|k|) \le 1$ with $\nu(\boldsymbol\gamma) := |\alpha|\gamma + |\beta|\gamma_3$.

Then we need to modify (1.53) replacing $|\alpha|\gamma$ by $\nu(\boldsymbol\gamma)$:

$$|k| \ge \hat{k}(\boldsymbol\gamma) := \mu h/\nu(\boldsymbol\gamma). \tag{5.22}$$

To understand how it affects a Tauberian estimate we need to consider the following cases:

(a) Micro-averaging has no effect as $\hat{k}(\gamma)\ge \epsilon\mu\ell$ i.e. as $\nu(\gamma)\le h\ell^{-1}$;

(b) Micro-averaging has a minimal effect (only decreasing the logarithmic factor) as $\epsilon\mu\ell\ge\hat{k}(\gamma)\ge\bar{k}$. Obviously in this case $\bar{k}=\epsilon\max\bigl(\mu^2h/\alpha,1\bigr)$.

So this case holds iff $h\ell^{-1}\le\nu(\gamma)\le\min(\mu h,\alpha\mu^{-1})$. Then the logarithmic factor becomes $\log\bigl(\hat{k}(\gamma)/\bar{k}\bigr)$.

(c) Micro-averaging kills logarithmic factor and further leads to $\bar{k}^{\frac{1}{2}}$ replaced by $\hat{k}(\gamma)^{\frac{1}{2}}$ as $1\le\hat{k}(\gamma)\le\bar{k}$. Obviously in this case $\bar{k}=\epsilon\min\bigl(\mu^2h/\alpha,\mu\ell\bigr)$.

So this case holds iff $\max(h\ell^{-1},\alpha\mu^{-1})\le\nu(\gamma)\le\mu h$. Then the right-hand expression in (5.24) becomes $C\mu^2h^{-1}\nu^{-\frac{1}{2}}\gamma^{-\frac{1}{2}}$;

(d) Micro-averaging resets $\bar{k}$ to 1 and brings factor $(\mu h/\nu(\gamma))^l$ as $\hat{k}(\gamma)\le 1$ i.e. as $\nu(\gamma)\ge\mu h$. Then the right-hand expression in (5.24) becomes $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}(\mu h/\nu(\gamma))^l$.

Thus we arrive to

Proposition 5.4.1. *For the pilot-model operator* (5.1) *with* $\tau\asymp 1$, $0\le\alpha\le 1$, $|\beta|\le 1$, $\mu\le h^{-1}$ *in* $\mathbb{R}^3$ *estimates*

$$\gamma^{-2}\gamma_3^{-1}|F_{t\to\hbar^{-1}\tau}\bigl(\bigl(\bar{\chi}_T(t)-\bar{\chi}_{\bar{T}}\bigr)\Gamma(U\psi_\gamma)\bigr)|\le C\mu R^{\mathsf{T}}(\gamma)\tag{5.23}$$

and

$$\gamma^{-2}\gamma_3^{-1}|F_{t\to\hbar^{-1}\tau}\bigl(\bar{\chi}_T(t)\Gamma(U\psi_\gamma)\bigr)|\le C\mu h^{-2}+C\mu R^{\mathsf{T}}(\gamma)\tag{5.24}$$

hold as $T=\epsilon\mu\ell$ *where*

$$R^{\mathsf{T}}(\gamma)=\begin{cases}\mu^{\frac{3}{2}}h^{-1}\alpha^{-\frac{1}{2}}\bigl(1+(\log(\alpha\ell/\mu h))_+\bigr) & \textit{as}\ \ \alpha\ell\ge\mu h,\ \ \nu(\gamma)\le h\ell^{-1},\\ \mu h^{-\frac{3}{2}}\ell^{\frac{1}{2}} & \textit{as}\ \ \alpha\ell\le\mu h,\ \ \nu(\gamma)\le h\ell^{-1},\\ \mu^{\frac{3}{2}}h^{-1}\alpha^{-\frac{1}{2}}\bigl(1+(\log(\alpha/\mu\nu(\gamma)))_+\bigr) & \textit{as}\ \ h\ell^{-1}\le\nu(\gamma)\le\min(\mu h,\alpha\mu^{-1}),\\ \mu h^{-1}\nu(\gamma)^{-\frac{1}{2}} & \textit{as}\ \ \max(h\ell^{-1},\alpha\mu^{-1})\le\nu(\gamma)\le\mu h,\\ \mu^{\frac{1}{2}}h^{-\frac{3}{2}}(\mu h/\nu(\gamma))^l & \textit{as}\ \ \nu(\gamma)\ge\mu h.\end{cases}\tag{5.25}$$

In the right-hand expression of (5.25) two first lines correspond to the case (a).

Corollary 5.4.2. *In frames of proposition 5.4.1 the following estimate holds*

$$\gamma^{-2}\gamma_3^{-1}|\Gamma\bigl(e(.,.,\tau)-e^{\mathsf{T}}(.,.,\tau)\bigr)\psi_\gamma| \le C\bigl(h^{-2}+R^{\mathsf{T}}(\gamma)\bigr)\ell^{-1} \tag{5.26}$$

Consider now how micro-averaging affects Weyl estimate (with the right-hand expression (5.19)). There is no effect as $\nu(\gamma)\le \mu h$ and factor $(\mu h/\nu(\gamma))^l$ comes out as $\nu(\gamma)\ge \mu h$. Thus we arrive to

Proposition 5.4.3. *For the pilot-model operator* (5.1) *with* $\tau\asymp 1$, $0\le \alpha\le 1$, $\mu\le h^{-1}$ *in* $\mathbb{R}^3$

(i) As $\mu^2 h\le \alpha\le 1$ *estimate holds:*

$$\begin{aligned}
&\mathsf{R}^{\mathsf{W}}_{x(r)}(\gamma) :=\\
&\qquad \gamma^{-2}\gamma_3^{-1}|\int \Bigl(e^{\mathsf{T}}(x,x,0)-h^{-3}\bigl(\mathcal{N}_x^{\mathsf{W}}+\mathcal{N}_{x,\mathsf{corr}(r)}\bigr)\Bigr)\psi_\gamma\,dx| \le\\
&\qquad Ch^{-2}+C\mu^{\frac32}h^{-\frac32}\bigl(\mu^2h/\alpha\bigr)^{r+\frac12}\begin{cases}(\mu h/\nu(\gamma))^l & \text{as}\ \ \nu(\gamma)\ge\mu h,\\ 1 & \text{as}\ \ \nu(\gamma)\le \mu h\end{cases}
\end{aligned}\tag{5.27}$$

(ii) As $\alpha\le \mu^2 h$ *estimate holds:*

$$\begin{aligned}
\mathsf{R}^{\mathsf{W}}_{x}(\gamma) := \gamma^{-2}\gamma_3^{-1}|\int \Bigl(e^{\mathsf{T}}(x,x,0)-h^{-3}\mathcal{N}_x^{\mathsf{W}}\Bigr)\psi_\gamma\,dx| \le&\\
Ch^{-2}+C\mu^{\frac32}h^{-\frac32}\begin{cases}(\mu h/\nu(\gamma))^l & \text{as}\ \ \nu(\gamma)\ge\mu h,\\ 1 & \text{as}\ \ \nu(\gamma)\le \mu h.\end{cases}&
\end{aligned}\tag{5.28}$$

5.5 Strong and superstrong magnetic field

5.5.1 Tauberian estimate

Consider strong $\mu h\asymp 1$ and superstrong $\mu h\gtrsim 1$ magnetic field for the Schrödinger-Pauli pilot-model operator

$$A=\bar{A} := h^2D_1^2+(hD_2-\mu x_1)^2+h^2D_3^2+2\alpha x_1+2\beta x_3-\mu h \tag{5.29}$$

which alternatively means that we assume that $|\tau-\mathfrak{z}\mu h|\le C$ in the framework of the standard pilot-model (5.1), $\mathfrak{z}=1$.

Proposition 5.5.1. *For a pilot-model operator* (5.1) *with* $\mu h \gtrsim 1$ *in* $\mathbb{R}^3$ *as* $|x| \le C_0$, $|y| \le C_0$, $\tau \le c\mu h$

(i) $e(x,y,\tau) \equiv 0 \mod O(\mu h^\infty)$ *as* $\tau \le \mu h - \epsilon_0$ *(lower spectral gap);*

(ii) As $\alpha > 0$ *the following estimates hold*

$$|e(x,x,\tau+h) - e(x,y,\tau)| \le C\mu h^{-1} \max_{j\in\mathbb{Z}^+}\bigl(\max(|V(x)+2j\mu h-\tau|, h)\bigr)^{-\frac12} \tag{5.30}$$

and

$$|F_{t\to\hbar^{-1}\tau} U(x,y,t)| \le C\mu^2 h^{-1} \max_{j\in\mathbb{Z}^+}\bigl(\max(|V(x)+2j\mu h-\tau|, h)\bigr)^{-\frac12} \tag{5.31}$$

and these estimate are sharp as $x = y$ *and* τ *is close to Landau level* μh. *Recall that in* (5.31) t *is rescaled.*

Proof. Statement (i) is obvious. To prove (ii) note that

$$e(x,y,\tau) = \int e_B(x_3,y_3,\tau-\tau')\, d_{\tau'} e_{(2)}(x',y',\tau') = \int e_{(2)}(x',y',\tau-\tau')\, d_{\tau'} e_B(x_3,y_3,\tau') \tag{5.32}$$

and therefore

$$e(x,y,\tau+h) - e(x,y,\tau) = \int \bigl(e_B(x_3,y_3,\tau-\tau'+h) - e_B(x_3,y_3,\tau-\tau')\bigr)\, \partial_{\tau'} e_{(2)}(x',y',\tau')\, d\tau'. \tag{5.33}$$

Recall that due to our analysis in section 1 $\partial_{\tau'} e_{(2)}(x',y',\tau')$ is essentially supported in $C\mu^{-\frac12}h^{\frac12}\alpha$-vicinity of Landau level and fast decays out of it.

Finally recall that due to the analysis of subsubsections 5.2.1–5.2.1

$$|e_B(x_3,y_3,\tau+h) - e_B(x_3,y_3,\tau)| \le C(\tau')^{-\frac12} \qquad \text{as } \tau' \ge C_0 h. \tag{5.34}$$

Combining it with estimate (1.73) for $|\partial_{\tau'} e_{(2)}(x',y',\tau')|$ we conclude that the left-hand expression of (5.30) does not exceed

$$C\mu^{\frac32}\alpha^{-1}h^{-\frac32}) \times \alpha\mu^{-\frac12}h^{\frac12} \times \max_{j\in\mathbb{Z}^+}\bigl(\max(|V(x)+2j\mu h-\tau|, h)\bigr)^{-\frac12}.$$

Estimate (5.31) is proven; estimate (5.31) follows from it. ☐

Further, as before $T^* = \epsilon_0\mu\ell$. Then we immediately arrive to

Corollary 5.5.2. *For a self-adjoint operator in domain* X*,* $B(0,\ell) \subset X \subset \mathbb{R}^2$*,* $\ell \ge C_0 h$*, coinciding in* $B(0,\ell)$ *with the pilot-model operator* (5.29) *with* $\tau \asymp 1$*,* $0 < \alpha \le 1$*,* $|\beta| \le 1$*,* $\mu \ge h^{-1}$

(i) Statements (i)-(ii) of proposition 5.5.1 remain true;

(ii) Formula (5.32) *holds modulo* $O(\mu h^{-\frac{3}{2}}\ell^{-1})$.

5.5.2 Micro-averaging

Let us consider micro-averaging. First let us estimate Fourier transform where as before we rescaled $t \mapsto \mu t$; formulae (5.16)–(5.17) imply immediately

Proposition 5.5.3. *For a pilot-model operator* (5.29) *in* $\mathbb{R}^3$ *with* $\mu h \ge 1$*,* $\gamma_3 \ge h$

$$\gamma^{-2}\gamma_3^{-1}|F_{t\to h^{-1}\tau}\bar{\chi}_T(t)\Gamma(U\psi_\gamma)| \le C\mu^2 h^{-1}\gamma_3^{-\frac{1}{2}} \tag{5.35}$$

Corollary 5.5.4. *In frames of corollary 5.5.2 as* $\ell \ge \gamma_3 \ge h$

$$\gamma^{-2}\gamma_3^{-1}|\Gamma\bigl(e(.,.,\tau) - e^{\mathsf{T}}(.,.,\tau)\bigr)\psi_\gamma| \le C\mu h^{-1}\gamma_3^{-\frac{1}{2}}\ell^{-1}. \tag{5.36}$$

5.6 Magnetic Weyl approximation

We can try to use a host of the different approximations but restrict ourselves now to the magnetic Weyl approximation. Recall (5.32) and note that

$$e^{\mathsf{MW}}(x,y,\tau) = \int e^{\mathsf{W}}_B(x_3,y_3,\tau-\tau')\,d_{\tau'}e^{\mathsf{MW}}_{(2)}(x',y',\tau') = \\ \int e^{\mathsf{MW}}_{(2)}(x',y',\tau-\tau')\,d_{\tau'}e^{\mathsf{W}}_B(x_3,y_3,\tau'). \tag{5.37}$$

5.6.1 Pointwise asymptotics

Proposition 5.6.1. *For a pilot-model operator* (5.1) *with* $\tau \le c$, $\mu h \lesssim 1$ *and for a pilot-model operator* (5.29) *with* $\tau \le c$, $\mu h \gtrsim 1$

$$|e^{\mathsf{MW}}(x,y,\tau) - e^{\mathsf{MW}}(x,y,\tau')| \le C \begin{cases} \mu^{\frac{3}{2}} h^{-1} & \text{as } \mu h \le 1,\\ \mu h^{-\frac{3}{2}} & \text{as } \mu h \ge 1 \end{cases} \tag{5.38}$$

as $|\tau - \tau'| \le h$.

Proof. Obviously the left-hand expression does not exceed

$$C\mu h^{-\frac{3}{2}} \sum_{j\le J} j^{-\frac{1}{2}} \asymp C\mu h^{-\frac{3}{2}} J^{\frac{1}{2}}$$

with $J = C_0 \max((\mu h)^{-1}, 1)$ which implies (5.38). □

Therefore one can hardly expect that the magnetic Weyl approximation provides a better error than the right hand expression of (5.38).

Proposition 5.6.2. *Let* $|\tau| \le \epsilon$. *Then for a pilot-model operator* (5.1) *as* $\mu h \lesssim 1$ *and for a pilot-model operator* (5.1) *as* $\mu h \gtrsim 1$

$$\mathsf{R}^{\mathsf{MW}} := |e^{\mathsf{T}}(x,x,0) - h^{-3}\mathcal{N}^{\mathsf{MW}}| \le Ch^{-2} + C\mu h^{-1} + C\mu h^{-\frac{5}{3}} |\beta|^{\frac{1}{3}}. \tag{5.39}$$

Proof. Without any loss of the generality we can assume that $x = 0$, $\beta > 0$. As contribution of k-th tick to the Tauberian expression is $O(\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}})$, its contribution to the error when we replace β by 0 does not exceed $C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}} \times \mu^{-3} h^{-1} \beta^2 |k|^3$ and summation with respect to

$$k : |k| \le \tilde{k}_1 := \mu \beta^{-\frac{2}{3}} h^{\frac{1}{3}} \tag{5.40}$$

returns

$$C\mu^{-\frac{3}{2}} h^{-\frac{5}{2}} \beta^2 |k|^{\frac{5}{2}} = C\mu h^{-\frac{5}{3}} \beta^{\frac{1}{3}} \tag{5.41}$$

On the other hand, summation of $C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}}$ as $|k| \ge \tilde{k}_1$ returns $C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} \tilde{k}_1^{-\frac{1}{2}}$ which is the same expression because $\tilde{k}_1$ was defined from $\mu^{-3} h^{-1} \beta^2 k^3 = 1$.

Meanwhile, contribution of k-th tick to the error when we replace α by 0 does not exceed $C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}} \times \mu^{-2} h^{-1} \alpha |k|$ and summation with respect to

$$k : |k| \le \tilde{k}_2 := \mu^2 h \alpha^{-1} \tag{5.42}$$

returns

$$C\mu^{-\frac{1}{2}}h^{-\frac{5}{2}}\alpha\tilde{k}_2^{\frac{1}{2}} = C\mu^{-\frac{1}{2}}h^{-2}\alpha^{\frac{1}{2}} \tag{5.43}$$

which is less than Ch^{-2}. On the other hand, summation of $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}|k|^{-\frac{3}{2}}$ as $|k| \ge \tilde{k}_2$ returns $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}\tilde{k}_2^{-\frac{1}{2}}$ which is the same expression because $\tilde{k}_2$ was defined from equation $\mu^{-2}h^{-1}\alpha\tilde{k}_2 = 1$. $\square$

Remark 5.6.3. (i) We summed with respect to all k while it would be enough only with respect to $k : |k| \le \epsilon\mu$. However it provides us by no improvement.

(ii) So far we have not used factor $C(\mu^2h/\alpha|k|)^{\frac{1}{2}}$; using it we acquire in (5.41) factor $(\mu^2h/\alpha\tilde{k}_1)^{\frac{1}{2}} = \mu^{\frac{1}{2}}h^{\frac{1}{3}}\beta^{\frac{1}{3}}\alpha^{-\frac{1}{2}}$ resulting in

$$\mathsf{R}^{\mathsf{MW}} \le Ch^{-2} + C\mu h^{-1} + C\mu^{\frac{3}{2}}h^{-\frac{4}{3}}\beta^{\frac{2}{3}}\alpha^{-\frac{1}{2}} \tag{5.44}$$

provided

$$\alpha \ge \mu h^{\frac{2}{3}}\beta^{\frac{2}{3}}. \tag{5.45}$$

(iii) In contrast to 2D the magnetic Weyl approximation now is better than Weyl approximation as $\mu \ge h^{-\frac{1}{3}}$ (when it matters) but Weyl approximation with the correction terms may be better still.

5.6.2 Micro-averaging

Consider now micro-averaging. We need to redo only the first step of the proof of proposition 5.6.2 and only in the case $\mu \ge h^{-\frac{1}{3}}\beta^{-2}$ and it is useful only if $\tilde{k}_1 \ge \hat{k}(\gamma) = \mu h/\nu(\gamma)$ or equivalently

$$\nu(\gamma) = \alpha\gamma + \beta\gamma_3 \ge \beta^{\frac{2}{3}}h^{\frac{2}{3}}. \tag{5.46}$$

In this case in the left-hand expression of (5.41) should be reset to $\tilde{k}_1$ replaced by $\hat{k}(\gamma)$ as $\hat{k}(\gamma) \ge 1$ or by 1 with an extra factor $(\mu h/\nu(\gamma))^l$ as $\hat{k}(\gamma) \le 1$ resulting in $CR_1^{\mathsf{MW}}(\gamma)$ with

$$R_1^{\mathsf{MW}}(\gamma) := \\
C\mu\begin{cases}
\beta^2\nu(\gamma)^{-\frac{5}{2}}\bigl(\mu h/\nu(\gamma)\bigr)^l & \text{as } \nu(\gamma) \ge \mu h,\ \alpha \le \mu^2 h,\\
\beta^2\nu(\gamma)^{-\frac{5}{2}} & \text{as } \nu(\gamma) \le \mu h,\ \mu\nu(\gamma) \ge \alpha,\\
\beta^2\nu(\gamma)^{-\frac{5}{2}}(\mu^2h/\alpha)^{\frac{1}{2}}\bigl(\mu h/\nu(\gamma)\bigr)^l & \text{as } \nu(\gamma) \ge \mu h,\ \alpha \ge \mu^2 h,\\
\beta^2\nu(\gamma)^{-\frac{5}{2}}(\mu\nu(\gamma)/\alpha)^{\frac{1}{2}} & \text{as } \nu(\gamma) \le \mu h,\ \mu\nu(\gamma) \le \alpha
\end{cases} \tag{5.47}$$

and therefore we arrive to

Proposition 5.6.4. *Let* $|\tau| \le \epsilon$ *and* (5.46) *be fulfilled. Then for a pilot-model operator* (5.1) *as* $\mu h \lesssim 1$ *and for a pilot-model operator* (5.1) *as* $\mu h \gtrsim 1$ *a*

$$\mathsf{R}^{\mathsf{MW}}(\gamma) := \gamma^{-2}\gamma_3^{-1}\left|\int \bigl(e^{\mathsf{T}}(x,x,0) - h^{-3}\int \mathcal{N}^{\mathsf{MW}}\bigr)\psi_\gamma\, dx\right| \le Ch^{-2} + C\mu h^{-1} + CR_1^{\mathsf{MW}}(\gamma) \tag{5.48}$$

with $R_1^{\mathsf{MW}}(\gamma)$ *defined by* (5.47).

Remark 5.6.5. If we replace only $e_{(2)}(.,.,.)$ by $e^{\mathsf{MW}}(.,.,.)$ we can skip the last term.

5.7 Geometric interpretation

Note that there are two classical dynamics: in x' and in x_1. The former has return times $t'_k := t'_k(\tau')$ where τ' is a corresponding part of energy and t'_k are defined in section 1 as $t_k(\tau)$ while the latter is

$$x_3(t) = x_3(0) + 2\xi_3(0)t - \beta t^2, \qquad \xi_3(t) = \xi_3(0) - \beta t. \tag{5.49}$$

and has return times (to 0) $t''_k(\tau-\tau') = 2\beta^{-1}(\tau-\tau')$ and therefore the total system has return times[16] t_k and return energy partitions $(\tau'_k, \tau-\tau'_k)$ defined from the pair of equations

$$t_k = t'_k(\tau'_k) = t''_k(\tau-\tau') \tag{5.50}$$

which is equivalent to (5.9)–(5.10).

Therefore relatively sets of return directions ξ_k is thinner in 3D than in 2D which explains errors acquiring lesser factor (in comparison with the principal part) in 3D than in 2D.

---

[16] To point 0.

In particular, as $\beta = 0$ 1D-dynamics does not return at all unless it stays at 0, therefore in this case (5.49) mean exactly that $\tau'_k = \tau$ and $t'_k(\tau) = t_k(\tau)$.

6 Pointwise asymptotics: general 3D-operators

6.1 Set-up

We assume that

$$F := \bigl(\sum_{jk} g_{jk} F^j F^k\bigr)^{\frac{1}{2}} \tag{6.1}$$

where $\mathbf{F} = (F^1, F^2, F^3)$ and F are vector and scalar intensities of the magnetic field respectively. Temporarily we assume that $F = 1$ (as we can reduce the general case to this one by dividing operator by F and using our standard arguments).

Further, without any loss of the generality we will assume locally that

$$F^1 = F^2 = 0. \tag{6.2}$$

Remark 6.1.1. (i) Due to Frobenius theorem we can make locally $g^{3j} = 0$ for $j = 1, 2$ and $F^1 = F^2 = 0$ simultaneously if and only if $f_* \wedge df_* = 0$ where $f_* = \sum_{j,k} F^j g_{jk} dx^k$ is 1-form.

(ii) As

$$F = \bigl(\frac{1}{2} \sum_{i,j,k,l} g^{ij} g^{kl} F_{ik} F_{jl}\bigr)^{\frac{1}{2}}, \qquad F_{jk} = \partial_k A_j - \partial_j A_k, \tag{6.3}$$

under assumption (6.2)

$$F = \bigl(g^{11} g^{22} - g^{12} g^{21}\bigr)^{\frac{1}{2}} |F_{12}|. \tag{6.4}$$

However

(6.5) We can assume that $F^1 = F^2 = 0$ locally and simultaneously that $g^{jk} = \delta_{jk}$ along a single magnetic line $\{x' = 0\}$.

Really, we can satisfy condition $f_* \wedge df_* = 0$ changing g^{jk} but preserving them at the chosen magnetic line. Note that F then acquires some scalar factor which does not affect this condition. So, we can assume $g^{3j} = 0$ for $j = 1, 2$ and $F^1 = F^2 = 0$ at the chosen magnetic line which (after shift) becomes then $\{x' = 0\}$. Changing $x_3 \mapsto \phi(x_3)$ makes $g^{33} = 1$ along this magnetic line. Then $F^3 = \pm F = 1$ (for an appropriate orientation). Changing $x' = B(x_3)x'$ with an appropriate matrix $B(x')$ makes $g^{jk} = \delta_{jk}$ along this line.

Further

(6.6) In frames of assumption (6.5) we can assume that $g^{31} = 0$.

Really, one can achieve it by $x' \mapsto x'$, $x_3 \mapsto \phi(x)$ with $\phi(x) = x_3 + O(|x'|^2)$.

Further, one can make

$$g^{jk} = \delta_{jk} + O(|x'|^2) \qquad \text{as } \; x_3 = 0 \quad j, k = 1, 2 \tag{6.7}$$

by $x' \mapsto x' + Q(x')$ with Q quadratic respect to x'.

By a gauge transform one can make $A_3 = 0$. Then $F^1 = F^2 = 0$ imply that A_1, A_2 do not depend on x_3 and therefore we can assume that $A_1 = 0$ as well:

$$A_1 = A_3 = 0, \qquad A_2 = A_2(x') = x_2 + O(|x'|^3) \tag{6.8}$$

where the last equality is due to $F = 1$ and assumption (6.5).

6.2 Classical dynamics

Let us consider a classical dynamics. We know that for time $|t| \le T = \epsilon_0$ Hamiltonian trajectory starting from $\{|x| \le c_0\mu^{-1}\}$ remains confined in $C_0\mu^{-1}$-tube $\{|x'| \le C_0\mu^{-1}\}$. Then in virtue of our assumptions

$$\frac{dx_3}{dt} = 2\xi_3 + O(\mu^{-1}) \qquad \frac{d\xi_3}{dt} = -V_{x_3}(0, 0, x_3) + O(\mu^{-1}) \tag{6.9}$$

and therefore

$$(x_3, \xi_3)(t) = (x_3^0, \xi_3^0)(t) + O(\mu^{-1}|t|) \tag{6.10}$$

where here $(x_3^0,\xi_3^0)(t)$ denotes dynamics for a 1D-Schrödinger $\xi_3^2+V(0,0,x_3)$ with the same initial data (as $t=0$).

Then one can prove easily that

$$|\xi_3(t)|\le C\bigl(|\beta|\cdot|t|+|\xi_3(0)|\bigr),\qquad |x_3|\le C\bigl(|\beta|t^2+|\xi_3(0)|\cdot|t|\bigr)\\ \text{if}\ \ x_3(0)=0\ \ \text{and}\quad |t|\le\epsilon \tag{6.11}$$

with

$$\beta:=-V_{x_3}(0). \tag{6.12}$$

Then

$$\xi_3=\xi_3(0)\bigl(1+O(t^2)\bigr)+2\beta t\bigl(1+O(t^2)\bigr), \tag{6.13}$$

$$x_3=\xi_3(0)t\bigl(1+O(t^2)\bigr)+\beta t^2\bigl(1+O(t^2)\bigr). \tag{6.14}$$

Let us analyze evolution of $(x',\xi')(t)$ but we start from $(p_1,p_2)(t)$. Recall that

$$\mu^{-1}\{p_1,p_2\}\equiv 1,\qquad g^{11}g^{22}-g^{12}g^{21}\equiv 1\mod O(\mu^{-2}) \tag{6.15}$$

in the tube in question.

Let $p_1':=p_1+\gamma p_3$; then $p_2':=(g^{12}p_1+g^{22}p_2+g^{23}p_3)\equiv -\frac{1}{2}\mu^{-1}\{a,p_1'\}$ $\mod O(\mu^{-1})$. Then

$$\frac{1}{2}\mu^{-1}\{a,p_2'\}\equiv (g^{11}p_1+g^{12}p_2+g^{13}p_3)g^{22}-(g^{12}p_1+g^{22}p_2+g^{23}p_3)g^{12}\equiv\\ p_1+(g^{13}g^{22}-g^{23}g^{12})p_3.$$

So we pick up $\gamma=g^{13}g^{22}-g^{23}g^{12}$ and then

$$p_1':=(g^{22})^{-\frac{1}{2}}(p_1+\gamma p_3),\qquad \gamma=g^{13}g^{22}-g^{23}g^{12}, \tag{6.16}$$

$$p_2':=(g^{22})^{-\frac{1}{2}}(g^{12}p_1+g^{22}p_2+g^{23}p_3), \tag{6.17}$$

satisfy

$$\frac{1}{2}\mu^{-1}\{a,p_1'\}\equiv -p_2',\qquad \frac{1}{2}\mu^{-1}\{a,p_2'\}\equiv p_1',\qquad \mu^{-1}\{p_1',p_2'\}\equiv 1 \tag{6.18}$$

modulo $O(\mu^{-1})$. We set $p_3'=p_3$.

We do not assume that $g^{13}=0$ identically; instead its choice will be different and it would be only $O(\mu^{-1})$ in the tube in question. Note that

$$\gamma \equiv g^{13}g^{22} \mod O(\mu^{-2}) \equiv g^{13} \mod O(\mu^{-2}+\mu^{-1}|x_3|), \tag{6.19}$$

$$\gamma_{x_j} \equiv g^{13} \mod O(\mu^{-1}). \tag{6.20}$$

Really, $g^{jk}=\delta_{jk}+O(|x'|^2+|x'|\cdot|x_3|)$ $(j=1,2)$, $g^{3j}=O(|x'|)$.

Let us consider again $\{a,p'_1\}$, this time more precisely. Note that

$$\{a,p'_1\}=-2\mu p'_2+2\sum_j \gamma_{x_j}p_jp_3-V_{x_1}+O(\mu^{-1}) \qquad \text{as } x_3=0$$

as $|x'|=O(\mu^{-1})$ due to our assumptions.

Therefore we arrive to the first equation below and the second is proven in the same way

$$\{a,p'_1\}\equiv -2\mu p'_2+2\sum_j g^{13}_{x_j}p_jp_3-V_{x_1}, \tag{6.21}$$

$$\{a,p'_2\}\equiv \quad 2\mu p'_1+2\sum_j g^{23}_{,x_j}p_jp_3-V_{x_2} \mod O(\mu^{-1}). \tag{6.22}$$

We want

$$g^{23}_{x_2}=g^{13}_{x_1}=0,\quad -g^{23}_{x_1}=g^{13}_{x_2}=\kappa(x_3) \qquad \text{as } x'=0 \tag{6.23}$$

which is possible to arrange as the only value at $x'=0$ which is fixed is

$$\kappa(x_3):=\frac{1}{2}\bigl(g^{13}_{x_2}-g^{23}_{x_1}\bigr)\bigr|_{x'=0}=-\frac{1}{2}\bigl(g_{13,x_2}-g_{23,x_1}\bigr)\bigr|_{x'=0}; \tag{6.24}$$

one can prove easily this by transformation $x_3\mapsto x_3+Q(x_3;x')$ with $Q(x_3;x')$ quadratic with respect to x'. Note that

$$f_*\wedge df_*=2\kappa(x_3)dx_1\wedge dx_2\wedge dx_3 \qquad \text{as } x'=0. \tag{6.25}$$

Then (6.21)–(6.22) become

$$\{a,p'_1\}\equiv -2(\mu+\kappa p_3)p'_2-V_{x_1}, \tag{6.26}$$

$$\{a,p'_2\}\equiv \quad 2(\mu+\kappa p_3)p'_1-V_{x_2} \qquad \mod O(\mu^{-1}). \tag{6.27}$$

Obviously

(6.28) We can replace in equations (6.26)–(6.27) V_{x_j} by $V_{x_j}(x_3) := V_{x_j}|_{x'=0}$ and p_3 by $p_3^0(t)$.

If $\kappa = 0$ and V_{x_j} did not depend on x_3 we would get the pilot-model operator. Unfortunately, it is not the case and (6.26), (6.27) coincide with those for the pilot-model only modulo $O(\mu^{-1} + |p_3| + |x_3|)$. However note that $\kappa(x_3)p_3 = \frac{d}{dt}K(x_3)$ where

$$K = \frac{1}{2}\int \kappa \, dx_3. \tag{6.29}$$

Therefore

$$p'(t) \equiv \Omega(t)\Omega^{-1}(0)p'(0) + \int_0^t \Omega(t)\Omega^{-1}(t')V'\,dt \qquad \mod O(\mu^{-1}|t|) \tag{6.30}$$

with

$$\Omega(t) = \begin{pmatrix} \cos(\mu\theta) & -\sin(\mu\theta) \\ \sin(\mu\theta) & \cos(\mu\theta) \end{pmatrix}, \quad \theta(t) = t + \mu^{-1}K(x_3(t)) \tag{6.31}$$

$p = (p_1, p_2)^t$, $V' = (\nabla' V)^t$, $\Omega(t)$ fundamental matrix of system (6.26)–(6.27). Expressing $\Omega^{-1}(t')$ via its derivative and integrating by parts we conclude that

$$p'(t) \equiv \Omega(t)p'(0) + \mu^{-1}\Omega(t)V'(x_3(0)) - \mu^{-1}V'(x_3(t)). \tag{6.32}$$

Meanwhile, as $j = 1, 2$

$$\{a, (-1)^j\mu x_{3-j} + p_j\} \equiv -\sum_{k,l} g^{kl}_{x_j} p_k p_l - V_{x_j} \quad \mod O(\mu^{-1}) \tag{6.33}$$

and we can replace in the right-hand expression x' by 0, p_k by p'_k and as $k = 1, 2$ we can replace p'_k by the corresponding component of $\Omega(t)p'(0)$. Then integrating by parts we arrive to

$$\begin{gathered}\bigl((-1)^j\mu x_{3-j} + p_j\bigr)\big|_{t=0}^{t=t} \equiv \\ -\int_0^t \Bigl(\frac{1}{2}\sum_{k,l} \bigl(g^{11} + g^{22}\bigr)_{x_j}(p_1^2 + p_2^2) + V_{x_j}(x_3(t))\Bigr)\,dt \equiv -\int_0^t V_{x_j}(x_3(t))\,dt\end{gathered} \tag{6.34}$$

modulo $O(\mu^{-1})$ as $\bigl(g^{11} + g^{22}\bigr)_{x_j} = O(\mu^{-1})$ in virtue of our assumptions including (6.15). Further, considering out-of integral terms we conclude that

(6.35) Equality (6.34) holds modulo $O\bigl(\mu^{-1}(|t| + |\sin(2\mu\theta) + |p_3|\bigr)$ (where $p_3(t) \equiv p_3(0) \mod |t|$).

6.3 Semiclassical approximation

So, after rescaling, the classical dynamics of the general operator is close to one of the *generalized pilot-model*

$$\tag{6.36} A^0 := h^2D_1^2 + (hD_2 - \mu x_1)^2 + h^2D_3^2 + V^0(x), \qquad V^0(x) := V(x_3) + \sum_{j=1,2} \alpha_j(x_3)x_j$$

with

$$\tag{6.37} V(x_3) = V(0,0,x_3), \qquad \alpha_j(x_3) = V_{x_j}(0,0,x_3), \quad j = 1,2$$

and the latter one is close albeit with a larger error to the dynamics of the pilot-model (5.1) with $V(x_3)$, $\alpha_j(x_3)$ replaced by $V(0) + \beta x_3$ ($\beta = V_{x_3}(0)$), $\alpha_j = \alpha_j(0)$ respectively.

Proposition 6.3.1. *(cf. proposition 2.2.1). Let conditions* (2.1)–(2.3), (6.7)–(6.8) *be fulfilled*[6]*. Then after rescaling*

(i) Uniformly with respect to $|t| \le \epsilon\mu$ *the propagator* $e^{i\hbar^{-1}tA}$ *is an* $\hbar$*-Fourier integral operator corresponding to the Hamiltonian flow* Ψ_t*;*

(ii) As $|\sin(2\theta_1)| \ge \epsilon$, $|t| \ge \epsilon_0$

$$\tag{6.38} U(x,y,t) \equiv \mu(2\pi\hbar)^{-\frac{3}{2}} t^{-\frac{1}{2}} i(\sin(\theta_1))^{-1} e^{i\hbar^{-1}\phi(x,y,t)} \Bigl(\sum_m b_m(x,y,t)\Bigr)\hbar^m$$

with θ_1 *defined by*

$$\tag{6.39} \theta_1(t) = \theta(t) + K\bigl(x_3(t)\bigr) - K\bigl(x_3(0)\bigr)$$

with $\theta(t)$, $K(x_3)$ *defined by* (2.20), (6.29) *respectively and with* ϕ *defined by* (2.25)–(2.29) *and satisfying (with all derivatives)*

$$\tag{6.40} \phi = \bar{\phi}(\theta) + O(\mu^{-1}t),$$

$$\tag{6.41} b_m = \delta_{0m} + O(\mu^{-1}t)$$

with

$$\bar{\phi} = \beta\mu^{-1}t(x_3 + y_3) + \frac{1}{8}(x_3 - y_3)^2 t^{-1} - \frac{2}{3}\beta^2\mu^{-2}t^3 + \bar{\phi}_2 \tag{6.42}$$

with $\bar{\phi}_2$ *defined by* (1.10).

Proof. Easy but tedious details we leave to the reader: comparing with the pilot-model or generalized pilot model we see that the structure of the canonical graph is close to one of the pilot-model. $\square$

Proposition 6.3.2. *(cf. proposition 2.2.2). Let conditions* (2.1)–(2.3), (6.7)–(6.8) *be fulfilled*6)*. Then after rescaling*

(i) Decomposition (6.39) *remains valid as* $|t| \ge \epsilon_0$ *and* $|\cos(\theta_1)| \le \epsilon$*;*

(ii) Further, after rescaling decomposition (6.39) *remains valid as* $|t| \ge \epsilon_0$ *and*

$$C\max\bigl(\hbar, \mu^{-2}|t|\bigr) \le |\sin(\theta_1)| \le \epsilon \tag{6.43}$$

albeit with an error not exceeding

$$C\hbar^{-\frac{3}{2}}|t|^{-\frac{1}{2}}|\sin(\theta_1)|^{-1}\bigl(\hbar/|\sin(\theta_1)|\bigr)^l \tag{6.44}$$

and with ϕ*,* b_m *such that*

$$|D^\beta(\phi - \bar{\phi})| \le C_\beta\mu^{-1}|t|\alpha^{-1}|\sin(\theta_1)|^{-|\beta|} \qquad \forall\beta, \tag{6.45}$$

$$|D^\beta(b_m - \delta_{m0})| \le C_{m\beta}\mu^{-1}|t||\sin(\theta_1)|^{-m-|\beta|} \qquad \forall\beta, m. \tag{6.46}$$

Proof. Again a tedious but an easy proof we leave to the reader. $\square$

6.4 Tauberian estimates

Remark 6.4.1. (i) Let us set $x = y$. Then we get expression (6.38) for $U(y, y, t)$ albeit now due to definition $\theta_1 = \theta$.

(ii) Now we can continue in the same manner as before but to get a better estimate one needs to observe that the shift in x' (after rescaling) is now $\mu^{-1}\int \nabla_{x'}V \cdot F^{-1}\,dt$ and its difference with $\mu^{-1}\alpha t$ does not exceed

$C\mu^{-1}|t|\cdot|\operatorname{osc}(\nabla' V)|$ where $\operatorname{osc}(W)$ means an oscillation of W along trajectory (until time t).

However this trajectory must return to $x_3=y_3$ and therefore one can see easily that along it $p_3=O(\beta\mu^{-1}t)$ and $x_3-y_3=O(\beta\mu^{-2}t^2)$ and we need to assume that this is less than $\epsilon\alpha$ i.e.

$$|t|\le T_*=\epsilon\mu\min(1,\alpha^{\frac{1}{2}}\beta^{-\frac{1}{2}}),\quad \alpha:=|\nabla' V(y)|,\ \beta:=|\partial_{y_3}V(y)|. \tag{6.47}$$

Therefore only as $|t|\le T_*$ one can use nondegeneracy condition but it makes sense to consider $t:\ T_*\le|t|\le T^*=\epsilon\mu$. Then we have the following counterpart of proposition 5.2.1 where we need to sum (5.15) to $k=T_*$ and compare $\bar{k}=\mu^2 h\alpha^{-1}$ with T: namely, $\bar{k}\le T_*$ iff

$$\alpha\ge\max\bigl(\mu h,\beta^{\frac{1}{3}}(\mu h)^{\frac{2}{3}},\mu^{-1}\bigr). \tag{6.48}$$

Setting here $\beta=1$ (as without spatial averaging or micro-averaging $\partial_{\|\mathbf{F}}V/F$ is our foe and we need only estimate it from above) and thus $T_*=\epsilon\mu\alpha^{\frac{1}{2}}$ we arrive to estimate (6.50) below.

On the other hand, if we do not use non-degeneracy condition we need to sum

$$C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}|k|^{-\frac{1}{2}} \tag{6.49}$$

which results in $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}T^{\frac{1}{2}}$ and we arrive to estimate (6.51) below.

Proposition 6.4.2. *(cf. proposition 5.2.1). Let conditions* (2.1)–(2.3), (6.7)–(6.8), (0.5)–(0.6) *be fulfilled*[6]*. Then*

(i) Assume that conditions (6.47)–(6.48) *be fulfilled as well. Then after rescaling as* T *is given by* (6.47) *with* $\beta=1$

$$|F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)\Gamma_x\mathrm{U}|\le C\mu h^{-2}+ C\left\{\begin{aligned}&\mu^{\frac{5}{2}}h^{-1}\alpha^{-\frac{1}{2}}\bigl(1+(\log(\alpha/(\mu h)^{\frac{2}{3}}))_+\bigr) &&\textit{as}\ \ \alpha\ge(\mu h)^{\frac{2}{3}},\\ &\mu^{\frac{3}{2}}h^{-\frac{3}{2}}T^{\frac{1}{2}} &&\textit{as}\ \ \alpha\le(\mu h)^{\frac{2}{3}};\end{aligned}\right. \tag{6.50}$$

(ii) In the general case as $|t|\le T^*$

$$|F_{t\to\hbar^{-1}\tau}\bar{\chi}_T(t)\Gamma_x\mathrm{U}|\le C\mu h^{-2}+C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}T^{\frac{1}{2}}. \tag{6.51}$$

Then picking-up $T = T_*$ and dividing the left-hand expression of (6.50) by T_* or picking-up $T = T^*$ and dividing the left-hand expression of (6.51) by T^* we arrive to estimates (6.52), (6.53) below.

Corollary 6.4.3. *(i) In frames of proposition 6.4.2(i) with* $\alpha \ge (\mu h)^{\frac{3}{2}}$ *(which implies* $\mu \le h^{-1}$*)*

$$\begin{aligned}\text{(6.52)}\quad \mathsf{R}^{\mathsf{T}} := |e(0,0,\tau) - e^{\mathsf{T}}(0,0,\tau)| \le \\ Ch^{-2}\alpha^{-\frac{1}{2}} + C\mu^{\frac{3}{2}}h^{-1}\alpha^{-1}\bigl(1 + (\log(\alpha/(\mu h)^{\frac{2}{3}}))_+\bigr);\end{aligned}$$

(ii) In the general case

$$\mathsf{R}^{\mathsf{T}} \le Ch^{-2} + C\mu h^{-\frac{3}{2}};\tag{6.53}$$

(iii) In particular, according to (6.52)*, as* $\mu \le h^{-1}$*,* $\alpha \asymp 1$

$$\mathsf{R}^{\mathsf{T}} \le Ch^{-2} + C\mu^{\frac{3}{2}}h^{-1}\bigl(1 + |\log \mu h|\bigr)\tag{6.54}$$

which is $O(h^{-2})$ *as* $\mu \le (h|\log h|)^{-\frac{2}{3}}$*;*

(iv) In particular, according to (6.53)*,* $\mathsf{R}^{\mathsf{T}} = O(h^{-2})$ *as* $\mu \le h^{-\frac{1}{2}}$*;*

(v) Estimate (6.52) *is better iff* $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ *and*

$$\alpha \ge \mu^{-2}h^{-1} + (\mu h)^{\frac{1}{2}}|\log(\mu h)|.\tag{6.55}$$

6.5 Weyl estimates

The proof of the following statement is rather obvious:

Proposition 6.5.1. *(cf. proposition 5.3.1). Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.47)–(6.48), (0.5)–(0.6) *be fulfilled*[6]*. Then*

(i) As $\mu^2 h \le \alpha \le 1$ *estimate* (5.18) *holds and correction term* $h^{-3}\mathcal{N}_{x,\mathsf{corr}}$ *is delivered by* r*-term stationary phase approximation* $h^{-3}\mathcal{N}_{x,\mathsf{corr}(r)}$ *with an error not exceeding* (5.19)*.*

(ii) As $\alpha \le \mu^2 h$ *estimate* (5.19) *holds;*

(iii) In particular without any nondegeneracy condition $\mathsf{R}^{\mathsf{W}}_x = O(\mu^{\frac{3}{2}}h^{-\frac{3}{2}})$ *and it is* $O(h^{-2})$ *as* $\mu \le h^{-\frac{1}{3}}$*;*

(iv) On the other hand, $\mathsf{R}^{\mathsf{W}}_x = O(\mu^{-\frac{1}{2}}h^{-\frac{3}{2}} + \mu^{\frac{5}{2}}h^{-1})$ *as* $\alpha \asymp 1$ *and* $\mu \le h^{-\frac{1}{2}}$ *and therefore* $\mathsf{R}^{\mathsf{W}}_x = O(h^{-2})$ *as* $\alpha \asymp 1$ *and* $\mu \le h^{-\frac{2}{5}}$*.*

Combining with corollary 6.4.3 we conclude that

Theorem 6.5.2. *Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.48), (0.5)–(0.6) *be fulfilled*6)*. Then*

(i) In the general case

$$|e(x,x,0) - h^{-3}\mathcal{N}_x^{\mathsf{W}}(x,x,0)| \le Ch^{-2} + C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}; \tag{6.56}$$

(ii) Under non-degeneracy condition

$$|\nabla_{\perp \mathsf{F}} V/F| \asymp 1 \tag{6.57}$$

as $\mu \le h^{-\frac{1}{2}}$ *estimates*

$$|e(x,x,0) - h^{-3}\mathcal{N}_x^{\mathsf{W}}(x,x,0)| \le Ch^{-2} + C\mu^{\frac{5}{2}}h^{-1} \tag{6.58}$$

and

$$\begin{multline} |e(x,x,0) - h^{-3}\mathcal{N}_x^{\mathsf{W}}(x,x,0) - h^{-3}\mathcal{N}_{x,\mathsf{corr}(r)}| \le \\ Ch^{-2} + C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}(\mu^2 h)^{r+\frac{1}{2}} \end{multline} \tag{6.59}$$

hold.

6.6 Successive approximations

Let us try successive approximation method. As an approximation we successively try the generalized pilot-model operator A^0, the pilot-model operator $\bar{A}$ and the magnetic Weyl approximation. As for $\mu \le h^{-\frac{1}{3}}$ Weyl approximation delivers $O(h^{-2})$ error we assume that

$$\mu \ge h^{-\frac{1}{3}}. \tag{6.60}$$

6.6.1 Generalized pilot-model approximation, $\mu h \le 1$

We claim that

(6.61) Effectively if $x = y (= 0)$ we can estimate $\|A - A^0\|$ by

$$\zeta(k) := C\bigl(\mu^{-2}|k| + (\mu h)^{\frac{1}{2}}\mu^{-1}|k|^{-\frac{1}{2}}\bigr). \tag{6.62}$$

Really, due to our assumptions $g^{jl} = \delta_{jl} + O(|x'|^2) + O(|x'|\cdot|x_3|)$ with $|x'| = O(\mu^{-1})$, $x_3 = O(\mu^{-1}|k|)$ $(j,l = 1,2)$, $g^{33} = 1 + O(\mu^{-2})$ (before rescaling) the only term which does not allow such estimate is $g^{j3}D_jD_3$ with $j = 1,2$.

However for a classical trajectory to return back to time $t \asymp k$ we need to have $p_3 = O(\mu^{-1}k)$; in our analysis we will need also to satisfy uncertainty principle: $p_3^2|k| \ge \mu h$; so we estimate p_3 by $C\mu^{-1}|k| + C(\mu h)^{\frac{1}{2}}|k|^{-\frac{1}{2}}$ and as $g^{j3} = O(|x'|)$ this leads to (6.62).

Then the error in k-th winding does not exceed

$$\tag{6.63} C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}|k|^{-\frac{3}{2}} \times \bigl(\mu^{-2}|k| + (\mu h)^{\frac{1}{2}}\mu^{-1}|k|^{-\frac{1}{2}}\bigr) \times \mu^{-1}h^{-1}|k| \times \\ \begin{cases} \bigl(\dfrac{\mu^2 h}{\alpha|k|}\bigr)^{r+\frac{1}{2}} & \text{as}\ \ \alpha^{-1}\mu^2 h \le |k| \le (\mu^3 h)^{\frac{1}{2}},\\ 1 & \text{as}\ \ |k| \le \min\bigl(\alpha^{-1}\mu^2 h, (\mu^3 h)^{\frac{1}{2}}\bigr). \end{cases}$$

Really, the successive approximation makes sense only as $\zeta(k)|k| \le \mu h$ which in view of (6.60) is equivalent to $|k| \le (\mu^3 h)^{\frac{1}{2}}$.

Here as $|k| \ge \alpha^{-1}\mu^2 h$ we apply r-term stationary phase approximation as well. Also recall that this interval originally was $\alpha^{-1}\mu^2 h \le |k| \le \epsilon\mu\alpha^{\frac{1}{2}}$ [17] but it is non-empty iff $\alpha \ge (\mu h)^{\frac{2}{3}}$.

Then $(\mu^3 h)^{\frac{1}{2}} \le \mu\alpha^{\frac{1}{2}}$ and further $(\mu^3 h)^{\frac{1}{2}} \ge \alpha^{-1}\mu^2 h$ iff $\alpha \ge (\mu h)^{\frac{1}{2}}$. Only under this assumption we need to consider the first case in (6.63) but it is the most important case as it happens for $\alpha \asymp 1$.

Let us break (6.63) into two expressions: one with the second factor $\mu^{-2}|k|$ and another with the second factor $(\mu h)^{\frac{1}{2}}\mu^{-1}|k|^{-\frac{1}{2}}$.

In the general case (when we do not know if $\alpha \ge (\mu h)^{\frac{1}{2}}$) summation of the first expression returns $Ch^{-2}(\mu^3 h)^{\frac{1}{4}}$ and of the second expression returns a lesser $Ch^{-2}(1 + \log(\mu^3 h))$. Contribution of $k : |k| \ge (\mu^3 h)^{\frac{1}{2}}$ to the Tauberian expression is a sum of $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}|k|^{-\frac{3}{2}}$, which gives us $C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}(\mu^3 h)^{-\frac{1}{4}}$ i.e. exactly the same answer. Therefore we arrive to

(6.64) The Tauberian expressions for the original operator and its generalized pilot-model approximation differ by no more than $Ch^{-2}(\mu^3 h)^{\frac{1}{4}}$.

---

[17] Here the upper limit meant that $|x_3| \le \alpha$ along trajectory before k-th winging. Since we look at the trajectory returning at k-th winging, $|x_3| \le (\mu^{-1}|k|)^2$ before it and we need to satisfy $(\mu^{-1}|k|)^2 \le \alpha$.

Assume now that $\alpha \ge (\mu h)^{\frac{1}{2}}$. Again break (6.63) into two expressions the same way as before. Obviously summation with respect to k of the second expression returns $Ch^{-2}\bigl(1+(\log(\mu^2 h/\alpha))_+\bigr)$ no matter what r is.

As $r=0$ summation of the first expression with respect to k returns $C\mu h^{-\frac{3}{2}}\alpha^{-\frac{1}{2}}$. One can see easily that the contribution of $k: |k| \ge (\mu^3 h)^{\frac{1}{2}}$ to the Tauberian expression does not exceed $C\mu h^{-\frac{3}{2}}\alpha^{-\frac{1}{2}}$.

Therefore we arrive to

(6.65) As $\alpha \ge (\mu h)^{\frac{1}{2}}$ the Tauberian expressions for the original operator and its generalized pilot-model approximation differ by no more than $Ch^{-2} + C\mu h^{-\frac{3}{2}}\alpha^{-\frac{1}{2}}$.

As $r \ge 2$ summation of the first expression with respect to k returns its value $(\times |k|)$ as $k = \max(1, \mu^2 h/\alpha)$:

$$C\mu^{\frac{3}{2}} h^{-1}\alpha^{-\frac{3}{2}} \times \begin{cases} \bigl(\dfrac{\mu^2 h}{\alpha}\bigr)^{r-1} & \text{as } 1 \ge \mu^2 h/\alpha, \\ 1 & \text{as } 1 \le \mu^2 h/\alpha \end{cases} \tag{6.66}$$

and as $r=1$ we get $C\mu^{\frac{3}{2}} h^{-1}\alpha^{-\frac{3}{2}}(1+\log(\alpha(\mu h)^{-\frac{1}{2}})$. One can see easily that the contribution of $k: |k| \ge (\mu^3 h)^{\frac{1}{2}}$ to the Tauberian expression does not exceed these expressions as $r \ge 1$.

Therefore we arrive to

(6.67) As $\alpha \ge (\mu h)^{\frac{1}{2}}$ the Tauberian expressions with the subtracted $h^{-3}\mathcal{N}_{x,\mathsf{corr}(r)}$ i.e.

$$e^{\mathsf{T}}(x,x,\tau) - h^{-3}\mathcal{N}_{x,\mathsf{corr}(r)} \tag{6.68}$$

for the original operator and its generalized pilot-model approximation differ by no more than

$$Ch^{-2} + Ch^{-2}(\log(\mu^2 h/\alpha))_+ + \\ C\mu^{\frac{3}{2}} h^{-1}\alpha^{-\frac{3}{2}} \begin{cases} \bigl(1+\log(\alpha(\mu h)^{-\frac{1}{2}})\bigr) & \text{as } r=1, \alpha \ge \mu^2 h, \\ \bigl(1+\log(\mu^3 h)\bigr) & \text{as } r=1, \alpha \le \mu^2 h, \\ (\mu^2 h/\alpha)^{r-1} & \text{as } r \ge 2, \alpha \ge \mu^2 h, \\ 1 & \text{as } r \ge 2, \alpha \le \mu^2 h. \end{cases} \tag{6.69}$$

Combining with corollary 6.4.3 we estimate expressions (6.70)–(6.72) below for different α; in particular we arrive to

Theorem 6.6.1. *Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.48), (0.5)–(0.6) *be fulfilled*[6]*. Let* $e^0(.,.,.)$ *be a Schwartz kernel of the spectral projector for operator* A^0 *defined by* (6.36)–(6.37). *Then as* $h^{-\frac{1}{3}} \le \mu \le h^{-1}$

(i) In the general case

$$|e(x,x,0) - e^0(x,x,0)| \le C\mu^{\frac{3}{4}} h^{-\frac{7}{4}}; \tag{6.70}$$

(ii) Under non-degeneracy condition (6.57) *estimates*

$$|e(x,x,0) - e^0(x,x,0)| \le Ch^{-2} + C\mu h^{-\frac{3}{2}} \tag{6.71}$$

and

$$\begin{gathered} |e(x,x,0) - e^0(x,x,0) - h^{-3}\bigl(\mathcal{N}_{x,\mathsf{corr}(r)} - \mathcal{N}^0_{x,\mathsf{corr}(r)}\bigr)| \le \\ Ch^{-2}\bigl(1 + (\log\mu^2 h)_+\bigr) + C\mu^{\frac{3}{2}} h^{-1}\bigl(1 + |\log\mu h|\bigr) + \\ C\mu^{\frac{3}{2}} h^{-1} \begin{cases} \bigl(1 + \log(\mu^3 h)\bigr) & \text{as } r = 1, \\ (\mu^2 h)^{r-1} & \text{as } r \ge 2, \mu \le h^{\frac{1}{2}}, \\ 1 & \text{as } r \ge 2, \mu \ge h^{-\frac{1}{2}} \end{cases} \end{gathered} \tag{6.72}$$

hold.

Remark 6.6.2. Replacing $\alpha(x_3)x_j$ by $\alpha(0)x_j$ we estimate the norm of the perturbation by $\zeta_1(k) = \mu^{-1} \cdot (\mu^{-2}|k|^2 + \mu^{-1})$ which is less than expression (6.62) and thus leads to the lesser error. On the other hand, this new approximation is already a direct sum $A = A_{(2)} + B$ with the pilot model operator $A_{(2)}$ and

$$B = h^2 D_3^2 + V^0(x_3). \tag{6.73}$$

Therefore all above estimate remain valid with A^0 defined by (6.36)–(6.37) albeit with $\alpha_j = V_{x_j}(0,0,0)$.

6.6.2 Pilot-model approximation, $\mu h \le 1$

Consider now the pilot-model approximation $\bar{A}$ which differ from the approximation A^0 of remark 6.6.2 by $O(x_3^2)$. Then

(6.74) Effectively if $x = y$ we can estimate $\|A^0 - \bar{A}\|$ by

$$\zeta(k) := C\bigl(\mu^{-4}k^4 + \mu^{-1}h|k|\bigr). \tag{6.75}$$

Really, we need to estimate $O(x_3^2)$ and according to above arguments we estimate $|x_3|$ by $C\bigl(\mu^{-2}k^2 + (\mu h)^{\frac{1}{2}}|k|^{-\frac{1}{2}} \times \mu^{-1}|k|\bigr)$ which leads to (6.74).

Then we have estimate of the error in k-th winding not exceeding

$$C\mu^{\frac{3}{2}}h^{-\frac{3}{2}}|k|^{-\frac{3}{2}} \times \bigl(\mu^{-4}k^4 + \mu^{-1}h|k|\bigr) \times \mu^{-1}h^{-1}|k| \times \begin{cases} \bigl(\dfrac{\mu^2 h}{\alpha|k|}\bigr)^{r+\frac{1}{2}} & \text{as } |k| \ge \alpha^{-1}\mu^2 h,\\ 1 & \text{as } |k| \le \alpha^{-1}\mu^2 h. \end{cases} \tag{6.76}$$

The perturbation factor is $O(1)$ iff $|k| \lesssim \mu h^{\frac{1}{5}}$ and we need to compare it with $\alpha^{-1}\mu^2 h$: we have cases $\alpha \ge \mu h^{\frac{4}{5}}$ when $\alpha^{-1}\mu^2 h$ is less and $\alpha \le \mu h^{\frac{4}{5}}$ when $\mu h^{\frac{4}{5}}$ is less.

Case $\alpha \ge \mu h^{\frac{4}{5}}$. Then we need to sum

(a) Expression (6.76) with the last factor 1 from $|k| = 1$ to $|k| = \alpha^{-1}\mu^2 h$ which returns $C\mu^{\frac{1}{2}}h^{-\frac{5}{2}}|k|^{\frac{1}{2}} \times \bigl(\mu^{-4}k^4 + \mu^{-1}h|k|\bigr)$ calculated as $|k| = \alpha^{-1}\mu^2 h$:
$$C\mu^{\frac{11}{2}}h^2\alpha^{-\frac{9}{2}} + C\mu^{\frac{5}{2}}\alpha^{-\frac{3}{2}};$$
as $\alpha \asymp 1$ the first term does not exceed $C\mu^{\frac{3}{2}}h^{-1}$ iff $\mu \le h^{-\frac{3}{4}}$ while for the second term it is always true;

(b) Expression (6.76) with the last factor $\bigl(\mu^2 h/\alpha|k|\bigr)^{r+\frac{1}{2}}$ from $|k| = \alpha^{-1}\mu^2 h$ to $k = \mu h^{\frac{1}{5}}$ which returns what we got earlier plus value of $C\mu^{\frac{1}{2}}h^{-\frac{5}{2}}|k|^{\frac{1}{2}} \times \bigl(\mu^{-4}k^4 + \mu^{-1}h|k|\bigr) \times (\mu^2 h/\alpha|k|)^{r+\frac{1}{2}}$ calculated at $k = \mu h^{\frac{1}{5}}$ (may be with the logarithmic term for exceptional values of $r = 4, 1$) i.e.
$$C\mu h^{-\frac{8}{5}}(\mu h^{\frac{4}{5}}/\alpha)^{r+\frac{1}{2}} + C\mu h^{-\frac{6}{5}}(\mu h^{\frac{4}{5}}/\alpha)^{r+\frac{1}{2}};$$
as $r = 1$ we get $C\mu^{\frac{5}{2}}h^{-\frac{2}{5}}\alpha^{-\frac{3}{2}}$; as $\alpha \asymp 1$ this is less than Ch^{-2} as $\mu \le h^{-\frac{16}{25}}$; note that the second term always is $O(\mu^{\frac{3}{2}}h^{-1} + h^{-2})$;

(c) $C\mu^{\frac{3}{2}}h^{\frac{3}{2}}|k|^{-\frac{3}{2}}(\mu^2 h/\alpha)^{r+\frac{1}{2}}$ as $|k| \ge C\mu h^{\frac{4}{5}}$ which returns the expression above.

In total, we get modulo $O\bigl(h^{-2}+\mu^{\frac{3}{2}}h^{-1}\bigr)$

$$C\mu^{\frac{11}{2}}h^2\alpha^{-\frac{9}{2}} + C\mu^{\frac{5}{2}}\alpha^{-\frac{3}{2}} + C\mu h^{-\frac{8}{5}}(\mu h^{\frac{4}{5}}/\alpha)^{r+\frac{1}{2}}. \tag{6.77}$$

Case $\alpha \le \mu h^{\frac{4}{5}}$. Then we need to sum

(a) Expression (6.76) with the last factor 1 from $|k|=1$ to $|k|=\mu h^{\frac{1}{5}}$ which returns $C\mu^{\frac{1}{2}}h^{-\frac{5}{2}}|k|^{\frac{1}{2}} \times \bigl(\mu^{-4}k^4 + \mu^{-1}h|k|\bigr)$ calculated at $k=\mu h^{\frac{1}{5}}$ i.e.

$$C\mu h^{-\frac{8}{5}} + C\mu h^{-\frac{6}{5}};$$

here the first term is larger than $C\mu^{\frac{3}{2}}h^{\frac{3}{2}} + Ch^{-2}$ as $\mu \ge h^{-\frac{2}{5}}$ but the second term is always smaller than this;

(b) $C\mu^{\frac{3}{2}}h^{\frac{3}{2}}|k|^{-\frac{3}{2}}$ as $|k| \ge C\mu h^{\frac{4}{5}}$ which returns an above expression.

So, we get in this case

$$C\mu h^{-\frac{8}{5}}. \tag{6.78}$$

Thus we arrive to

Theorem 6.6.3. *Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.48), (0.5)–(0.6) *be fulfilled*6)*. Let $\bar{e}(.,.,.)$ be a Schwartz kernel of the spectral projector for the pilot-model operator $\bar{A}$. Then as $h^{-\frac{1}{3}} \le \mu \le h^{-1}$*

(i) In the general case

$$|e(x,x,0) - \bar{e}(x,x,0)| \le C\mu^{\frac{3}{4}}h^{-\frac{7}{4}} + C\mu h^{-\frac{8}{5}}; \tag{6.79}$$

(ii) Under non-degeneracy condition (6.57) *as $\mu \le h^{-\frac{4}{5}}$ estimates*

$$|e(x,x,0) - \bar{e}(x,x,0)| \le Ch^{-2} + C\mu h^{-\frac{3}{2}} + C\mu^{\frac{11}{2}}h^2 + C\mu^{\frac{5}{2}} + C\mu^{\frac{3}{2}}h^{-\frac{6}{5}} \tag{6.80}$$

and

$$
\begin{gathered}
|e(x,x,0)-\bar{e}(x,x,0)-h^{-3}\bigl(\mathcal{N}_{x,\mathsf{corr}(r)}-\mathcal{N}^0_{x,\mathsf{corr}(r)}\bigr)|\le \\
Ch^{-2}\bigl(1+(\log\mu^2h)_+\bigr)+C\mu^{\frac{3}{2}}h^{-1}\bigl(1+|\log\mu h|\bigr)+ \\
C\mu^{\frac{3}{2}}h^{-1}\begin{cases}\bigl(1+\log(\mu^3h)\bigr) & \textit{as}\quad r=1,\\ (\mu^2h)^{r-1} & \textit{as}\quad r\ge 2,\mu\le h^{\frac{1}{2}},\\ 1 & \textit{as}\quad r\ge 2,\mu\ge h^{-\frac{1}{2}}\end{cases}\qquad + \\
C\mu^{\frac{11}{2}}h^2+C\mu^{\frac{5}{2}}C\mu h^{-\frac{8}{5}}(\mu h^{\frac{4}{5}})^{r+\frac{1}{2}}.
\end{gathered}
\tag{6.81}
$$

hold.

Let us improve the above results. Note that we need only consider components of the estimate which are (in unrescaled x_3, p_3) due to $p_3 = O(\mu^{-1}k)$, $x_3 = O(\mu^{-2}k^2)$ rather than those which are due to the uncertainty principle $p_3 = O((\mu h)^{\frac{1}{2}}|k|^{-\frac{1}{2}})$, $x_3 = O(\mu^{-1}(\mu h)^{\frac{1}{2}}|k|^{\frac{1}{2}})$ as the latter brought a proper estimate in the above analysis.

Consider only case $\alpha \asymp 1$, $\beta \lesssim 1$. Then we need to consider only $\mu \ge h^{-\frac{16}{25}}$. Note that actually instead of $p_3 = O(t)$, $x_3 = O(t^2)$ we can use estimates $p_3 = O(\beta t)$, $x_3 = O(\beta t^2)$ and then our estimates acquire factor β^2 (recall that we are not discussing terms which are due to the uncertainty principle).

Case $h^{-16/25} \le \mu \le h^{-4/5}$. Consider (6.77) with $r = 1$; then the last term is the largest and the estimate with above improvement is $O(\beta^2\mu^{5/2}h^{-2/5})$ and it is $O(\mu^{3/2}h^{-1})$ as $\beta \le (\mu h^{\frac{3}{5}})^{-\frac{1}{2}}$. So we need to consider case

$$
\beta \ge (\mu h^{\frac{3}{5}})^{-\frac{1}{2}} \qquad (\ge h^{1/10}).
\tag{6.82}
$$

As $r = 0$ we get $\mu^{\frac{3}{2}}h^{-\frac{6}{5}}$ which is less than $\mu h^{-\frac{3}{2}}$ in (6.71);

Case $h^{-4/5} \le \mu \le h^{-1}$. Then estimate is (modulo $O(\mu^{-3/2}h^{-1})$) $O(\beta^2\mu h^{-8/5})$ and it is $O(\mu^{3/2}h^{-1})$ as $\beta \le (\mu h^{\frac{6}{5}})^{\frac{1}{4}}$. So we need to consider case

$$
\beta \ge (\mu h^{\frac{6}{5}})^{\frac{1}{4}} \qquad (\ge h^{1/10}).
\tag{6.83}
$$

Therefore we conclude that

Theorem 6.6.4. *As* $\beta := |\nabla_{\|\mathsf{F}} V/F| \le h^{-1/10}$ *estimates* (6.70)–(6.72) *of theorem 6.6.1 remain true for* A^0 *replaced by* $\bar{A}$.

So we need to consider the case when 1D-Schrödinger operator is almost non-degenerate and the corresponding dynamics is confined to zone $x_3 \le \beta$; rescaling $x_3 \mapsto x_{3,\mathsf{new}} = x_3/\beta$ we arrive to operator

$$\beta^2\Bigl(h_3^2 D_3^2 + V_3(x_3)\Bigr), \qquad h_3 = h/\beta^2 \ll 1, \quad V_3(x_3) = \beta^{-2}V(x_3) \tag{6.84}$$

satisfies $\partial_3 V_3(x_3) \asymp 1$. Then we can construct its solution by WKB method. Note that before any rescaling

$$U_{(1)}(x_3,y_3,t) = (2\pi h)^{-1}\int \exp\Bigl(ih^{-1}\bigl(S(x_3,t,\zeta) - y_3\zeta\bigr)\Bigr)B(x_3,t,\zeta)\,d\zeta \tag{6.85}$$

where S solves

$$S_t = S_{x_3}^2 + V(x), \qquad S|_{t=0} = x_3\zeta \tag{6.86}$$

and $B \sim \sum_k B_k h^k$ in the standard way.

Then one can prove easily that

$$S = \bar{S} + O\Bigl(x_3^2|t| + |x_3|(|\zeta| + |t|)|t| + \zeta^2|t|^3 + |t|^5\Bigr) \tag{6.87}$$

where

$$\bar{S} = x_3\zeta + t(\zeta^2 - \beta x_3) + \zeta\beta t^2 + \frac{2}{3}\beta^2 t^3 \tag{6.88}$$

solves the same problem for pilot-model $V = -\beta x_3$. Rescaling $x_3 \to x_3/\beta$, $h \mapsto h/\beta^2$ shows one can calculate S and $\bar{S}$ for $\beta = 0$, then plug $x_3 = x_3/\beta$, $\zeta = \zeta/\beta$ and multiply the result by β^2; (6.87) obviously survives but (6.88) improves to

$$S = \bar{S} + O\Bigl(x_3^2|t| + |x_3|(|\zeta| + \beta|t|)|t| + \zeta^2|t|^3 + |t|^5\Bigr). \tag{6.87*}$$

Obviously phase function

$$\bar{\phi}(x_3,y_3,t,\zeta) := \bar{S}(x_3,t,\zeta) - y_3\zeta \tag{6.89}$$

is equivalent to phase function

$$\bar{\phi} := (\zeta + \beta t)x_3 - (\zeta - \beta t)y_3 - 2\zeta^2 t - \frac{2}{3}\beta^2 t^3 \tag{6.90}$$

we used earlier.

Furthermore, obviously

(6.91) Phase function

$$\phi(x_3, y_3, t, \zeta) := S(x_3, t, \zeta) - y_3\zeta \tag{6.92}$$

is 1D-action and therefore the total exponent is also 3D-action associated with the generalized pilot-model.

Note that for a phase function (6.89) $\bar{\phi}_\zeta = 0$ iff $x_3 - y_3 + 2t\zeta + \beta t^2 = 0$ and then $\bar{\phi}_{\zeta\zeta} = -4t$ and therefore for $|t| \gg h$ we can apply a stationary phase method for both S and $\bar{S}$ and while the principal term would be of magnitude $(h|t|)^{-\frac{1}{2}}$ the error in l-term approximation will be of magnitude $(h|t|)^{-\frac{1}{2}}(h/|t|)^l$ or after all the substitution to U and rescalings it will be

$$C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}} \times (\mu h/|k|)^l \tag{6.93}$$

in the Tauberian expression. While almost useless for μh close to 1 and $k = 1$ the last factor is very important for larger k.

Then the contribution of zone $\{k : |k| \ge m\}$ to such error does not exceed (as $l = 1$)

$$C\mu^{\frac{5}{2}} h^{-\frac{1}{2}} m^{-\frac{3}{2}} \tag{6.94}$$

and it is $O(\mu^{-\frac{3}{2}} h^{-1})$ as $m \ge \mu^{\frac{2}{3}} h^{\frac{1}{3}}$. Obviously $m \ll \mu h^{\frac{1}{5}}$ and $m \ll \mu^2 h$ which means that in zone $k : |k| \le m$ we could perfectly deal with successive approximations.

On the other hand, replacing B by 1 and $\phi_{\zeta\zeta}$ by $-4t$ leads to an error

$$C\mu^{\frac{3}{2}} h^{-\frac{3}{2}} |k|^{-\frac{3}{2}} \times \mu^{-2} k^2 \times \min\bigl((\mu^2 h/|k|)^{\frac{1}{2}}, 1\bigr) \tag{6.95}$$

which after summation becomes $\mu^{\frac{1}{2}} h^{-1} \times \mu$ which is not as good as the Tauberian estimate. This spares us from more complicated formula. We leave to the reader

Problem 6.6.5. *(i) Write the approximation due to described combination of successive approximations and WKB method.*

(ii) Consider $\alpha \le 1$*.*

6.6.3 Magnetic Weyl approximation

In virtue of the above results the result of subsubsection 5.6.1 hold for general operators.

6.7 Strong and super-strong magnetic field

The standard method of successive approximations shows that replacing operator by the generalized pilot-model brings an error with the contribution of $\{t : |t| \asymp T\}$ (before rescaling) not exceeding

$$C\mu h^{-2} \times \bigl(\mu^{-1}h + C\mu^{-\frac{1}{2}}h^{\frac{1}{2}}(T^{\frac{1}{2}} + h^{\delta})\bigr) \tag{6.96}$$

and summation with respect to T running from h to 1 returns

$$C\mu h^{-2} \times \bigl(\mu^{-1}h + C\mu^{-\frac{1}{2}}h^{\frac{1}{2}}(1 + h^{\delta}|\log h|)\bigr) \asymp C\mu^{\frac{1}{2}}h^{-\frac{3}{2}} + C\mu^{\frac{1}{2}}h^{\delta-\frac{3}{2}}|\log \mu|$$

which is less than the Tauberian estimate. Therefore

Theorem 6.7.1. *Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.48), (0.5)–(0.6) *be fulfilled*6)*. Then as* $\mu h \gtrsim 1$

$$|e(x,x,0) - e^{0}(x,x,0)| \le C\mu h^{-\frac{3}{2}}. \tag{6.97}$$

Similarly, transition to the pilot model brings an error not exceeding

$$C\mu h^{-2} \times \bigl(\beta^{2}T^{4} + h^{\frac{4}{3}}\bigr) \tag{6.98}$$

and summation with respect from h to T_* returns $C\mu h^{-\frac{3}{2}} + C\mu h^{-2}\beta^{2}T_*^{4}$. On the other hand, Tauberian approach shows that the contribution of $|t| \ge T_*$ should not exceed $C\mu h^{-\frac{3}{2}}T_*^{-1}$ and minimizing it sum we arrive to $T_* = \beta^{-\frac{2}{5}}h^{\frac{1}{10}}$ and it is

$$C\mu h^{-\frac{3}{2}}\Bigl(1 + \beta^{\frac{2}{5}}h^{-\frac{1}{10}}\Bigr). \tag{6.99}$$

Therefore

Theorem 6.7.2. *Let conditions* (2.1)–(2.3), (6.7)–(6.8), (6.48), (0.5)–(0.6) *be fulfilled*6)*. Then as* $\mu h \gtrsim 1$

$$|e(x,x,0) - \bar{e}(x,x,0)| \le C\mu h^{-\frac{8}{5}}. \tag{6.100}$$

Further, if we pick up $T_* = \min(\beta^{-\frac{1}{2}}h^{\frac{1}{8}}, 1)$ we conclude that the contribution of zone $\{t : T_* \le |t| \le \epsilon\}$ to an error does not exceed $C\mu h^{-\frac{3}{2}}$. So we need to consider zone $\{t : |t| \ge h^{\frac{1}{8}}\}$. However in this zone we can apply stationary phase expression for $U_{(31)}$ which gives an error

$$C\int_T^1 \mu h^{-2}\times \bigl(\frac{h}{t}\bigr)^{\frac{3}{2}}\,dt \asymp C\mu h^{-2}\bigl(\frac{h}{T}\bigr)^{\frac{3}{2}}$$

which is $O(\mu h^{-\frac{3}{2}})$ as $T \ge h^{\frac{2}{3}}$.

Problem 6.7.3. *Construct this stationary phase approximation.*

Again for the magnetic Weyl approximation error we refer to subsubsection 5.2.

6.8 Micro-averaging

We are interested only in Tauberian, Weyl and magnetic Weyl estimates as we consider pilot-model and generalized pilot-model approximations a bit too complicated. We consider here only isotropic micro-averaging (i.e. with $\gamma_3 = \gamma$).

6.8.1 Tauberian estimates

Basically everything remains as in section 5.4 but we need to answer a question: "what is ℓ?" Or rather "what T^* we need to take?" Clearly $T^* = \epsilon\min(\alpha,\beta)$ fits but can we do better than this?

(i) Let $\beta \le \alpha$. Then as we take $\gamma_3 = \gamma$ we have $\nu(\boldsymbol{\gamma}) \asymp \alpha\gamma$ so averaging with respect to x_3 does not matter and therefore we do not need to assume that $\gamma_3 \le \epsilon\beta$ (which in this case would mean $\gamma \le \epsilon\beta$). In this case our assumption is $\gamma \le \epsilon\alpha$.

Then as $\nabla_{\perp\mathsf{F}} V/F$ has a less oscillation then $\epsilon\alpha$ with $\alpha = |\nabla_{\perp\mathsf{F}} V/F|\bigr|_{t=0}$ for time $T^* = \epsilon_1\alpha$ we can take $T^* = \epsilon_1\alpha$.

(ii) Let $\beta \ge \alpha$. Then as we take $\gamma_3 = \gamma$ we have $\nu(\boldsymbol{\gamma}) \asymp \beta\gamma$ so averaging with respect to x' does not matter as well as the shift and therefore we do not need to assume that $\gamma \le \epsilon\alpha$ (which in this case would mean $\gamma_3 \le \epsilon\beta$).

Then as $\nabla_{\|\mathsf{F}} V/F$ has a less oscillation then $\epsilon\beta$ with $\beta = |\nabla_{\|\mathsf{F}} V/F|\bigr|_{t=0}$ for time $T^* = \epsilon_1\beta$ we can take $T^* = \epsilon_1\beta$. However in this case we should not include factor $(\mu h/\alpha|k|)^{\frac{1}{2}}$ which is a loss as $\mu h/\alpha < h/\beta\gamma$ i.e. as $\mu\gamma \le \alpha\beta$.

So, in both cases we can take $\gamma_3 = \gamma$ and $T^* \asymp |\nabla V/F|$ (but we obviously need $\gamma \le |\nabla V/F|$).

6.8.2 Weyl and magnetic Weyl estimates

As we do not care about T^* anymore we just take $\gamma_3 = \gamma \le |\nabla V/F|$.

Problem 6.8.1. *As β is rather an obstacle in the pilot-model approximation, it would be interesting investigate this approximation with micro-averaging.*

7 Dirac energy: 3D-estimates

In this and the next sections we consider asymptotics of I defined by (0.2).

7.1 Tauberian formula

Let us consider first contribution of zone $\{(x,y):\ |x-y| \ge C\gamma\}$.

Proposition 7.1.1. *(cf. proposition 3.1.1). Let $\mu \le h^{-1}$. Then under conditions* (0.5)–(0.7) *the contribution of zone $\{(x,y):\ |x-y| \ge C\gamma\}$, to the remainder is $O(h^{-2}\gamma^{-\kappa})$ while the main part is given by the same expression* (0.1) *with $e(x,y,0)$ replaced by its standard implicit Tauberian approximation with $T \asymp \epsilon$* (0.11).

Proof. Proof repeats one of proposition 3.1.1 albeit according to Chapter 13 estimate (3.4) is replaced by

$$\|E(\tau,\tau')\bar{\psi}\|_1 \le Ch^{-3}\bigl(|\tau-\tau'| + CT^{-1}h\bigr) \qquad \forall \tau,\tau' \in [-\epsilon,\epsilon],\ \ T = \epsilon\mu \tag{7.1}$$

with $T = \epsilon$. □

Proposition 7.1.2. *(cf. proposition 3.1.3). Let conditions* (0.5)–(0.7) *be fulfilled. Then*

(i) As $0 < \kappa < 3$ *and* *either* $\kappa \ne 1, 2$ *or* $\kappa = 1, 2$ *and* $\omega(x, y)$ *is replaced by* $\omega(x, y) - \varkappa(\frac{1}{2}(x + y))|x - y|^{-\kappa}$ *with an appropriate smooth coefficient* $\varkappa(x)$*, with the error* $O(h^{-2-\kappa})$ *one can replace* $e(x, y, \tau)$ *by its standard Tauberian expression* (0.11) *in the formula* (0.11) *for* I*.*

(ii) As $\kappa = 1, 2$ *and* $\omega = \varkappa(\frac{1}{2}(x+y))|x-y|^{-\kappa}$*, with the error* $O(h^{-2-\kappa}|\log h|)$ *one can replace* $e(x, y, \tau)$ *by its standard Tauberian expression* (0.11) *in the formula* (0.11) *for* I*.*

Proposition 7.1.3. *(cf. proposition 3.1.6). Let conditions* (0.5) *and* (0.6) *be fulfilled.*

(i) Further, let *either* *condition* (0.8)[18] *be fulfilled* *or* $\mu \le h^{\delta-1}|\log h|^{-1}$ *with an arbitrarily small exponent. Then* (0.11) *and statements (i), (ii) of proposition 7.1.2 hold.*

(ii) Furthermore, let $h^{-1-\delta}\mu \le h^{-1}$*. Then* (0.11) *and statements (i), (ii) of proposition 7.1.2 hold with an extra factor* $(1 + \mu h^{1-\delta})$ *in the right-hand expressions.*

7.2 Superstrong magnetic field

Consider Schrödinger-Pauli operator as $\mu h \ge \epsilon_0$ which is a bit more tricky:

Proposition 7.2.1. *(cf. proposition 3.2.1). Let* $\mu h \ge \epsilon_0$*. Then*

(i) Contribution of zone $\{(x, y) :\ |x-y| \ge \gamma\}$ *to* I *does not exceed* $C\mu h^{-2}\gamma^{-\kappa}$*;*

(ii) Further,

$$|\mathsf{I}| \le C\mu h^{-2}\begin{cases} h^{-\kappa} & \text{as } 0 < \kappa < 1,\\ h^{-1}\bigl(1 + (\log \mu h)_+\bigr) & \text{as } \kappa = 1,\\ \mu^{\frac{1}{2}(\kappa-1)}h^{-\frac{1}{2}(\kappa+1)} & \text{as } 1 < \kappa < 3.\end{cases} \tag{7.2}$$

(iii) Furthermore, $\mathsf{I} = O(\mu h^{\infty})$ *under condition* (3.14)*; in particular it is the case as* $\mathfrak{z} < 1$ *and* $\mu h \ge C_0$*.*

---

[18] Which actually could be replaced by much weaker non-degeneracy condition of Chapter 13 for $d = 3$.

Proof. Statement (i) trivially follows from the fact that $\mathscr{L}^2$-norm of $e(.,.,\tau)$ does not exceed $C\mu^{\frac{1}{2}}h^{-1}$.

Meanwhile estimate of the contribution of the zone

$$\{(x,y): x\in B(0,1), y\in B(0,1), |x-y|\le\gamma\}$$

is more subtle: if $0<\kappa<1$ $\omega=\partial_{x_3}\omega'$ with $\omega'=O(\gamma^{1-\kappa})$ as $|x-y|\le\gamma$. Then

$$\begin{gathered}I_\gamma=\int |e(x,y,\tau)|^2\omega(x,y)\psi'_\gamma(x-y)\,dxdy=\\ -2h^{-1}\operatorname{Re} i\int hD_{x_3}e(x,y,\tau)\cdot e(y,x,\tau)\omega'(x,y)\psi'_\gamma(x-y)\,dxdy-\\ \int |e(x,y,\tau)|^2\omega(x,y,\tau)\cdot\partial_{x_3}\psi'_\gamma(x-y)\,dxdy\end{gathered}$$

with the first and second terms not exceeding $C\gamma^{1-\kappa}h^{-1}\times\mu h^{-2}$ and $C\gamma^{-\kappa}\times\mu h^{-2}$ respectively where we used the fact that $\mathscr{L}^2$-norm of $P_je(.,.,\tau)$ does not exceed $C\mu^{\frac{1}{2}}h^{-1}$. Setting $\gamma=h$ finishes the proof of (ii) in this case.

As $1\le\kappa<3$ we consider a partition of $\{(x,y):\ |x-y|\le h\}$ to subzones $\{(x,y):\ |x'-y'|\gtrsim\sigma\}$ and $\{(x,y):\ |x'-y'|\lesssim\sigma\}$ and in the former zone we repeat the same arguments as before albeit with $\omega'=O(\log|x'-y'|/h)$, $\omega'=O(|x'-y'|^{1-\kappa})$ for $\kappa=1$ and $1<\kappa<3$ respectively; then contribution of this zone does not exceed $Ch^{-1}|\log\sigma/h|\times\mu h^{-2}$ and $Ch^{-1}\sigma^{1-\kappa}\times\mu h^{-2}$ respectively; as $\sigma=\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ we arrive to estimate (7.2) for contribution of this zone as well.

Finally, contribution of the zone $\{(x,y):\ |x'-y'|\le\sigma, |x_3-y_3\le\gamma\}$ does not exceed $C\mu^2h^{-4}\times\int|z|^{-\kappa}\,dz$ with the integral taken over cylinder $\{|z'|\le\sigma, |z_3\le\gamma\}$ and not exceeding $C(1+|\log\sigma/\gamma|)\sigma^2$ and $C\sigma^{2-\kappa}$ respectively as $|e(x,y,\tau)=O(\mu h^{-2})$. Again for selected γ,σ we get estimate (7.2) for contribution of this zone as well.

Finally, statement (iii) is due to the fact that $e(x,y,0)=O(\mu h^\infty)$ as condition (3.14) is fulfilled in one of points x,y; see subsection 13.5.4. ☐

Proposition 7.2.2. *(cf. proposition 3.2.2). Let conditions* (0.1)–(0.5) *be fulfilled. Let $\mu h\ge\epsilon$ and one of the nondegeneracy conditions* (3.15*) or* (3.17*) be fulfilled.*

(i) As $\sigma \ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ *contribution of the zone*

$$\{(x,y):\ |x'-y'|\ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}, |x-y|\ge\gamma\}$$

to I *does not exceed* $C\mu h^{-1}\gamma^{-\kappa}$*;*

(ii) As $\gamma\ge C_0h$ *contribution of the zone*

$$\{(x,y):\ |x'-y'|\le C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}, |x_3-y_3|\ge\gamma\}$$

to I *does not exceed* $C\mu h^{-1}\gamma^{-1-\kappa}$*;*

(iii) Estimate

$$|\mathsf{I}-\mathsf{I}^{\mathsf{T}}|\le C\mu h^{-1}\left\{\begin{aligned} &h^{-\kappa} && \text{as}\ \ 0<\kappa<1,\\ &h^{-1}\bigl(1+(\log\mu h)_+\bigr) && \text{as}\ \ \kappa=1,\\ &\mu^{\frac{1}{2}(\kappa-1)}h^{-\frac{1}{2}(\kappa+1)} && \text{as}\ \ 1<\kappa<3,\ \kappa\ne2,\\ &\mu^{\frac{1}{2}}h^{-\frac{3}{2}}\bigl(1+(\log\mu h)_+\bigr) && \text{as}\ \ \kappa=2\end{aligned}\right.\tag{7.3}$$

holds.

Proof. To prove statement (i) recall that drift speed does not exceed $C\mu^{-1}$ and therefore Hilbert-Schmidt norm of $\psi E(\tau)\psi'$ does not exceed $C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}$ as ψ,ψ' are $\mathscr{L}_0^\infty$-functions with $\mathsf{dist}(\mathsf{supp}\,\psi,\mathsf{supp}\,\psi')\ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$.

Really, it is true for a Hilbert-Schmidt norm of $\bigl(E(\tau)-E(\tau')\bigr)\psi$ with $|\tau-\tau'|\le h$ and then by Tauberian theorem it is true for a Hilbert-Schmidt norm of $\bigl(E(\tau)-E^{\mathsf{T}}(\tau)\bigr)\psi'$ with E^{T} operator with the Schwartz kernel e^{T} with time $T\asymp1$:

$$\|\bigl(E(\tau)-E^{\mathsf{T}}(\tau)\bigr)\psi'\|_{\mathsf{HS}}\le C\mu^{\frac{1}{2}}h^{-\frac{1}{2}}.\tag{7.4}$$

However $\psi E^{\mathsf{T}}\psi'$ is negligible as $T\le\epsilon$ due to propagation results. This implies (i) obviously. Statement (ii) is proven in the same way but now $T\asymp\gamma$.

To prove statement (iii) we need to estimate contribution to $\mathsf{I}-\mathsf{I}^{\mathsf{T}}$ of the zones $\{(x,y):\ |x'-y'|\le h, h\le|x_3-y_3|\le\epsilon\}$ and $\{(x,y):\ |x-y|\le h\}$. However estimate of the contribution of the former is trivial due to estimate (7.4). So, consider zone $\{(x,y):\ |x-y|\le h\}$.

(a) As $0<\kappa<1$ we can replace as in the proof of proposition 7.2.1 $\omega(x,y)$ by $\omega'(x,y)=O(h^{1-\kappa})$ in this zone and one copy of $\bigl(e(x,y,\tau)-e^{\mathsf T}(x,y,\tau)\bigr)$ to $D_3\bigl(e(x,y,\tau)-e^{\mathsf T}(x,y,\tau)\bigr)$; however subsection 13.5.4 implies the following generalization of (7.4):

$$\|P_3^kP'^j\bigl(E(\tau)-E^{\mathsf T}(\tau)\bigr)\psi'\|_{\mathsf{HS}}\le C\mu^{\frac12}h^{-\frac12}(\mu h)^j \tag{7.5}$$

with $P'=(P_1,P_2)$ and therefore contribution of this zone to $(\mathsf I-\mathsf I^{\mathsf T})$ does not exceed $C\mu h^{-1-\kappa}$. In this case estimate (7.3) is proven.

(b) Let $1\le\kappa<3$. Then the above arguments imply

(7.6) As $1<\kappa<3$ contribution of the zone $\{(x,y):\ |x'-y'|\ge\sigma\}$ with $\sigma\ge C_0\mu^{-\frac12}h^{\frac12}$ to $\mathsf I$ does not exceed $C\mu h^{-2}\sigma^{1-\kappa}$; as $\kappa=1$ it does not exceed $C\mu h^{-2}(1+|\log\mu h|)$.

Note that with $\sigma=C_0\mu^{-\frac12}h^{\frac12}$ we get exactly the right-hand expression in (7.3). Therefore we need to estimate contribution of the zone $\{(x,y):|x'-y'\le\sigma,|x_3-y_3|\le h\}$ with $\sigma=C_0\mu^{-\frac12}h^{\frac12}$.

Consider case $1\le\kappa<2$ first. Note that $\omega=(3-\kappa)^{-1}\sum_j\partial_{z_j}z_j\omega$ [19] and therefore we can replace ω either by $(x_j-y_j)\omega=O(|x'-y'|\cdot|x-y|^{-\kappa})$ with $j=1,2$ or by $(x_3-y_3)\omega=O(|x-y|^{1-\kappa})$, simultaneously applying $h^{-1}P_j$ to one of the factors $\bigl(e(x,y,\tau)-e^{\mathsf T}(x,y,\tau)\bigr)$. Then we can replace $(x_j-y_j)\omega$ by $(x_j-y_j)\omega'$ simultaneously applying $h^{-1}P_j$ to one of the factors $\bigl(e(x,y,\tau)-e^{\mathsf T}(x,y,\tau)\bigr)$ (the same as before or another one).

So as $j=1,2$ we estimate the term by $C\mu h^{-1}\times h^{-2}(\mu h)^{\frac12}\times\sigma\times\sigma^{1-\kappa}$ with factors $(\mu h)^{\frac12}$ and σ coming from P_j and (x_j-y_j); as $\kappa=1$ the last factor is replaced by $|\log\sigma/\gamma|$. As $j=3$ these factors are replaced by 1 and h respectively with the same product. Note that when we drag the last ∂_{x_3} through x_3 they may cancel one another but the power becomes $(1-\kappa)$ which we treated already (and while the gain in this power is exactly compensated by an extra factor h^{-1}.)

Plugging σ we get a proper estimate (7.3).

(c) The same arguments work for $2\le\kappa<3$ as well albeit now $\omega=(3-\kappa)^{-2}\sum_{j,k}\partial_{z_j}\partial_{z_k}z_jz_k\omega$. Note that when we drag the last ∂_{x_3} through x_3 they may cancel one another but the power becomes $(1-\kappa)$ which we treated already (and while the gain in this power is exactly compensated by an extra factor h^{-1}.) However as $\kappa = 2$ we get power -1 and then logarithm comes in. $\square$

---

[19] Recall that we replace (x,y) by $(x,z)=(\frac12(x+y),x-y)$.

8 Dirac energy: 3D-calculations

8.1 Pilot Model

8.1.1 Transformations

Consider first the pilot-model operator (5.1).

Let us rescale as before[20]. Then $U(x, y, t)$ in comparison with $U_{(2)}(x', y', t)$ has factor $U_{(1)}(x_3, y_3, t)$ defined by (5.4). Let us consider effect of all these changes ignoring other variables; again without any loss of the generality we assume that $\omega(x, y)$ is given by (4.3);

We can replace variables x, y with new variables $\mathsf{x} := \frac{1}{2}(x + y)$ and $z := (x - y)$ and rescale. Note that the phase of $U_{(1)}$ is linear with respect to x_3 and we can get rid off $\mu^{-1}d\mathsf{x}_3$ integration in the same manner we got rid off $\mu^{-1}\,d\mathsf{x}_1$: we replace $\Omega(\dots, \mathsf{x}_3, .)$ by its partial Fourier transform $\mathsf{x}_3 \to -\mu^{-1}h^{-1}(t' - t'')$ with an extra factor 2π; thus in comparison with 2D case we have

- $\hat{\Omega}\bigl(2\mu^{-1}h^{-1}\alpha(t' - t''), -\mu^{-1}h^{-1}\beta(t' - t''), z\bigr)$
  where $\hat{\Omega}(., ., z)$ is a partial Fourier transform with respect to $(\mathsf{x}_1, \mathsf{x}_3)$ (instead of $\hat{\Omega}\bigl(2\mu^{-1}h^{-1}\alpha(t' - t''), z\bigr)$);

- an extra factor which after easy reductions becomes

$$\frac{1}{4}h^{-1}(t't'')^{-\frac{1}{2}}\exp\Bigl(i\hbar^{-1}(t'-t'')\bigl(-\frac{1}{8}(t't'')^{-1}z_3^2-\frac{2}{3}\beta^2\mu^{-2}(t'^2+t't''+t''^2)\bigr)\Bigr),$$

- integration with respect to dz_3 which enters both this factor and $\hat{\Omega}(., ., z)$.

---

[20] We need to add factor $\mu^{-6+\kappa} = \mu^{-3} \times \mu^{-3} \times \mu^{-\kappa}$ where two factors μ^{-3} are coming from dx and dy and μ^{κ} from ω.

Taking in account all these modifications we need to replace (4.4) by

$$\begin{gathered}
\mathsf{I} = \\
\frac{1}{8}(2\pi)^{-3}\hbar^{-3}\mu^{\kappa+3}\iint\iiint \hat{\Omega}(4\hbar^{-1}\alpha s, -\hbar^{-1}\beta s, z)\times \\
\csc(t+s)\csc(t-s)(t+s)^{-\frac{3}{2}}(t-s)^{-\frac{3}{2}}\times \\
\exp\Bigl(i\hbar^{-1}\Bigl[-\frac{1}{4}\bigl(\cot(t+s)-\cot(t-s)\bigr)\bigl(z_1^2+(z_2+2t\mu^{-1}\alpha)^2+4s^2\mu^{-2}\alpha^2\bigr) \\
-\bigl(\cot(t+s)+\cot(t-s)\bigr)s\mu^{-1}\alpha - 2s(\tau-\mu^{-2}\alpha^2)+ \\
2s\bigl(-\frac{1}{8}(t+s)^{-1}(t-s)^{-1}z_3^2-\frac{2}{3}\beta^2\mu^{-2}(3t^2+s^2)\bigr)\Bigr]\Bigr)\,dtds\,\psi(\mu^{-1}z)dz_1dz_2dz_3.
\end{gathered}\tag{8.1}$$

8.1.2 Case $\lambda^2 \gg \mu h$

Assume first that

$$\lambda^2 := \alpha^2+\beta^2 \ge h^{1-\delta}.\tag{8.2}$$

Remark 8.1.1. (cf. remark 4.1.2). (i) We take λ-admissible with respect to x function $\Omega_\lambda = \Omega(\mathsf{x}/\lambda, .)$.

Then the virtue of the factor $\hat{\Omega}(4\hbar^{-1}\alpha\gamma s, -\hbar^{-1}\beta\gamma s, \cdot)$ under assumption (8.2) we need to consider only $|s| \le h^\delta$ and therefore we can consider separately $|t'| \le \epsilon_0, |t''| \le \epsilon_0$ and $|t'| \ge \epsilon_0, |t''| \ge \epsilon_0$;

(ii) Note that due to section 6.3 contribution of zone $\{|t'| \le \epsilon_0, |t''| \le \epsilon_0\}$ defined by integral expressions (8.1) etc with an extra factor $\bar{\chi}_{\epsilon_0}(t')$ or $\bar{\chi}_{\epsilon_0}(t'')$ or $\bar{\chi}_{\epsilon_0}(t)$ differs from the same expression for non-magnetic Schrödinger operator by $O(\mu h^{-2-\kappa}\times\hbar^\kappa) = O(\mu^{\kappa+1}h^{-2})$ as $\kappa \ne 1$ and by $O(\mu^2 h^{-2}|\log\mu|)$ as $\kappa = 1$.

(iii) Furthermore, if we remove from this expression for a non-magnetic Schrödinger operator cut-off $\{|t'| \ge \epsilon_0\}$ then the error would not exceed the same expression as well.

Let us consider contribution of zone $\{|t'| \ge \epsilon_0, |t''| \ge \epsilon_0\}$ defined by an integral expressions (8.1) etc with an extra factor $\bigl(1-\bar{\chi}_{\epsilon_0}(t')\bigr)$. Due to remark 8.1.1(i) we need to consider only t', t'' belonging to the same tick.

Let us consider first zone

$$\{|s|\geq \hbar/\lambda^2, |\sin(t)|\geq C|s|\}. \tag{8.3}$$

Then integration by parts with respect to z' delivers one of the factors (4.7), (4.8). Thus integrating by parts many times in the zone where both of these factors are less than 1 we acquire factors (4.9), (4.10) respectively.

On the other hand, integration by parts with respect to z_3 delivers factor $\hbar|s|^{-1}t^2|z_3|^2$; therefore integrating by parts many times in the zone $\{|z_3|\geq \hbar^{\frac{1}{2}}|s|^{-\frac{1}{2}}|t|\}$ delivers factor

$$\bigl(1+|s|\hbar^{-1}t^{-2}|z_3|^2\bigr)^{-l}. \tag{8.4}$$

Case $1<\kappa<3$. Multiplying by $|z|^{-\kappa}$ and integrating we get after multiplication by $|\sin(t)|^{-2}$ expression (4.11) albeit with κ replaced by $(\kappa-1)$ i.e.

$$\asymp |\sin(t)|^{1-\kappa}|s|^{-\frac{3}{2}+\frac{1}{2}\kappa}\hbar^{\frac{3}{2}-\frac{1}{2}\kappa}. \tag{8.5}$$

Integrating by t over k-th tick intersected with $\{t:\ |\sin(t)|\geq |s|\}$ we get (4.12) modified the same way i.e.

$$C\hbar^{\frac{3}{2}-\frac{1}{2}\kappa}\left\{\begin{aligned} &|s|^{-\frac{3}{2}+\frac{1}{2}\kappa} && 1<\kappa<2,\\ &|s|^{-\frac{1}{2}}(1+|\log|s||) && \kappa=2,\\ &|s|^{\frac{1}{2}-\frac{1}{2}\kappa} && 2<\kappa<3.\end{aligned}\right. \tag{8.6}$$

This expression (8.6) must be either integrated with respect to s over $\{|s|\leq \hbar/\lambda^2\}$ or multiplied by $(\hbar/\lambda^2)^l|s|^{-l}$ due to factor $\hat{\Omega}$ and integrated over $\{|s|\geq \hbar/\lambda^2\}$, resulting in both cases in the same answer which is the value of (8.6) $\times\ |s|$ calculated as $s=\hbar/\lambda^2$ which is similarly modified expression (4.13) i.e.

$$C\hbar\left\{\begin{aligned} &\lambda^{1-\kappa} && 1<\kappa<2,\\ &(1+|\log(\hbar/\lambda^2)|)\lambda^{-1} && \kappa=2,\\ &\hbar^{2-\kappa}\lambda^{-3+\kappa} && 2<\kappa<3.\end{aligned}\right. \tag{8.7}$$

In addition to zone (8.3) we need to consider zone (4.14) defined as $\{|\sin(t')|\asymp |s|,\ |\sin(t'')|\leq |s|\}$; its tween $\{|\sin(t')|\asymp |s|,\ |\sin(t'')|\leq |s|\}$ is considered in the same way.

In zone (4.14) $|\cot(t')-\cot(t'')|\asymp|\sin(t'')|^{-1}$ and in this case factors (4.7), (4.8) are replaced by (4.7)′, (4.8)′ and (4.9), (4.10) by 4.9, (4.10)′ while factor (8.4) is preserved.

Then, multiplying by $|z|^{-\kappa}$ and integrating we get after multiplication by $|\sin(t'')|^{-1}|\sin(t')|^{-1}$ similarly modified expression (4.11)′

$$\hbar^{\frac{3}{2}-\frac{1}{2}\kappa}|\sin(t'')|^{\frac{1}{2}-\frac{1}{2}\kappa}|s|^{-1}; \tag{8.5$'$}$$

then integrating by $|t''|$ over one tick but with $|\sin(t'')|\le|s|$ we get modified expression (4.12)′ i.e.

$$\hbar^{\frac{3}{2}-\frac{1}{2}\kappa}|s|^{\frac{1}{2}-\frac{1}{2}\kappa}. \tag{8.6$'$}$$

Finally, either integrating over $\{|s|\le h/\varepsilon\lambda\}$ or multiplying by $|s|^{-l}(h/\varepsilon\lambda)^l$ and integrating over $\{|s|\ge h/\varepsilon\lambda\}$ we get in both cases the same answer (8.6)′ × $|s|$, calculated as $s=\hbar/\lambda^2$ not exceeding (8.7).

Therefore the total contribution of zones (8.3) and (4.14) is given by expression (8.7). Then multiplying by $|k|^{-3}\mu^{\kappa}h^{-3}\lambda^3$ we get after summation with respect to $k:|k|\ge 1$ the value as $k=1$ i.e. (8.7) × $\mu^{\kappa}h^{-3}\lambda^3$.

Therefore we arrive to analogue of proposition 4.1.3

$$|\mathsf{I}_l ambda^{\mathsf{T}}-\mathsf{I}^{\mathsf{T}}_\lambda{}'|\le R^{\mathsf{W}}(\lambda):= \\ C\mu^{\kappa+1}h^{-2}\lambda^3\begin{cases}\lambda^{1-\kappa} & 1<\kappa<2,\\ (1+|\log(\hbar/\lambda^2)|)\lambda^{-1} & \kappa=2,\\ \mu^{2-\kappa}h^{2-\kappa}\lambda^{-3+\kappa} & 2<\kappa<3\end{cases} \tag{8.8}$$

where $\mathsf{I}^{\mathsf{T}}_\alpha{}'$ is a Tauberian expression for I albeit with $T=\epsilon_0\mu^{-1}$.

Case $0<\kappa\le 1$. In this case integration over $\{z:|z|\le C_0\}$ [21] and multiplication by $|\sin(t)|^{-2}$ results in

$$C|\sin(t)|^{-2}\rho^2\begin{cases}\zeta^{1-\kappa} & 0<\kappa<1,\\ (1+(\log\zeta/\rho))_+) & \kappa=1\end{cases} \tag{8.9}$$

with

$$\zeta=\zeta(s,k)=\min\bigl((\hbar/|s|)^{\frac{1}{2}}|k|,1\bigr), \tag{8.10}$$

$$\rho=\min\bigl((\hbar/|s|)^{\frac{1}{2}},1\bigr) \tag{8.11}$$

---

[21] Corresponding to $|x-y|\le C_0\mu^{-1}$.

(cf. expression (8.5)).

One can see easily that

(8.12) Integral by t over one tick intersected with $\{t : |\sin(t)| \ge |s|\}$ does not exceed $C\hbar|s|^{-1}\zeta^{1-\kappa}$ as $|s| \ge \hbar$ and $C(\hbar|s|^{-1})^{\frac{1}{2}}\zeta^{1-\kappa}$ as $|s| \le \hbar$. As $\kappa = 1$ one needs to include an extra factor $|\log \zeta/\rho|$, ρ calculated as $|\sin(t)| = 1$ for $|s| \ge \hbar$ and $\sin(t)| = (|s|/\hbar)^{\frac{1}{2}}$ for $|s| \le \hbar$.

Meanwhile in the zone (4.14) integration over $\{z : |z| \le C_0\}$ and multiplication by $s^{-1}|\sin(t'')|^{-1}$ results in expression (8.11) and integration by $\{t'' : |\sin(t'')| \le |s|\}$ results in much lesser term $C\hbar\zeta^{1-\kappa}$, with some logarithmic factor as $\kappa = 1$.

Now integral by $\{s : |s| \lesssim 1\}$ does not exceed $C\hbar|k|^{1-\kappa}$ as $0 < \kappa < 1$ and $C\hbar(1 + |\log \hbar| + |\log |k|)$ as $\kappa = 1$. Finally, multiplication by $\mu^{\kappa+3}\hbar^{-3}|k|^{-2}$ and summation with respect to $k : |k| \ge 1$ results in $C\mu^{\kappa+1}h^{-2}$ as $0 < \kappa < 1$ and $C\mu^2 h^{-2}|\log \mu h|$ as $\kappa = 1$.

Therefore we arrive to analogue of proposition 4.1.3:

$$|\mathsf{I}_lambda^{\mathsf{T}} - \mathsf{I}^{\mathsf{T}\prime}_{\lambda}| \le R^{\mathsf{W}}(\lambda) := \\ C\mu^{\kappa+1}h^{-2}\lambda^3 \begin{cases} 1 & 0 < \kappa < 1, \\ (1 + |\log(\mu h)|) & \kappa = 1. \end{cases} \tag{8.13}$$

Combining with (8.8) we arrive to

Proposition 8.1.2. *(cf. corollary 4.1.3). (i) For the pilot-model operator with* $1 \le \mu \le h^{-1}$ *under assumption* (8.2) *estimates* (8.8), (8.13) *hold for* $1 < \kappa < 3$, $0 < \kappa \le 1$ *respectively.*

(ii) In particular, as $\lambda = 1$

$$|\mathsf{I}^{\mathsf{T}} - \mathsf{I}^{\mathsf{T}\prime}| \le R^{\mathsf{W}} := \\ C\mu^{\kappa+1}h^{-2} \begin{cases} 1 & 0 < \kappa < 2, \kappa \ne 1, \\ (1 + |\log(\mu h)|) & \kappa = 1, 2, \\ \mu^{2-\kappa}h^{2-\kappa} & 2 < \kappa < 3. \end{cases} \tag{8.14}$$

Following the same proof as in proposition 4.1.4 one can prove easily that

$$|\mathsf{I}^{\mathsf{T}\prime} - \mathcal{I}^{\mathsf{W}}| \le C\mu^2 h^{-1-\kappa} \tag{8.15}$$

which does not exceed $R^{\mathsf{W}}(\lambda)$ implying

Corollary 8.1.3. *(cf. corollary 4.1.5). In frames of proposition 8.1.2*

$$|\mathsf{I}^{\mathsf{T}}_{\lambda} - \mathcal{I}^{\mathsf{W}}_{\lambda}| \le CR^{\mathsf{W}}(\lambda). \tag{8.16}$$

Remark 8.1.4. After summation with respect to partition we restore estimate (8.14) under some non-degeneracy assumption. Comparing with the Tauberian remainder estimate $O(h^{-2-\kappa})$ we conclude that R^{W} is lesser as $\mu \le h^{-\kappa/(\kappa+1)}$ $(0<\kappa<2)$ and $\mu \le h^{-\frac{2}{3}}$ $(2\le \kappa <3)$ (with some correction for $\kappa = 1,2$).

In particular, $\mu \le h^{-\frac{1}{2}}$ would suffice for $\kappa > 1$ and therefore improvement as in subsubsection 4.1.3. "Improvement" is not needed.

8.1.3 Case $\lambda^2 \not\gg \mu h$

Assume now that $\mu^{-1} \le \lambda$ and $\lambda^2 \le \mu h^{1+\delta}$. Then in contrast to the previous we will need to compute contributions of pair of ticks (k', k'') with $k' \ne k''$. Then if t', t'' belong to k'-th and k''-th ticks respectively we denote $r = k' - k''$ and $s = t' - t'' - 2\pi r$.

However we still need to distinguish cases $1 < \kappa < 3$ and $0 < \kappa \le 1$.

Case $1<\kappa<3$. In this case we according to the previous subsection need just to take results for $d=2$, plug $\kappa := (\kappa - 1)$, $\alpha = \lambda$ and multiply by $h^{-1}\lambda$ resulting in

Proposition 8.1.5. *(cf. proposition 4.1.10). For the pilot-model operator with* $1 \le \mu \le h^{-1}$, $1<\kappa<3$

$$|\mathsf{I}^{\mathsf{T}}_{(\lambda)} - \mathcal{I}^{\mathsf{W}}_{(\lambda)}| \le (8.8) + (8.18) \qquad \textit{as}\ \ \lambda^2 \ge \mu h, \tag{8.17}$$

with

$$C\mu^{\frac{1}{2}\kappa - \frac{1}{2}} h^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^3 (\mu h/\lambda^2)^l \tag{8.18}$$

and

$$|\mathsf{I}^{\mathsf{T}}_{(\lambda)} - \mathcal{I}^{\mathsf{W}}_{(\lambda)}| \le (8.20) + (8.21) + (8.22) \qquad \textit{as}\ \ \lambda^2 \le \mu h \tag{8.19}$$

with

$$C\mu^{\frac{1}{2}\kappa+\frac{3}{2}}h^{-\frac{3}{2}-\frac{1}{2}\kappa}\lambda^3,\tag{8.20}$$

$$C\mu^{\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{3}{2}-\frac{1}{2}\kappa}\lambda^3\bigl(1+(\log(\mu h/\lambda^2))_+\bigr)^2,\tag{8.21}$$

$$C\mu^{\frac{1}{2}\kappa-\frac{1}{2}}h^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^3\bigl(1+(\log(\mu h/\lambda^2))_+\bigr)\tag{8.22}$$

Proof. Easy details are left to the reader. □

Case $0<\kappa\le 1$. In this case we need to modify arguments of the previous subsubsection.

(i) Consider case $k'-k''=r=0$ first. As we did not use $\lambda^2\gg\mu h$ estimate (8.13) for contributions of such terms remains valid.

(ii) Consider now case $k'-k''=r\ne 0$. Note that then according to (8.4) one should use $\zeta=|k'+k''|\cdot(\hbar|k-k'|)^{\frac{1}{2}}$

As $\lambda^2\ge\mu h$ the last expression before summation with respect to k $(=k')$ acquires factor $(|k'-k''|\lambda^2/\mu h)^{-l}$ and we get after summation with respect to $k'\ne 0$, $k''\ne 0$ the same right-hand expression of (8.13).

Meanwhile, as $\lambda^2\le\mu h$ we get before summation (as $0<\kappa<1$)

$$C\mu^{\kappa+1}h^{-2}\lambda^3|k'|^{-1}|k''|^{-1}\bigl(|k'+k''|/|k'-k''|^{\frac{1}{2}})\bigr)^{1-\kappa}\times\\ \bigl(1+|k'-k''|\lambda^2/\mu h\bigr)^{-l}\tag{8.23}$$

(with a logarithmic factor as $\kappa=1$).

It is enough to sum with as $1\le k''$, $k'=k''+r$, $r\ge 1$; then the sum is

$$\asymp C\mu^{\kappa+1}h^{-2}\lambda^3\sum_{r\ge 1}r^{-\frac{1}{2}-\frac{1}{2}\kappa}\bigl(1+r\lambda^2/\mu h\bigr)^{-l}\asymp C\mu^{\kappa+1}h^{-2}\lambda^3(\mu h/\lambda^2)^{\frac{1}{2}-\frac{1}{2}\kappa};$$

so we get

$$C\left\{\begin{aligned}&\mu^{\frac{3}{2}+\frac{1}{2}\kappa}h^{-\frac{3}{2}-\frac{1}{2}\kappa}\lambda^{2+\kappa} && 0<\kappa<1,\\ &\mu^2h^{-2}\lambda^3\bigl(1+\log(\mu h)\bigr)^2 && \kappa=1\end{aligned}\right.\tag{8.24}$$

where case $\kappa=1$ we left to the reader.

(iii) Consider now $k'=0, k''\ne 0$; its tween case $k'\ne 0, k''=0$ is addressed in the same way. To do this we need to modify our expression fo I: namely, we do not divide by t' and then take Fourier transform $F_{t'\to\hbar^{-1}\tau}$; instead we take Fourier transform $F_{t'\to\hbar^{-1}\tau'}$, then integrate by τ' to τ and divide by $\hbar$; modifying this way (8.1) we arrive instead of (8.1) to the expression similar to (4.30).

Then repeating the similar arguments of section 4 with the above modification, we arrive to

$$C\mu^\kappa h^{-3}\lambda^3|k''|^{-\frac{1}{2}-\frac{1}{2}\kappa}\bigl(1+|k''|\lambda^2/\mu h\bigr)^{-l}$$

instead of (8.23) and summation with respect to $k''\,:\,|k''|\ge 1$ returns $C\mu^\kappa h^{-3}\lambda^3(\mu h/\lambda^2)^l$ as $\lambda^2\ge\mu h$ and $C\mu^\kappa h^{-3}\lambda^3(\mu h/\lambda^2)^{\frac{1}{2}-\frac{1}{2}\kappa}$ as $\lambda^2\le\mu h$.

So we arrive to

$$C(\mu h/\lambda^2)^l\mu^\kappa h^{-3}\lambda^3\begin{cases}1 & 0<\kappa<1,\\ \bigl(1+\log(\mu h)\bigr)^2 & \kappa=1\end{cases}\tag{8.25}$$

as $\lambda^2\ge\mu h$ and

$$C\mu^{\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^{2+\kappa}\begin{cases}1 & 0<\kappa<1,\\ \bigl(1+\log(\mu h)\bigr)^2 & \kappa=1\end{cases}\tag{8.26}$$

as $\lambda^2\le\mu h$.

Thus we proved

Proposition 8.1.6. *For the pilot-model operator with* $1\le\mu\le h^{-1}$, $0<\kappa\le 1$

$$|\mathsf{I}^{\mathsf{T}}_{(\lambda)}-\mathcal{I}^{\mathsf{W}}_{(\lambda)}|\le (8.13)+(8.25)\qquad\textit{as}\quad\lambda^2\ge\mu h,\tag{8.27}$$

and

$$|\mathsf{I}^{\mathsf{T}}_{(\lambda)}-\mathcal{I}^{\mathsf{W}}_{(\lambda)}|\le (8.13)+(8.24)+(8.26).\qquad\textit{as}\quad\lambda^2\le\mu h\tag{8.28}$$

8.2 General operators

Consider now general operators satisfying conditions (0.5)–(0.6) and either (0.7) or (0.8) as $\tau=0$.

First, using proposition 7.1.1 and propagation results we conclude that

Proposition 8.2.1. *(cf. proposition 4.2.1). Let conditions* (0.5)–(0.6) *be fulfilled. Then*

(i) Contribution of the zone $\{(x,y):\ |x-y|\ge \epsilon_0\mu^{-1}\}$ *(before rescaling) to* I^{T} *does not exceed* $C\mu^{\kappa+1}h^{-2}$ *and*

(ii) Under condition (0.7) *contribution of time* $\{|t|\ge\epsilon_0\}$ *(before rescaling) to* I^{T} *does not exceed* $C\mu^{\kappa+1}h^{-2}$.

Note that $O(\mu^{\kappa+1}h^{-1})$ either coincides or is smaller than the remainder estimate (8.14) and as long as we are interested in Weyl approximation $\mathcal{I}^{\mathsf{W}}$ we should be completely happy with it.

Under condition (0.7) we are done arriving to statement (i) of theorem 8.2.3 below.

Under condition (0.8) let us introduce the scaling function

$$lambda(x)=\epsilon_0|\nabla VF^{-1}|+\frac{1}{2}\bar\lambda,\qquad \bar\lambda=C_0\mu^{-1}\tag{8.29}$$

and consider covering of $B(0,1)$ by λ-admissible elements. In virtue of proposition 8.2.1(i) we need to consider only pairs (x,y) belonging to the same element. Due to the rescaling arguments, the contribution of one such element to $|I^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed $C\mu h^{-1-\kappa}\lambda^3$ and therefore

(8.30) Under condition (0.8) contribution of the zone

$$\{(x,y):|x-y|\le C_0\mu^{-1},\ |\nabla VF^{-1}|\le\lambda\}$$

to $|I^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed $C\mu h^{-2-\kappa}\lambda^3$;

as $\lambda=\bar\lambda$ we get $O(\mu^{-2}h^{-2-\kappa})$.

Therefore we need to consider only zone

$$\{(x,y):|x-y|\le C_0\mu^{-1},\ |\nabla VF^{-1}|\ge\bar\lambda\}.$$

We are going to prove that

Proposition 8.2.2. *(cf. proposition 4.2.2). Let conditions* (0.5)–(0.6) *be fulfilled. Further, let* $\mu\le h^{-1}$. *Then for each* z *contribution of zone* $\{(x,y)\in B(z,\lambda(z)),\ |x-y|\le C_0\mu^{-1}|\}$ *with* $\lambda(z)\ge C_0\mu^{-1}$ *to* $|\mathsf{I}^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ *does not exceed* *(i)* $R^{\mathsf{W}}(\lambda)$ *which is defined as the right-hand expression of* (8.8) *or* (8.13) *for* $1<\kappa<3$*,* $0<\kappa\le 1$ *provided* $\lambda\ge (\mu h)^{\frac{1}{2}-\delta}$*;*

(ii) the right-hand expression of (8.17) *or* (8.27) *for* $1<\kappa<3$*,* $0<\kappa\le 1$ *provided* $\lambda\ge (\mu h)^{\frac{1}{2}}$*;*

(iii) the right-hand expression of (8.19) *or* (8.28) *for* $1<\kappa<3$*,* $0<\kappa\le 1$ *provided* $\lambda\le (\mu h)^{\frac{1}{2}}$*.*

Consider components of $R^{\mathsf{W}}(\lambda)$. Note that expressions (8.8), (8.13) and (8.20)–(8.22), (8.24), (8.26) contain λ in the positive powers while expressions (8.18), (8.25) contain λ in the negative powers. Therefore after integration with respect $\lambda^{-1}\,d\lambda$ we get expressions (8.8) and (8.13) as $\lambda=1$ (i.e. we get (8.14) and we get all other expressions as $\lambda=(\mu h)^{\frac{1}{2}}$; one can check easily that these latter expressions are less than (8.14) + $Ch^{-2-\kappa}$.

Therefore we arrive to

Theorem 8.2.3. *(cf. theorem 4.2.3). Let conditions* (0.5)*–*(0.6)*. Let* $\mu\le h^{-1}$*. Then*

(i) Under condition (0.7) *estimate*

$$|\mathsf{I}-\mathsf{I}^{\mathsf{W}}|\le CR^{\mathsf{W}}+Ch^{-2-\kappa}; \tag{8.31}$$

(ii) Under condition (0.8)

$$|\mathsf{I}-\mathsf{I}^{\mathsf{W}}|\le CR^{\mathsf{W}}+C\mu^{\frac{5}{2}}h^{-\frac{1}{2}-\kappa-\delta}+Ch^{-2-\kappa}. \tag{8.32}$$

Proof of proposition 8.2.2. First of all applying proposition 2.2.2 and the same arguments as in the analysis of section 2 we arrive to

(8.33) Contribution of the zone $\{|\sin(\theta(t'))|\ge C_0\hbar,\ |\sin(\theta(t''))|\ge C_0\hbar\}$ to the error $|\mathsf{I}^{\mathsf{T}}-\mathcal{I}^{\mathsf{W}}|$ does not exceed $R^{\mathsf{W}}(\lambda)$.

Therefore we need to explore contributions of three remaining zones

$$\{|\sin(\theta(t'))|\le 2C_0\hbar,\ |\sin(\theta(t''))|\le 2C_0\hbar\} \tag{4.43}$$

and

$$\{|\sin(\theta(t'))|\ge 2C_0\hbar,\ |\sin(\theta(t''))|\le C_0\hbar\} \tag{4.44}$$

(the same analysis would cover its tween zone $\{|\sin(\theta(t'))| \ge 2C_0\hbar,\ |\sin(\theta(t''))| \le C_0\hbar\}$).

We would like to follow the arguments of the proof of proposition 4.2.2 but there is a problem: the magnetic drift is controlled by α and it may be much less than λ. However let us apply few general arguments.

Zone (4.43) as $\lambda^2 \ge \mu h$. First, if we consider zone $\{(x,y):\ |x-y| \ge \ell\}$ (in the non-rescaled coordinates) we estimate the contribution of the corresponding pair (k',k'') by $C\hbar^2\ell^{-\kappa}|k'|^{-1}|k''|^{-1} \times h^{-d}\lambda^d$ and summation with respect to $|k'| \le \mu\lambda$, $|k''| \le \mu\lambda$ returns

$$C\mu^2 h^{2-d}\ell^{-\kappa}(1+|\log \mu\lambda|)^2. \tag{8.34}$$

Meanwhile, if we consider zone $\{(x,y):\ |x-y| \ge \ell\}$ (in the non-rescaled coordinates as well) then estimating $U(x,y,t)$ by h^{-d} (no rescaling, $d = 2,3$) we we estimate the contribution of the corresponding pair (k',k'') by $C\hbar^2 \times h^{-2d} \times \ell^{d-\kappa}\lambda^d$ and again summation with respect to k', k'' returns

$$C\mu^2 h^{2-2d}\ell^{d-\kappa}\lambda^d(1+|\log \mu\lambda|)^2. \tag{8.35}$$

Comparing (8.34) and (8.35) we see that their sum is minimal

$$C\mu^2 h^{2-d-\kappa}\lambda^d(1+|\log \mu\lambda|)^2 \tag{8.36}$$

as $\ell = h$.

Comparing with (8.8) and (8.13) we conclude that for $d = 3$ this is less than R^{W} as $0 < \kappa < 1$, $\lambda \ge \mu h$. To deal with $1 \le \kappa < 3$ let us note first that there must be a factor

$$C\bigl(1+|k'-k''|\lambda^2/\mu h\bigr)^{-l} \tag{8.37}$$

before summation. Really, as $|\xi_3+\eta_3| \ge \epsilon_0\lambda$ this factor is due to the shift in (x_3+y_3) and as $|\xi_3+\eta_3| \ge \epsilon_0\lambda$ but $|\nabla' V/F| \asymp \lambda$ or $|\nabla_{x_3} V/F| \asymp \lambda$ this factor is due to the shift in $(\xi_1+\eta_1)$ or $(\xi_3+\eta_3)$ respectively.

Therefore

(8.38) As $\lambda^2 \ge \mu h^{1-\delta}$ contribution of $k' \ne k''$ is negligible.

This allows us also eliminate logarithmic factors in (8.36) and (8.37).

Let us apply cut-off Q by hD_{x_3} and Q' by hD_{xy_3}. Assume first that either $|\xi_3| \le C\rho$ on the support of (the symbol of) Q or $|\eta_3| \le C\rho$ on the support of (the symbol of) Q', in our estimate we acquire factor ρ and we get an expression below $R^{\mathsf{W}}(\lambda)$.

Therefore we can assume that

$$|\xi_3|\ge \rho := \max\bigl((\mu h)^{\frac{1}{2}},\mu^{-1}\bigr) \text{ on the support of (the symbol of) } Q \tag{8.39}$$

(or $|\eta_3|\ge \max\bigl((\mu h)^{\frac{1}{2}},\mu^{-1}\bigr)$ on the support of (the symbol of) Q'.

Let us consider corresponding classical dynamics. Assumption (8.39) implies that in k winding it travels the "observable" distance $k\xi_3$ and if there was no "turn back" the contribution of k-th winding to estimate would not exceed

$$C\mu^2h^{-1}|k|^{-2-\kappa}\int_{|\xi_3|\ge\rho}|\xi_3|^{-\kappa}\,d\xi_3\lambda^3$$

where integral is $\asymp 1$ as $0<\kappa<1$, $\asymp \log\rho$ as $\kappa=1$ and $\asymp \rho^{1-\kappa}$ as $1<\kappa<3$. Obviously multiplied by μ^2h^{-1} it does not exceed $R^{\mathsf{W}}(\lambda)$ and therefore summation with respect to k returns $O(R^{\mathsf{W}}(\lambda))$.

Now we need to consider effect of the "turn back". Let $\beta=|\nabla_{x_3}V/F|$. Note first that as $|\xi_3|\ge\beta\lambda$ there is no turn back for time $|t|\le\epsilon\lambda$. Therefore we need to consider only $|\xi_3|\le\beta\lambda$.

There is come back on k-th winding as ξ_3 is close to $\zeta^*(k)\sim \mathsf{const}\beta\mu^{-1}k$ and the distance $|x-y|\asymp\mu^{-1}|k||\xi_3-\zeta(k)|$ and then contribution of k-th winding (as $|x-y|\ge Ch$) does not exceed

$$C\mu^{2+\kappa}h^{-1}|k|^{-2-\kappa}\int_{|\xi_3-\zeta(k)|\ge\mu h|k|^{-1}}|\xi_3-\zeta(k)|^{-\kappa}\,d\xi_3\lambda^3\le\\
C\lambda^3\begin{cases}\mu^{2+\kappa}h^{-1}|k|^{-2-\kappa} & 0<\kappa<2,\\ \mu^4h^{1-\kappa}|k|^{-4} & 2<\kappa<3,\end{cases}$$

with a logarithmic factor $(1+\log|k|+|\log\mu h|)$ as $\kappa=2$. The sum of the expression above is the same as this expression for $k=1$ (we denote it temporarily by $R\lambda^3$) which obviously less than $R^{\mathsf{W}}(\lambda)$.

Finally, we need to consider the contribution of k-th winding as $|x-y|\le h$. However it is done by the standard way as $U(x,y,t)$ after cut by ξ_3 and η_3 acquires factor Ch^2.

Therefore, under assumption $\lambda^2\ge\mu h^{1-\delta}$ we are done.

Case $\lambda^2\ge\mu h$. As $\lambda^2\ge\mu h$ (but $\lambda^2\not\gg\mu h$) terms with $k'\ne k''$ are no more negligible but in the estimate we can ignore due to the factor (8.37) them except when $k'=0$ or $k''=0$.

Applying the same arguments as in 2D-case i.e. replacing in term with $k'=0$ factor $|k'|^{-1}$ by factor $\hbar^{-1}$ we arrive to $R\lambda^3\times 1/(\mu h)\times (\mu h/\lambda^2)^l$ which is less than term (8.18) or (8.25) as $1<\kappa<3$ or $0<\kappa\le 1$ respectively.

Zone (4.43) as $\lambda^2\le \mu h$. In this case we apply the same arguments but now effectively we sum over k',k'' with $|k'-k''|\le \mu h/\lambda^2$ which effectively boils out to the summation over $k'=1$, $k''=1,\dots,\mu h/\lambda^2$ or $k'=0$, $k''=1,\dots,\mu h/\lambda^2$; the former results in $R\lambda^3\times \mu h/\lambda^2$ and the latter results $R\lambda^3\times \mu h/\lambda \times 1/(\mu h)$ and one can see easily that the result is less than what we claimed.

Zone (4.44) *as* $\lambda^2\ge \mu h$. Now there is always shift with respect to $(x-y)$ which is at least $\asymp |\sin(t')|$ and therefore as $1<\kappa<3$ contribution of this zone does not exceed contribution of zone (4.43). As $0<\kappa<1$ we get $C\mu^{\kappa+1}h^{-2}\lambda^3|k|^{-2}$ and summation returns $C\mu^{\kappa+1}h^{-2}\lambda^3$ which is the answer as $\lambda^2\gg \mu h$; let us denote it by $R\lambda^3$; for $\kappa=1$ it includes logarithmic factor $(1+|\log(\mu h)|)$.

The same answer estimates contribution of $k'\ne 0, k''\ne 0$ as $\lambda^2\ge \mu h$; contributions of $k'=0, k''\ne 0$ and $k''\ne 0, k'=0$ do not exceed $R\lambda^3\times 1/(\mu h)\times (\mu h/\lambda^2)^l$ and one can see easily that the result is less than what we claimed.

Zone (4.44) *as* $\lambda^2\le \mu h$. Again as $1<\kappa<3$ contribution of (4.44) does not exceed contribution of (4.43) and as $0<\kappa\le 1$ we estimate contribution of $k'\ne 0, k''\ne 0$ by $R\lambda^3\times \mu h/\lambda^2$, and contributions of $k'\ne 0, k''=0$ and $k'=0, k''\ne 0$ by $R\lambda^3\times \mu h/\lambda^2\times 1/(\mu h)$ and one can see easily that the result is less than what we claimed. □

8.3 Perturbations

Consider perturbations. However due to the presence of x_3, hD_3 there are rather different components here than in subsection 4.3.

8.3.1 Perturbations. I

Let us start from the terms $O(|x'-y'|)$. During evolution these terms are $O(\mu^{-1})$ and therefore we need to multiply by $O(\mu^{-2}h^{-1}|k|)$.

$1<\kappa<3$. Consider first case $k'=k''=k$. Assume that $\lambda^2\ge \mu h$. Recall that k-th tick' contribution to I^{T} was not exceeding expression (8.7) multiplied by $\mu^\kappa h^{-3}\lambda^3|k|^{-3}$ i.e.

$$C\lambda^3|k|^{-3}\left\{\begin{aligned}&\mu^{\kappa+1}h^{-2}\lambda^{1-\kappa} &&1<\kappa<2,\\ &\mu^3h^{-2}(1+|\log(\hbar/\lambda^2)|)\lambda^{-1} &&\kappa=2,\\ &\mu^3h^{-\kappa}\lambda^{-3+\kappa} &&2<\kappa<3\end{aligned}\right. \tag{8.40}$$

(as $|\sin(t')|\le\hbar$ or $|\sin(t'')|\le\hbar$ one should use proposition 8.2.2) and multiplying by $C\mu^{-2}h^{-1}|k|^{-1}$ we get after summation with respect to $k:|k|\ge 1$

$$C\lambda^3\left\{\begin{aligned}&\mu^{\kappa-1}h^{-3}\lambda^{1-\kappa} &&1<\kappa<2,\\ &\mu h^{-3}(1+|\log(\hbar/\lambda^2)|)\lambda^{-1} &&\kappa=2,\\ &\mu h^{-1-\kappa}\lambda^{-3+\kappa} &&2<\kappa<3\end{aligned}\right. \tag{8.41}$$

and summation with respect to λ returns the same expression as $\lambda=1$ which does not exceed $Ch^{-2-\kappa}$.

Next we need to consider $0\ne k'\ne k''\ne 0$. Recall that the contribution of such pair to I^{T} does not exceed

$$C\lambda^3|k'|^{-\frac{3}{2}}|k''|^{-\frac{3}{2}}(\mu h/\lambda^2|k'-k''|)^l\times \left\{\begin{aligned}&\mu^{\kappa+1}h^{-2}\lambda^{1-\kappa} &&1<\kappa<2,\\ &\mu^3h^{-2}(1+|\log(\mu h/\lambda^2)|)\lambda^{-1} &&\kappa=2,\\ &\mu^3h^{-\kappa}\lambda^{-3+\kappa} &&2<\kappa<3.\end{aligned}\right. \tag{8.42}$$

Multiplying by $C\mu^{-2}h^{-1}|k'|$ (as the perturbation goes to the first "factor") we get

$$C\lambda^3|k'|^{-\frac{1}{2}}|k''|^{-\frac{3}{2}}(\mu h/\lambda^2|k'-k''|)^l\times \left\{\begin{aligned}&\mu^{\kappa-1}h^{-3}\lambda^{1-\kappa} &&1<\kappa<2,\\ &\mu h^{-3}(1+|\log(\mu h/\lambda^2)|)\lambda^{-1} &&\kappa=2,\\ &\mu h^{-1-\kappa}\lambda^{-3+\kappa} &&2<\kappa<3.\end{aligned}\right. \tag{8.43}$$

Then summation with respect to k' of $|k'|^{-\frac{1}{2}}(\mu h/\lambda^2|k'-k''|)^l$ returns $|k''|^{-\frac{1}{2}}(\mu h/\lambda^2)^l$ as $\lambda^2\ge\mu h$ and summation with respect to k'' then returns the same expression as we got considering $k'=k''\ne 0$ with an extra factor $(\mu h/\lambda^2)^l$.

On the other hand, as $\lambda^2 \le \mu h$ one should replace (8.7) by $C(\mu h)^{\frac{3}{2}-\frac{1}{2}\kappa}$; multiplying by $\mu^\kappa h^{-3}\lambda^3|k'|^{-\frac{3}{2}}|k''|^{-\frac{3}{2}}(1+\lambda^2|k'-k''|/\mu h)^{-l} \times \mu^{-2}h^{-1}|k'|$ we get

$$C\mu^{-\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^3|k'|^{-\frac{1}{2}}|k''|^{-\frac{3}{2}}(1+\lambda^2|k'-k''|/\mu h)^{-l} \tag{8.44}$$

and one can see easily that summation with respect to k', k'' returns

$$C\mu^{-\frac{1}{2}+\frac{1}{2}\kappa}h^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^3(\mu h/\lambda^2)^{\frac{1}{2}} \asymp C\mu^{\frac{1}{2}\kappa}h^{-2-\frac{1}{2}\kappa}\lambda^2$$

and summation with respect to λ returns $C\mu^{\frac{3}{2}\kappa}h^{-1-\frac{1}{2}\kappa}$ which again is less than $Ch^{-2-\kappa}$.

If $k' \ne 0$, $k''=0$ we recall that $|k''|^{-\frac{3}{2}}$ should be replaced by $\hbar^{-\frac{3}{2}}$ in (8.42)–(8.43) and perturbation brings factor $\mu^{-2}h^{-1}|k'|$ so we get instead of (8.43)

$$C\lambda^3|k'|^{-\frac{1}{2}}\mu^{-\frac{3}{2}}h^{-\frac{3}{2}}(\mu h/\lambda^2|k'|)^l\times \\ \begin{cases} \mu^{\kappa-1}h^{-3}\lambda^{1-\kappa} & 1<\kappa<2,\\ \mu h^{-3}(1+|\log(\mu h/\lambda^2)|)\lambda^{-1} & \kappa=2,\\ \mu h^{-1-\kappa}\lambda^{-3+\kappa} & 2<\kappa<3\end{cases}$$

and summation with respect to $|k'|$ and then $\lambda \ge (\mu h)^{\frac{1}{2}}$ returns $C\mu^{\frac{1}{2}\kappa-\frac{1}{2}}h^{-\frac{5}{2}-\frac{1}{2}\kappa}$.

Similarly, as $\lambda^2 \le \mu h$ we start from (8.44) which is replaced by

$$C\mu^{-2+\frac{1}{2}\kappa}h^{-4-\frac{1}{2}\kappa}\lambda^3|k'|^{-\frac{1}{2}}(1+\lambda^2|k'|/\mu h)^{-l}$$

and summation with respect to k' and returns $C\mu^{-\frac{3}{2}+\frac{1}{2}\kappa}h^{-\frac{7}{2}-\frac{1}{2}\kappa}\lambda^2$ and then summation with respect to $\lambda \le (\mu h)^{\frac{1}{2}}$ returns the same answer $C\mu^{\frac{1}{2}\kappa-\frac{1}{2}}h^{-\frac{5}{2}-\frac{1}{2}\kappa}$.

As $|k'|<|k''|$ the estimate of the contribution of the pair (k',k'') to an error is obviously less than the same estimate for the pair (k'',k'); this takes care of $k'=0$, $k''\ne 0$. Finally for $k'=k''=0$ we can use the standard arguments to estimate the same way.

As the result we conclude that the error will not exceed $Ch^{-2-\kappa}$.

$0<\kappa\le 1$. The similar arguments work for $0<\kappa<1$. However one must reconsider zone $\{(x,y):\ |x-y|\ge C_0\mu^{-1}\}$ where contribution of the pair (k',k'') to Weyl asymptotics does not exceed

$$C\mu^2h^{-2}\zeta^{1-\kappa}|k|^{-3}\lambda^3 \asymp C\mu^{\kappa+1}h^{-2}|k|^{-2-\kappa}\lambda^3 \tag{8.45}$$

with $\zeta\asymp\mu^{-1}|k|$ as $k'=k''=k\ne 0$. Summation with respect to $k:|k|\ge 1$ returns then $C\mu^{\kappa+1}h^{-2}\lambda^3$ as it should.

As $k'=k''=k\ne 0$ it results in the approximation error not exceeding

$$C\mu^2h^{-2}\zeta^{1-\kappa}|k|^{-3}\lambda^3\times\mu^{-2}h^{-1}|k|\asymp C\mu^{\kappa-1}h^{-3}|k|^{-1-\kappa}\lambda^3 \tag{8.46}$$

and summation with respect to k,λ returns $C\mu^{\kappa-1}h^{-3}$.

As $0\ne k'\ne k'\ne 0$, $|k'|\ge|k''|$ it results in the approximation error not exceeding

$$\begin{multline}C\mu^2h^{-2}\bigl(\mu^{-1}|k'|)\bigr)^{1-\kappa}|k'|^{-\frac{3}{2}}|k''|^{-\frac{3}{2}}\times\\ (1+\lambda^2|k'-k''|/\mu h)^{-l}\lambda^3\times\mu^{-2}h^{-1}|k'|\asymp\\ C\mu^{\kappa-1}h^{-3}|k'|^{\frac{1}{2}-\kappa}|k''|^{-\frac{3}{2}}(1+\lambda^2|k'-k''|/\mu h)^{-l}\lambda^3\end{multline} \tag{8.47}$$

and the sum with respect to (k',k'') and $\lambda:\lambda^2\ge\mu h$ does not exceed $C\mu^{\kappa-1}h^{-2}$.

Meanwhile as $\lambda^2\le\mu h$ the sum with respect to (k',k'') does not exceed $C\mu^{\kappa-1}h^{-3}(\mu h/\lambda^2)^{\frac{3}{2}-\kappa}\lambda^3$ and the sum with respect to λ is less than $C\mu^{\kappa-1}h^{-3}$.

As $k'\ne 0$, $k''=0$ (8.47) becomes

$$C\mu^{\kappa-1}h^{-3}|k'|^{\frac{1}{2}-\kappa}(\mu h)^{-\frac{3}{2}}(1+\lambda^2|k'|/\mu h)^{-l}\lambda^3 \tag{8.48}$$

and then summation with respect to k' returns $C\mu^{\kappa-1}h^{-3}(\mu h)^{-\frac{3}{2}}(\mu h/\lambda^2)^l\lambda^3$ as $\lambda^2\ge\mu h$ and then the sum with respect to λ is less than $C\mu^{\kappa-1}h^{-3}$.

Meanwhile as $\lambda^2\le\mu h$ the sum with respect to k' does not exceed $C\mu^{\kappa-1}h^{-3}(\mu h/\lambda^2)^{\frac{3}{2}-\kappa}(\mu h)^{-\frac{3}{2}}\lambda^3$ and then the sum with respect to λ is less than $C\mu^{\kappa-1}h^{-3}$.

As $|k'|<|k''|$ the estimate of the contribution of the pair (k',k'') to an error is obviously less than the same estimate for the pair (k'',k'); this takes care of $k'=0$, $k''\ne 0$. Finally, for $k'=k''=0$ we can use the standard arguments to estimate contribution by $Ch^{-2-\kappa}$.

For $\kappa=1$ we apply the same arguments with the obvious modifications, concluding that the error does not exceed $Ch^{-3}(1+|\log\mu h|)$.

Note that in contrast to the case $1<\kappa<3$ now the error is larger than $Ch^{-2-\kappa}$.

To improve this remainder estimate let us notice that the second term in the approximation is 0 and the third term also acquires factor $\mu^{-2}h^{-1}|k|$ (as $k'=k''=k\ne 0$); so we get

$$C\mu^{\kappa-1}h^{-3}|k|^{-1-\kappa}\times\mu^{-2}h^{-1}|k|\lambda^3\asymp C\mu^{\kappa-3}h^{-4}|k|^{-\kappa}\lambda^3$$

which we should sum as $|k|\le \mu^2 h$; we also should sum

$$C\mu^2h^{-2}\zeta^{1-\kappa}|k|^{-3}\lambda^3 \asymp C\mu^{\kappa+1}h^{-2}|k|^{-2-\kappa}\lambda^3$$

as $|k|\ge \mu^2 h$ and both sums return $C\mu^{\kappa+1}h^{-2}(\mu^2h)^{-1-\kappa}\lambda^3\asymp C\mu^{-1-\kappa}h^{-3-\kappa}\lambda^3$. Summation with respect to λ returns $C\mu^{-1-\kappa}h^{-3-\kappa}$. One can estimate contribution of all all other pairs (k',k'') in the same way. As $\kappa=1$ we get $C\mu^{-2}h^{-4}(1+|\log \mu h|)$.

So we arrive to

Theorem 8.3.1. *(cf. theorem 4.3.6). Let conditions* (0.5)–(0.6) *and either* (0.7) *or* (0.8) *be fulfilled. Let* $h^{-\frac{1}{2}}\mu\le h^{-1}$. *Then*

(i) As $1<\kappa<3$

$$|\mathsf{I}-\mathsf{I}^0|\le Ch^{-2-\kappa} \tag{8.49}$$

where

$$\mathsf{I}^0 := \int \Omega_{1,1}\bigl(\frac{1}{2}(x+y),x-y\bigr)\cdot |e^0_{\frac{1}{2}(x+y)}(x,y,0)|^2\,dxdy \tag{8.50}$$

and $e^0_z(x,y,\tau)$ *is the Schwartz kernel of the spectral projector of the generalized pilot-model operator*

$$A^0_z := h^2D_1^2+(hD_2-\mu x_1)^2+h^2D_3^2+V^0(x),\qquad V^0(x):=V(z_1,z_2,x_3) \tag{8.51}$$

which in contrast to (6.36) V^0 *does not contain linear terms with respect to* (x_1,x_2);

(ii) As $0<\kappa<1$

$$|\mathsf{I}-\mathsf{I}^0|\le Ch^{-2-\kappa}+C\mu^{-1-\kappa}h^{-3-\kappa} \tag{8.52}$$

and as $\kappa=1$

$$|\mathsf{I}-\mathsf{I}^0|\le Ch^{-3}+C\mu^{-2}h^{-4}|\log\mu h|. \tag{8.53}$$

Remark 8.3.2. Note that the remainder estimate does not exceed $Ch^{-2-\kappa}$ for $\mu\ge h^{-1/(1+\kappa)}$ as $0<\kappa<1$ and for $\mu\ge h^{-\frac{1}{2}}(1+|\log\mu h|)^{\frac{1}{2}}$ as $\kappa=1$.

8.3.2 Perturbations. II

If we go from the generalized pilot-model approximation to the magnetic Weyl approximation then we need to counter $(x_3 - y_3)\lambda$ terms and this is $O(\mu^{-1}|k'|\lambda)$ due to the finite speed of propagation. Then dealing as before we acquire an extra factor $|k'|\lambda$.

$1 < \kappa < 3$. Then as $k' = k'' = k \ne 0$ we have $|k|^{-1}$ rather than $|k|^{-2}$ and the sum with respect to k contains an extra logarithmic factor $(1+\log(\mu^2 h))\lambda$ in comparison to what we got in subsubsection 8.3.1 "Perturbations. I".

Consider case $0 \ne k' \ne k'' \ne 0$, $|k'| \ge |k''|$. Then as $\lambda^2 \ge \mu h$ we arrive to the same expression (but with an extra factor $(\mu h/\lambda^2)^l$). However as $\lambda^2 \le \mu h$ we need to consider $|k'|^{-\frac{3}{2}}|k''|^{-\frac{3}{2}} \times |k'|^2\lambda$ with $k', k'' : |k' - k''| \le \mu h/\lambda^2$ which in comparison to what we got before after summation returns an extra factor $\mu h/\lambda^2 \times \lambda$ and the sum with respect to λ does not exceed what we got above.

Consider $0 \ne k'$, $k'' = 0$. Again no change as $\lambda^2 \ge \mu h$ and an extra factor $(\mu h/\lambda^2) \times \lambda$ as $\lambda^2 \le \mu h$ and the sum with respect to λ does not exceed what we got above.

Therefore, as a result we get $C\mu^{\kappa-1}h^{-3}(1 + |\log \mu^2 h|)$ as $1 < \kappa < 2$ and $C\mu h^{-1-\kappa}(1 + |\log \mu^2 h|)$ as $2 < \kappa < 3$ with an extra factor $(1 + |\log \mu h|)$ as $\kappa = 2$. In both cases it is less than $Ch^{-2-\kappa}$ unless $\mu \ge h^{-1}|\log h|^{-r}$.

To improve our result in this latter case we apply three term approximation and note that the second term is 0 while the third term contains summation of $R^{\mathsf{W}}\lambda^3|k|^{-3} \times (\mu^{-2}h|k|^2\lambda)^2$ for $k : |k| \le \mu h^{\frac{1}{2}}$ and summation of $R^{\mathsf{W}}\lambda^3|k|^{-3}$ for $k : |k| \ge \mu h^{\frac{1}{2}}$ as $k' = k'' = k \ne 0$ where $R^{\mathsf{W}} = \mu^3 h^{-\kappa}$ as $2 < \kappa < 3$, $R^{\mathsf{W}} = \mu^{\kappa+1}h^{-2}$ as $1 < \kappa < 2$ with an extra factor $(1 + |\log \mu h|)$ as $\kappa = 2$.

Other pairs (k', k'') should be modified in the same way and we leave easy but tedious details to the reader. As a result we get $R^{\mathsf{W}}\mu^{-2}h^{-1}$ without any logarithmic factor thus arriving to the same estimate as before but without logarithmic factor; as $1 < \kappa < 3$ it does not exceed $Ch^{-2-\kappa}$. So we arrive to statement (i) of theorem 8.3.3 below.

$0 < \kappa \le 1$. Now expression (8.46) acquires factor $|k|\lambda$ and becomes $C\mu^{\kappa-1}h^{-3}|k|^{-\kappa}\lambda^4$ and summation with respect to $k : |k| \le \mu h^{\frac{1}{2}}\lambda^{-\frac{1}{2}}$ returns

$$C\mu^{\kappa-1}h^{-3}(\mu h^{\frac{1}{2}}\lambda^{-\frac{1}{2}})^{1-\kappa}\lambda^4 \asymp Ch^{-\frac{5}{2}-\frac{1}{2}\kappa}\lambda^{-\frac{7}{2}+\frac{1}{2}\kappa}.$$

We also should sum $C\mu^{\kappa+1}h^{-2}|k|^{-2-\kappa}\lambda^3$ for $|k|\ge \mu h^{\frac{1}{2}}\lambda^{-\frac{1}{2}}$ with the same result.

Finally summation with respect to λ returns $Ch^{-\frac{5}{2}-\frac{1}{2}\kappa}$.

Other pairs (k', k'') should be modified in the same way and we leave easy but tedious details to the reader. Three term approximation does not improve results. As $\kappa=1$ we acquire factor $(1+|\log h|)$. So we arrive to statement (ii) of theorem 8.3.3 below.

Theorem 8.3.3. *(cf. theorem 4.3.4). Let conditions* (0.5)–(0.6) *and either* (0.7) *or* (0.8) *be fulfilled. Let* $h^{-\frac{1}{2}}\le \mu\le h^{-1}$. *Then*

(i) As $1<\kappa<3$

$$|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|\le Ch^{-2-\kappa}; \tag{8.54}$$

(ii) As $0<\kappa<1$

$$|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|\le Ch^{-\frac{5}{2}-\frac{1}{2}\kappa} \tag{8.55}$$

and as $\kappa=1$

$$|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|\le Ch^{-3}(1+|\log \mu h|). \tag{8.56}$$

8.3.3 Perturbations. III

Now in the case $0<\kappa\le 1$ we consider the pilot-model approximation. As (8.55) is larger than (8.53) we consider improvement of the previous subsubsection only. Now as perturbation is $(x_3-y_3)^2$ which translates into $\mu^{-2}|k'|^2$ rather than μ^{-1} expression (8.46) acquires factor $\mu^{-1}|k'|^2$ and becomes $C\mu^{\kappa-2}h^{-3}|k|^{1-\kappa}\lambda^3$ and we need to sum this as $|k|\le \mu h^{\frac{1}{3}}$ resulting in

$$C\mu^{\kappa-2}h^{-3}(\mu h^{\frac{1}{3}})^{2-\kappa}\lambda^3 \asymp Ch^{-\frac{7}{3}-\frac{1}{3}\kappa}\lambda^3.$$

We should sum $C\mu^{\kappa+1}h^{-2}|k|^{-2-\kappa}\lambda^3$ for $|k|\ge \mu h^{\frac{1}{3}}$ with the same result.

Finally, summation with respect to λ returns $Ch^{-\frac{7}{3}-\frac{1}{3}\kappa}$.

Other pairs (k', k'') should be modified in the same way and we leave easy but tedious details to the reader. Three term approximation does not improve results. As $\kappa=1$ we acquire factor $(1+|\log h|)$. So we arrive to

Theorem 8.3.4. *Let conditions* (0.5)–(0.6) *and either* (0.7) *or* (0.8) *be fulfilled. Let* $h^{-\frac{1}{2}} \le \mu \le h^{-1}$*. Then as* $0 < \kappa < 1$

$$|\mathsf{I} - \bar{\mathsf{I}}| \le Ch^{-2-\kappa} + Ch^{-\frac{7}{3}-\frac{1}{3}\kappa} + C\mu^{-1-\kappa}h^{-3-\kappa} \tag{8.57}$$

and as $\kappa = 1$

$$|\mathsf{I} - \bar{\mathsf{I}}| \le Ch^{-3} + C\mu^{-2}h^{-4}(1 + |\log \mu h|) \tag{8.58}$$

where $\bar{\mathsf{I}}$ *is defined for a simplified pilot-model operator*

$$\begin{aligned} \bar{A}_z := h^2D_1^2 + (hD_2 - \mu x_1)^2 + h^2D_3^2 + \bar{V}(x),\\ \bar{V}(x) := V(z) + \partial_{z_3}V(z)(x_3 - z_3). \end{aligned} \tag{8.59}$$

8.4 Superstrong magnetic field

Consider pilot-model Schrödinger-Pauli operator (1.72). Recall that according to propositions 7.2.1(ii), 7.2.2(ii) respectively

$$|\mathsf{I}| \le C\mu h^{-2} \left\{\begin{aligned} &h^{-\kappa} &&\text{as}\ \ 0 < \kappa < 1,\\ &h^{-1}\bigl(1 + (\log \mu h)_+\bigr) &&\text{as}\ \ \kappa = 1,\\ &\mu^{\frac{1}{2}(\kappa-1)}h^{-\frac{1}{2}(\kappa+1)} &&\text{as}\ \ 1 < \kappa < 3 \end{aligned}\right. \tag{7.2}$$

and

$$\begin{aligned} |\mathsf{I} - \mathsf{I}^{\mathsf{T}}| \le \\ C\mu h^{-1} \left\{\begin{aligned} &h^{-\kappa} &&\text{as}\ \ 0 < \kappa < 1,\\ &h^{-1}\bigl(1 + (\log \mu h)_+\bigr) &&\text{as}\ \ \kappa = 1,\\ &\mu^{\frac{1}{2}(\kappa-1)}h^{-\frac{1}{2}(\kappa+1)} &&\text{as}\ \ 1 < \kappa < 3,\ \ \kappa \ne 2,\\ &\mu^{\frac{1}{2}}h^{-\frac{3}{2}}\bigl(1 + (\log \mu h)_+\bigr) &&\text{as}\ \ \kappa = 2. \end{aligned}\right. \end{aligned} \tag{7.3}$$

Let us assume that $F = 1$ to avoid some unpleasant correction terms as reduced to $F = 1$.

(i) Consider the generalized pilot-model approximation (8.51). Then $\sigma \asymp h$ and under condition

$$|V + 2m\mu h| \ge \epsilon_0 \qquad \forall m \in \mathbb{Z}^+ \tag{8.60}$$

the shift with respect to x_3 is $\asymp |t|$ which implies that we need to consider only $t'\asymp t''$ as the total contribution of other intervals will be less, and that contribution of intervals $|t'|\le C_0h$, $|t''|\le C_0h$ does not exceed the right-hand of (8.2) multiplied by Ch, and that contribution of intervals $t'\asymp T$, $t''\asymp T$ ($T\ge C_0h$) does not exceed $C\mu h^{-1}\times T^{-\kappa}$ so summation with respect to $T^{-1}dT$ returns $C\mu h^{-1-\kappa}$.

Then the error does not exceed the right-hand expression of (8.3).

(ii) Consider the magnetic Weyl approximation. In this case $\sigma\asymp T$ and again under assumption (8.60) we can consider only $t'\asymp t''$, and we estimate contribution of intervals $|t'|\le C_0h$, $|t''|\le C_0h$ as above, but contribution of intervals $t'\asymp T$, $t''\asymp T$ ($T\ge C_0h$) does not exceed $C\mu h^{-2}\times T^{1-\kappa}$ and summation with respect to $T^{-1}dT$ returns $C\mu h^{-1-\kappa}$ as $1<\kappa<3$. As $0<\kappa<1$ we sum this way only for $T\le h^{\frac{1}{2}}$ resulting in $C\mu h^{-\frac{3}{2}-\frac{1}{2}\kappa}$ and we sum $C\mu h^{-1}T^{-\kappa}$ for $T\ge h^{\frac{1}{2}}$ with the same result. As $\kappa=1$ we get $C\mu h^{-2}(1+|\log h|)$.

(iii) Consider the pilot-model approximation as $0<\kappa<1$. In this case $\sigma\asymp T^2$ (as we pass from the generalized pilot-model approximation) and again under assumption (8.60) we can consider only $t'\asymp t''$, and we estimate contribution of intervals $|t'|\le C_0h$, $|t''|\le C_0h$ as above, but contribution of intervals $t'\asymp T$, $t''\asymp T$ ($T\ge C_0h$) does not exceed $C\mu h^{-2}\times T^{2-\kappa}$ summation with respect to $T^{-1}dT$ returns $C\mu h^{-\frac{4}{3}-\frac{1}{3}\kappa}$ as we sum with respect to $T^{-1}dT$ for $T\le h^{\frac{1}{3}}$; for $T\ge h^{\frac{1}{3}}$ we sum $C\mu h^{-1}T^{-1-\kappa}$ with the same result.

Then we arrive to

Theorem 8.4.1. *(i) For the Schrödinger-Pauli operator under conditions* (0.5)–(0.6) *and* (8.60)

$$|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|\le C\begin{cases}\mu h^{-\frac{3}{2}-\frac{1}{2}\kappa} & 0<\kappa<1,\\ \mu h^{-2}(1+|\log h|) & \kappa=1,\\ \mu^{\frac{1}{2}(\kappa+1)}h^{-\frac{1}{2}(\kappa+3)} & 1<\kappa<3,\ \kappa\ne 2,\\ \mu^{\frac{3}{2}}h^{-\frac{5}{2}}(1+(\log\mu h)_+) & \kappa=2.\end{cases}\tag{8.61}$$

(ii) Further, as $0<\kappa\le 1$

$$|\mathsf{I}-\mathsf{I}^0|\le C\begin{cases}\mu h^{-1-\kappa} & 0<\kappa<1,\\ \mu h^{-2}(1+(\log\mu h)_+) & \kappa=1\end{cases}\tag{8.62}$$

and

$$|\mathsf{I}-\bar{\mathsf{I}}|\le C\begin{cases}\mu h^{-1-\kappa}+\mu h^{-\frac{4}{3}-\frac{1}{3}\kappa} & 0<\kappa<1,\\ \mu h^{-2}(1+|\log h|) & \kappa=1\end{cases}\tag{8.63}$$

Let us get rid off condition (8.60). First, we still can assume without any loss of the generality that (8.60) holds for all $m\in\mathbb{Z}^+$ save $m=\bar{m}$. Then making ℓ-admissible partitions in x and y with

$$\ell=\epsilon|V+2\bar{m}\mu h|+h^{\frac{2}{3}}\tag{8.64}$$

one can replace (8.60) by (0.7) and then making ℓ-admissible partitions in x and y with

$$\ell=\epsilon\bigl(|V+2\bar{m}\mu h|+|\nabla V|^2\bigr)^{\frac{1}{2}}+h^{\frac{1}{2}}\tag{8.65}$$

one can replace (0.7) by (0.8) arriving to

Theorem 8.4.2. *Theorem 8.4.1 remains valid with assumption* (8.60) *replaced by* (0.8)*.*

Easy but tedious details are left to the reader.

8.5 Problems and remarks

Remark 8.5.1. The main difference between cases $2<\kappa<3$, $1<\kappa<2$ and $0<\kappa<1$ that in the first case the main contribution to the remainder is delivered by (x,y) close to one another ($|x-y|\ll\mu^{-1}$), in the second case by (x,y) with $|xy|\asymp\mu^{-1}$ and in the third case by (x,y) with $|xy|\asymp 1$.

We can calculate $\mathcal{I}^{\mathsf{MW}}$ and $\bar{\mathsf{I}}$ plugging corresponding expressions $e_x^{\mathsf{MW}}(x,x,0)$ and $\bar{e}_x(x,x,0)$ into I.

Problem 8.5.2. *Find nice expressions for $\mathcal{I}^{\mathsf{MW}}$ and $\bar{\mathsf{I}}$.*

Problem 8.5.3. *As $\mu h\le 1$ get rid off condition (0.5); according to Chapter 13 we do not need it for estimate $|\mathsf{I}-\mathsf{I}^{\mathsf{T}}|$ but we want to get rid off it in estimates for $|\mathsf{I}-\mathcal{I}^{\mathsf{W}}|$, $|\mathsf{I}-\mathcal{I}^{\mathsf{MW}}|$ and $|\mathsf{I}-\bar{\mathsf{I}}|$. To do this*

(i) Assume first that condition (0.7) is fulfilled and consider the scaling function $\ell(x)=\max\bigl(\epsilon|V(x)|,\mu h, C\mu^{-1}\bigr)$.

(ii) Assuming then that condition (0.8) is fulfilled and consider the scaling function $\ell(x) = \max\bigl(\epsilon(|V(x)| + |\nabla V(x)|^2)^{\frac{1}{2}}, (\mu h)^{\frac{1}{2}}, C\mu^{-1}\bigr)$.

Problem 8.5.4. *Weaken non-degeneracy condition (0.8) which seems to be excessive here (in contrast to* 2D *case).*

Problem 8.5.5. *Get rid off condition (0.6) assuming instead that* $|\mathbf{F}| + |\nabla\otimes\mathbf{F}| \ge \epsilon$.

9 Estimates of expression (0.4)

9.1 Basics

In the multiparticle quantum theory we need to calculate expression (0.2) approximating energy of the electron-electron interaction or, more often, to estimate expression

$$\mathsf{K} := \iint \bigl(e(x,x,\tau) - h^{-d}\mathcal{N}_x(\tau)\bigr)\bigl(e(y,y,\tau) - h^{-d}\mathcal{N}_x(\tau)\bigr)\omega(x,y)\,dxdy \tag{0.4}$$

where $\omega(x,y) = \omega_\kappa(x,y)$ satisfies assumption (0.3) and $\mathcal{N}$ is *some* approximation. We concentrate on (0.4); asymptotics of (0.2) could be proven by the same arguments and we leave them to the reader.

Applying partition of unity we note that both of them depend on asymptotics of expression

$$\Gamma(e\psi) = \int e(x,x,\tau)\psi_\gamma(x)\,dx \tag{9.1}$$

with γ-admissible function ψ_γ.

Let us denote by $\gamma^d R(\alpha,\gamma)$ an upper estimate of an error occurring when we replace in (9.1) $e(x,x,0)$ by $h^{-d}\mathcal{N}$ and let

$$M = Ch^{-d} + C\mu h^{1-d} \tag{9.2}$$

be an upper estimate for $e(x,x,0)$.

Proposition 9.1.1. *Under condition* (0.7)

$$|\mathsf{K}| \le C\int \bigl(R(1,\gamma)\bigr)^2\gamma^{d-1-\kappa}\,d\gamma \tag{9.3}$$

and

$$|\mathsf{J}-\mathcal{J}| \le CM\int R(1,\gamma)\gamma^{d-1-\kappa}\,d\gamma \tag{9.4}$$

and under condition (0.8)

$$\begin{multlined}|\mathsf{K}| \le C\iint_{\{\gamma\le\alpha\}} \bigl(R(\alpha,\gamma)\bigr)^2\gamma^{d-1-\kappa}\alpha^{d-1}\,d\gamma d\alpha+\\ C\iint_{\{\alpha\le\beta\}} R(\alpha,\alpha)R(\beta,\beta)\alpha^{d-1}\beta^{d-1-\kappa}\,d\alpha d\beta\end{multlined} \tag{9.5}$$

and

$$\begin{multlined}|\mathsf{J}-\mathcal{J}| \le CM\iint_{\{\gamma\le\alpha\}} R(\alpha,\gamma)\gamma^{d-1-\kappa}\alpha^{d-1}\,d\gamma d\alpha+\\ CM\iint_{\{\alpha\le\beta\}} R(\alpha,\alpha)\alpha^{d-1}\beta^{d-1-\kappa}\,d\alpha d\beta\end{multlined} \tag{9.6}$$

where for $\alpha \le C_0\mu^{-1}$ *we need to modify* $\alpha^{-1}R(\alpha,\gamma)$ *(see below)*[22]*.*

Proof. Due to representation (3.1) and definition of $R(\alpha,\gamma)$ and M contribution of the zone $\{(x,y):\ |x-z|\asymp\gamma, |y-z|\asymp\gamma\}$ (note that $R(\alpha,\gamma)\le M$) to K does not exceed $C(R(\alpha,\gamma))^2\gamma^{2d-\kappa}$ for fixed z and γ where $\alpha=\alpha(z)$ provided $\gamma\le 3\epsilon\alpha$.

Under assumption (0.7) summation over z results in $C(R(1,\gamma))^2\gamma^{d-\kappa}$ and summation over γ results in estimate (9.3); estimate (9.4) is proven in the same way.

On the other hand, under assumption (0.8) summation over $z:\alpha(z)\asymp\alpha$ results in $C(R(\alpha,\gamma))^2\alpha^d\gamma^{d-\kappa}$ and summation over γ, and then over α results the first term in the right-hand expression of (9.5); it estimates contribution of zone $\{(x,y):\ |x-y|\le 3\epsilon\alpha\alpha(x)\}$ (in which $\alpha(y)\asymp\alpha(x)$).

---

[22] As $d=3$ we denote by α what we used to denote by γ in sections 5–8 (i.e. $(\alpha^2+\beta^2)^{\frac12}$ in these sections notations.

Further, under assumption (0.8) if $|x-y|\ge 2\epsilon\alpha$, $\alpha:=\alpha(x)$, $\beta:=\alpha(y)$ then $|x-y|\asymp \max(\alpha,\beta)$. Therefore as $\beta\ge\alpha$ the contribution of zone

$$\{(x,y):|x-y|\asymp\beta,\alpha(y)\asymp\alpha,\alpha(y)\asymp\beta\}$$

does not exceed $CR(\alpha,\alpha)\alpha^d\times R(\beta,\beta)\beta^{d-\kappa}$ and summation over α, β returns the second term in the in the right-hand expression of (9.5); it estimates contribution of zone $\{(x,y):\ |x-y|\ge\epsilon\bigl(\alpha(x)+\alpha(y)\bigr)\}$. $\square$

9.2 Case $d=2$

9.2.1 Tauberian asymptotics

(i) Consider first case $\mu\le h^{-1}$. Now according to (1.58) we need just to plug

$$R(\alpha,\gamma)=\mu^{-1}h^{-1}+\alpha^{-1}R^{\mathsf{T}}(\alpha,\gamma) \tag{9.7}$$

as $h^{-2}\mathcal{N}_x=e^{\mathsf{T}}(x,x,0)$. Obviously the effect of $C\mu^{-1}h^{-1}$ term to both (9.3) and (9.5) will be term $C\mu^{-2}h^{-2}$.

Plugging into (9.3) or (9.5) $R(\alpha,\gamma)=\alpha^{-1}R^{\mathsf{T}}(\alpha,\gamma)$ with $R^{\mathsf{T}}(\alpha,\gamma)$ defined by (1.57) we obviously eliminate from integration the last two zones (defined in (1.57)) as a power of γ is negative [23] and the first one as a power of γ is positive. Thus γ snaps to its minimal value in the last two zones and to its maximal value in the first one, and only the second zone matters.

Further, in virtue of (1.57) $R^{\mathsf{T}}(\alpha,\gamma)=C\mu^{\frac12}\gamma^{-\frac12}$ in the remaining second zone $\{h\le\gamma\le\min(\mu h/\alpha,\mu^{-1})\}$.

Then the power of γ is positive, 0 or negative as $0<\kappa<1$, $\kappa=1$ or $1<\kappa<2$ respectively. Thus γ snaps to $\min(\mu h/\alpha,\mu^{-1})$ as $0<\kappa<1$ and γ snaps to h as $1<\kappa<2$.

Then (9.3) becomes

$$|\mathsf{K}^{\mathsf{T}}|\le C\mu^{-2}h^{-2}+C\begin{cases}\mu h^{1-\kappa} & 1<\kappa<2,\\ \mu\min(|\log\mu h|,|\log\mu|) & \kappa=1,\\ \mu\min\bigl((\mu h)^{1-\kappa},\mu^{-1+\kappa}\bigr) & 0<\kappa<1.\end{cases} \tag{9.8}$$

Consider the first term in the right-hand expression of estimate (9.5). If $1<\kappa<2$ as we mentioned γ snaps to h thus resulting in the integrand $C\mu h^{1-\kappa}$ and in $C\mu h^{1-\kappa}|\log h|$ after integration with respect to $\alpha^{-1}d\alpha$.

---

[23] Here and below talking about powers of α or β or γ we mean in the integral with respect to $\gamma^{-1}d\gamma$ or $\alpha^{-1}d\alpha$ or $\beta^{-1}d\beta$.

Let $0 < \kappa \le 1$. Then γ snaps to $\gamma(\alpha) := \min(\mu h/\alpha, \mu^{-1})$, thus resulting in the integrand $C\mu \min\bigl((\mu h/\alpha)^{1-\kappa}, \mu^{\kappa-1}\bigr)$ as $0 < \kappa < 1$ and in $C\mu \log \min\bigl(\mu/\alpha, (\mu h)^{-1}\bigr)\bigr)$ as $\kappa = 1$.

Then after integration with respect to $\alpha^{-1}d\alpha$ we get $O(\mu^{\kappa}|\log h|)$ as $0 < \kappa < 1$ or $O(\mu^{\kappa}|\log h| \cdot |\log(\mu h)|)$ as $\kappa = 1$ and this is less than the estimate of the contribution of zone (9.9) below unless $\mu \ge h^{-1}|\log h|^{-1}$.

However in (9.5) we need also calculate the contribution of the zone

$$\{(x,y):\ \alpha(x) \le C_0\mu^{-1},\ \alpha(y) \le C_0\mu^{-1}\} \tag{9.9}$$

which is not covered by our arguments and also the second term.

The former obviously does not exceed $C\mu^2h^{-2} \times \mu^{-4+\kappa} = C\mu^{-2+\kappa}h^{-2}$ where the second factor is $\int |x-y|^{-\kappa}dxdy$ over zone (9.9) and this also covers zone $\{\beta \le C_0\mu^{-1}\}$ in the second term of (9.5). Obviously $C\mu^{-2+\kappa}h^{-2}$ is larger than anything we got before.

Consider the second term in the right-hand expression of (9.5). Here we have $R^{\mathsf{T}}(\alpha,\alpha) = \alpha^{-1}$ as $(\mu h)^{\frac{1}{2}} \ge \alpha \ge \mu^{-1}$ and $\mu^{-1}\alpha^{-1}h^{-1}\min\bigl((\mu^2h/\alpha)^{\frac{1}{2}}, 1\bigr) \times (\mu h/\alpha)^{l}$ as $(\mu h)^{\frac{1}{2}} \le \alpha \le 1$ and therefore β is in the negative power and snaps to α; so we arrive to

$$\int \bigl(R^{\mathsf{T}}(\alpha,\alpha)\bigr)^2\alpha^{1-\kappa}\,d\alpha.$$

In this integral α is in the negative power and snaps to μ^{-1} resulting in $\bigl(R^{\mathsf{T}}(\mu^{-1},\mu^{-1})\bigr)^2\mu^{\kappa-2} = O(\mu^{\kappa})$ which is less than we have already.

Finally, we need to consider contribution of zone

$$\{(x,y):\ \alpha(x) \le C_0\mu^{-1}, \alpha(y) \ge C_0\mu^{-1}\}. \tag{9.10}$$

So, we get $C\mu^{-2} \times \mu h^{-1}\int R^{\mathsf{T}}(\beta,\beta)\beta^{1-\kappa}\,d\beta$; in this integral β is in the negative power for sure as $1 < \kappa < 2$ and it snaps to $\beta = \mu^{-1}$ resulting in $C\mu^{-1}h^{-1}R^{\mathsf{T}}(\mu^{-1},\mu^{-1})\mu^{\kappa-2}$ which is less than we got already.

As $0 < \kappa < 1$, β is also in the negative power except zone $\{\mu^{-1} \le \beta \le (\mu h)^{\frac{1}{2}}\}$ which is possible as $\mu \ge h^{-\frac{1}{3}}$; then as β snaps to $(\mu h)^{\frac{1}{2}}$ resulting in $C(\mu h)^{-\frac{1}{2}-\frac{1}{2}\kappa}$ and one can see easily that it is less than what we got already. The same is true for $\kappa = 1$ as well.

So we arrive to estimate

$$|\mathsf{K}^{\mathsf{T}}| \le C\mu^{-2+\kappa}h^{-2} + C|\log h|\mu\bigl(\mu^{\kappa-1} + h^{1-\kappa}\bigr). \tag{9.11}$$

(ii) Let $\mu \ge h^{-1}$. Now we need just to plug $R(\alpha,\gamma)=\alpha^{-1}R^{\mathsf{T}}(\alpha,\gamma)\ge 1$ with $R^{\mathsf{T}}(\alpha,\gamma)$ delivered by (1.76). Then in (9.3) we need to snap γ to $\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ and set $R(\gamma)=\mu^{\frac{1}{2}}h^{-\frac{1}{2}}$. So (9.3) becomes

$$|\mathsf{K}^{\mathsf{T}}|\le C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}. \tag{9.12}$$

Meanwhile, under assumption $(0.8)'$ in the first term of the right-hand expression of (9.5) α is in power 0 and logarithmic factor appears; we get $C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}|\log\mu|$.

Further, in (9.5) we need to estimate also contribution of zones

$$\{(x,y):\ \alpha(x)\le C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}},\alpha(y)\le C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}\}, \tag{9.13}$$

$$\{(x,y):\ \alpha(x)\le C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}},\alpha(y)\ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}\} \tag{9.14}$$

and of the second term in the right-hand expression and one can see easily that these contributions do not exceed $C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}|\log\mu|$; and so we get

$$|\mathsf{K}^{\mathsf{T}}|\le C\mu^{\frac{1}{2}\kappa}h^{-\frac{1}{2}\kappa}|\log\mu|. \tag{9.15}$$

Thus we proved

Proposition 9.2.1. *(i) For $\mu\le h^{-1}$ under condition* (0.7) *estimate* (9.8) *holds and under condition* (0.8) *estimate* (9.11) *holds;*

(ii) Let $\mu\ge h^{-1}$. Then under condition $(0.7)'$ *estimate* (9.12) *holds and under condition* $(0.8)'$ *estimate* (9.15) *holds.*

Problem 9.2.2. *Prove that estimates* (9.8) *and* (9.12) *hold under conditions* $(0.8)^{+}$ *and* $(0.8)^{+\prime}$ *respectively.*

9.2.2 Weyl approximation

Now we should add to what we got for K^{T} corresponding expressions with $R(\alpha,\gamma)=R^{\mathsf{W}}(\alpha,\gamma)$ calculated according to (1.80)–(1.81) and therefore γ must be snapped to $\min(\alpha,\mu h\alpha^{-1})$. Then (9.3) becomes

$$\begin{aligned}|\mathsf{K}^{\mathsf{W}}|\le C\mu^{-2}h^{-2}+C(R^{\mathsf{W}})^2(\mu h)^{2-\kappa}=\\ C\mu^{-2}h^{-2}+C\begin{cases}\mu^4h^{-1}(\mu h)^{2-\kappa} & \text{as }\ \mu\le h^{-\frac{1}{2}},\\ \mu^2h^{-2}(\mu h)^{2-\kappa} & \text{as }\ h^{-\frac{1}{2}}\le\mu\le h^{-1}\end{cases}\end{aligned} \tag{9.16}$$

which obviously is larger than the right-hand expressions of (9.8) and so we can neglect the Tauberian error.

Consider the first term in the right-hand expression of (9.5). Note that for $\alpha \le (\mu h)^{\frac{1}{2}}$ γ snaps to α and then α is in the positive power. Also note that as $\alpha \le \mu^2 h$ (1.80) defines $R^{\mathsf{W}} = \mu h^{-1}$ and α is in the positive power as well. Therefore as $h^{-\frac{1}{2}} \le \mu \le h^{-1}$ we snap α to 1 resulting in $C\mu^2 h^{-2}(\mu h)^{2-\kappa}$ prescribed by (9.16) and $\mu \le h^{-\frac{1}{2}}$ we need to consider interval $\max(\mu^{-1}, \mu^2 h) \le \alpha \le 1$; then automatically $\alpha \ge (\mu h)^{\frac{1}{2}}$ and we arrive to

$$C\int \mu^4 h^{-1}(\mu h/\alpha)^{2-\kappa} d\alpha. \tag{9.17}$$

Here as $1 < \kappa < 2$, α is in the positive power and snaps to 1 resulting in $C\mu^4 h^{-1}(\mu h)^{2-\kappa}$ prescribed by (9.16).

On the other hand, as $0 < \kappa < 1$, α is in the negative degree and snaps to $\max(\mu^{-1}, \mu^2 h)$ resulting in $C\mu^4 h^{-1}(\mu h)^{2-\kappa}\mu^{1-\kappa}$ for $\mu \le h^{-\frac{1}{3}}$ and $C\mu^4 h^{-1}(\mu h)^{2-\kappa}(\mu^2 h)^{\kappa-1}$ for $h^{-\frac{1}{3}} \le \mu \le h^{-\frac{1}{2}}$ and while the former expression is less than $C\mu^{-2+\kappa}h^{-2}$ the latter is not (at their respective intervals).

As $\kappa = 1$ we get $C\mu^5(1 + |\log \mu^2 h|)$.

Consider the second term in the right-hand expression of (9.5). Then both $\alpha \le \beta$ effectively are bounded from above by $(\mu h)^{\frac{1}{2}}$ which leaves interval empty for $\mu \le h^{-\frac{1}{3}}$. On the other hand then $\beta \le \mu^2 h$ for $h^{-\frac{1}{3}} \le \mu \le h^{-1}$ and we arrive to

$$C\mu^2 h^{-2} \iint_{\{\alpha \le \beta \le (\mu h)^{\frac{1}{2}}\}} \alpha\beta^{1-\kappa}\, d\alpha d\beta \asymp C\mu^2 h^{-2}(\mu h)^{2-\frac{1}{2}\kappa}$$

which is less than the right-hand expression of (9.16) unless $h^{-\frac{1}{3}} \le \mu \le h^{-\frac{1}{2}}$ and $0 < \kappa < 1$ in which case it is less than $C\mu^4 h^{-1}(\mu h)^{2-\kappa}(\mu^2 h)^{\kappa-1}$.

Therefore we arrive to

$$|\mathsf{K}^{\mathsf{W}}| \le C\mu^{-2+\kappa}h^{-2} + C\begin{cases} \mu^4 h^{-1}(\mu h)^{2-\kappa} & \text{as } \mu \le h^{-\frac{1}{2}}, \\ \mu^2 h^{-2}(\mu h)^{2-\kappa} & \text{as } h^{-\frac{1}{2}} \le \mu \le h^{-1} \end{cases} + \\ C\begin{cases} \mu^4 h^{-1}(\mu h)^{2-\kappa}(\mu^2 h)^{\kappa-1} & 0 < \kappa < 1, \quad \mu \le h^{-\frac{1}{2}}, \\ \mu^5(1 + |\log \mu^2 h|) & \kappa = 1, \quad \mu \le h^{-\frac{1}{2}}, \\ 0 & \text{otherwise}. \end{cases} \tag{9.18}$$

Thus we proved

Proposition 9.2.3. *For* $\mu \le h^{-1}$ *under condition* (0.7) *estimate* (9.16) *holds and under condition* (0.8) *estimate* (9.18) *holds.*

9.2.3 Magnetic Weyl approximation

Finally, let us deal with magnetic Weyl approximation. Now we should add to what we got for K^{T} corresponding expressions with $R(\alpha,\gamma)=R^{\mathsf{MW}}(\alpha,\gamma)$ calculated according to (1.83) for $\mu \le h^{-1}$ and (1.85) for $\mu \ge h^{-1}$.

(i) Consider case $\mu \le h^{-1}$ first. Then according to (1.83) γ snaps to $\min(\mu^{-1}, \mu h\alpha^{-1})$ and $R(\alpha,\gamma)=\mu h^{-1}$ and (9.3) becomes

$$|\mathsf{K}^{\mathsf{MW}}| \le C\mu^{-2}h^{-2} + C\mu^2h^{-2}\left\{\begin{aligned} &(\mu h)^{2-\kappa} && \mu \le h^{-\frac{1}{2}},\\ &\mu^{\kappa-2} && h^{-\frac{1}{2}} \le \mu \le h^{-1}.\end{aligned}\right. \tag{9.19}$$

Meanwhile in the first term of (9.5) α snaps to its largest possible value i.e 1. Further, in the second term α snaps to its largest value which is β and then β snaps to its largest value $(\mu h)^{\frac{1}{2}}$ as $h^{-\frac{1}{3}} \le \mu \le h^{-1}$ and it becomes $C\mu^2h^{-2}(\mu h)^{2-\frac{1}{2}\kappa}$.

Furthermore, contributions of zones (9.9) and (9.10) are estimated as before; we arrive in the end to

$$\begin{gathered}|\mathsf{K}^{\mathsf{MW}}| \le C\mu^{-2+\kappa}h^{-2} + C\mu^2h^{-2}\left\{\begin{aligned} &(\mu h)^{2-\kappa} && \mu \le h^{-\frac{1}{2}},\\ &\mu^{\kappa-2} && h^{-\frac{1}{2}} \le \mu \le h^{-1}\end{aligned}\right. +\\ C\left\{\begin{aligned} &\mu^2h^{-2}(\mu h)^{2-\frac{1}{2}\kappa} && h^{-\frac{1}{3}} \le \mu \le h^{-1},\\ &0 && \text{otherwise.}\end{aligned}\right.\end{gathered} \tag{9.20}$$

(ii) As $\mu h \ge 1$ γ snaps to $\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ and (9.3) becomes

$$|\mathsf{K}^{\mathsf{MW}}| \le C\mu^{1+\frac{1}{2}\kappa}h^{-1+\frac{1}{2}\kappa} \tag{9.21}$$

Consider the first term in expression (3.5); integrand contains α in the positive power and therefore α snaps to 1. Meanwhile in the second term both $\alpha \ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ and $\beta \ge C_0\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$ are in the negative powers and snap to $\mu^{-\frac{1}{2}}h^{\frac{1}{2}}$. Estimating as before contributions of zones (9.13) and (9.14) we arrive to estimate (9.21) again.

All these above estimates are worse than for K^{T}; thus we arrive to

Proposition 9.2.4. *(i) For $\mu \le h^{-1}$ under condition* (0.7) *estimate* (9.19) *holds and under condition* (0.8) *estimate* (9.20) *holds;*

(ii) For $\mu \ge h^{-1}$ under either condition $(0.7)'$, $(0.8)'$ *estimate* (9.21) *holds.*

9.3 Case $d = 3$

We consider case $d = 3$. Note that

Remark 9.3.1. (i) We use notations in (9.5), (9.6): λ instead of α and λ' instead of β (in the second term only);

(ii) We cannot expect estimate better than $O(h^{-4})$ for $\mu h \le 1$. Meanwhile under condition (0.8) contribution of zone

$$\{(x,y):\ \lambda(x) \le \bar{\lambda},\ \lambda(y) \le \bar{\lambda}\} \tag{9.22}$$

with $\bar{\lambda} = C_0\mu^{-1}$ to any estimate does not exceed $C\mu^2 h^{-4} \times \mu^{-6+\kappa} = O(h^{-4})$. We can even take $\bar{\lambda} = \mu^{-2/(6-\kappa)}$ here.

9.3.1 Tauberian estimates

We will use results of subsection 5.4 combined with arguments of subsubsection 6.8.1 always setting $\gamma_3 = \gamma$.

Case $\mu h \le 1$ As $\gamma_3 = \gamma$ we have $\nu(\boldsymbol{\gamma}) \asymp \lambda\gamma$, $\ell \asymp \lambda$ and (5.25) becomes

$$R^{\mathsf{T}}(\gamma) = \begin{cases} \mu h^{-\frac{3}{2}}\lambda^{\frac{1}{2}} & \text{as } \lambda^2\gamma \le h,\\ \mu h^{-1}\lambda^{-\frac{1}{2}}\gamma^{-\frac{1}{2}} & \text{as } \lambda^2\gamma \ge h, \lambda\gamma \le \mu h,\\ \mu^{\frac{1}{2}} h^{-\frac{3}{2}}(\mu h/\lambda\gamma)^{l} & \text{as } \lambda\gamma \ge \mu h. \end{cases} \tag{9.23}$$

To estimate $|\mathsf{K}^{\mathsf{T}}|$ we need to calculate

$$K(\lambda) := \int \bigl(R^{\mathsf{T}}(\gamma)\bigr)^2 \gamma^{2-\kappa}\, d\gamma \tag{9.24}$$

and as $\lambda \geq \mu^{-1}$ it is equal to $C\mu^2 h^{-2}\lambda^{-1}\gamma^{2-\kappa}$ calculated for $\gamma = \min(\mu h/\lambda, \lambda)$ as $0 < \kappa < 2$ and for $\gamma = h/\lambda^2$ as $2 < \kappa < 3$ resulting in

$$\tag{9.25} K(\lambda) \leq Ch^{-4}\lambda^{3-\kappa} + C\begin{cases} C\mu^2 h^{-2}\lambda^{-1}(h/\lambda^2)^{2-\kappa} & 2 < \kappa < 3, \\ C\mu^2 h^{-2}\lambda^{-1}(1 + |\log \lambda\mu|) & \kappa = 2, \\ \mu^2 h^{-2}\lambda^{-1}\min(\mu h/\lambda, \lambda)^{2-\kappa} & 0 < \kappa < 2. \end{cases}$$

In particular, we conclude that

(9.26) As $\mu h \leq 1$, $|\nabla V/F| \asymp 1$

$$\tag{9.27} |\mathsf{K}^\mathsf{T}| \leq Ch^{-4} + C\begin{cases} \mu^2 h^{-\kappa} & 2 < \kappa < 3, \\ \mu^2 h^{-2}|\log \mu| & \kappa = 2, \\ 0 & 0 < \kappa < 2. \end{cases}$$

On the other hand, under assumption (0.8) we need to calculate the first term in the right-hand expression of (9.5) which is equal to $\int K(\lambda)\lambda^2\, d\lambda$ and one can see easily that it is (9.25) calculated as $\lambda = 1$.

Finally, one needs to calculate the second term in the right-hand expression of (9.5) and we leave to the reader to prove that it allows the same estimate. Thus we arrive to statement (i) of proposition 9.3.2 below.

On the other hand, assume that assumption (6.57) is fulfilled i.e. $|\nabla_{\perp \mathsf{F}} V/F| \asymp 1$. Then (5.25) becomes

$$\tag{9.28} R^\mathsf{T}(\gamma) = \begin{cases} \mu^{\frac{3}{2}} h^{-1}\bigl(1 + |\log(\mu h)|\bigr) & \text{as} \quad \gamma \leq h, \\ \mu^{\frac{3}{2}} h^{-1}\bigl(1 + (\log(1/\mu\gamma))_+\bigr) & \text{as} \quad h \leq \gamma \leq \min(\mu h, \mu^{-1}), \\ \mu h^{-1}\gamma^{-\frac{1}{2}} & \text{as} \quad \mu^{-1} \leq \gamma \leq \mu h, \\ \mu^{\frac{1}{2}} h^{-\frac{3}{2}}(\mu h/\gamma))^l & \text{as} \quad \gamma \geq \mu h. \end{cases}$$

and in this case instead of $\gamma = h$ the lower pick is $\gamma = \mu^{-1}$ in $C\mu^2 h^{-2}\gamma^{2-\kappa}$ which becomes $Ch^{-2}\mu^\kappa$ and we arrive to statement (ii) of proposition 9.3.2:

Proposition 9.3.2. *Let $\mu h \le 1$. Then*

(i) Under either non-degeneracy condition either (0.7) *or* (0.8) *estimate* (9.27) *holds.*

In particular, $|\mathsf{K}^\mathsf{T}| = O(h^{-4})$ as *either* $0 < \kappa < 2$ *or* $\kappa = 2$ *and* $\mu \le (h|\log h|)^{-1}$ *or* $2 < \kappa < 3$ *and* $\mu \le h^{-2+\frac{1}{2}\kappa}$ *(and this is true for sure as* $\mu \le h^{-\frac{1}{2}}$*).*

(ii) Under non-degeneracy condition (6.57)*,* $2 < \kappa < 3$

$$|\mathsf{K}^\mathsf{T}| \le Ch^{-4} + Ch^{-2}\mu^\kappa. \tag{9.29}$$

In particular, $|\mathsf{K}^\mathsf{T}| = O(h^{-4})$ *as* $\mu \le h^{-2/\kappa}$ *(and this is true for sure as* $\mu \le h^{-\frac{2}{3}}$*).*

Case $\mu h \ge 1$

Proposition 9.3.3. *Let $\mu h \ge 1$. Then under either non-degeneracy condition* $(0.7)'$ *or* $(0.8)'$ *estimate*

$$|\mathsf{K}^\mathsf{T}| \le C\mu^2 h^{-2} + C\begin{cases} \mu^2 h^{-\kappa} & 2<\kappa<3,\\ \mu^2 h^{-2}|\log\mu| & \kappa = 2,\\ 0 & 0<\kappa<2. \end{cases} \tag{9.30}$$

holds.

Proof. An easy proof based on proposition 5.5.3 is left to the reader. □

9.3.2 Weyl asymptotics

Proposition 9.3.4. *Let $\mu h \le 1$. Then*

(i) Under non-degeneracy condition (0.7)

$$|\mathsf{K}^\mathsf{W}| \le Ch^{-4} + C\mu^3 h^{-3}(\mu h)^{3-\kappa}; \tag{9.31}$$

(ii) Under non-degeneracy condition (0.8)

$$|\mathsf{K}^\mathsf{W}| \le Ch^{-4} + C\mu^3 h^{-3}(\mu h)^{\frac{1}{2}(3-\kappa)}; \tag{9.32}$$

(iii) As $\mu \le h^{-\frac{1}{2}}$ under non-degeneracy condition (6.57)

$$|\mathsf{K}^\mathsf{W}| \le Ch^{-4} + C\mu^5 h^{-2}(\mu h)^{3-\kappa}. \tag{9.33}$$

Proof. An easy proof based on proposition 5.4.3 is left to the reader. □

9.3.3 Magnetic Weyl asymptotics

We will apply proposition 5.6.4. In the isotropic settings $R^{\mathsf{MW}}(\gamma)$ given by (5.47) becomes

$$R^{\mathsf{MW}}(\gamma) := \\ C\mu\begin{cases} \lambda^{-\frac{1}{2}}\gamma^{-\frac{5}{2}}\bigl(\mu h/\lambda\gamma\bigr)^{l} & \text{as } \gamma \ge \mu h/\lambda,\\ \lambda^{-\frac{1}{2}}\gamma^{-\frac{5}{2}} & \text{as } \lambda^{-\frac{1}{3}}h^{\frac{2}{3}} \le \gamma \le \mu h/\lambda,\\ \lambda^{\frac{1}{3}}h^{-\frac{5}{3}} & \text{as } \gamma \le \lambda^{-\frac{1}{3}}h^{\frac{2}{3}}, \end{cases} \tag{9.34}$$

which immediately implies statements (i), (ii) of proposition 9.3.5 below.

Meanwhile as $\alpha \asymp 1$, $\mu h \le 1$ (5.47) becomes

$$R_1^{\mathsf{MW}}(\gamma) := \\ C\mu\begin{cases} \gamma^{-\frac{5}{2}}(\mu h/\gamma)^{l} & \text{as } \gamma \ge \mu h,\ 1 \le \mu^2 h,\\ \gamma^{-\frac{5}{2}} & \text{as } \gamma \le \mu h,\ \mu\gamma \ge 1,\\ \gamma^{-\frac{5}{2}}(\mu^2 h)^{\frac{1}{2}}\bigl(\mu h/\gamma\bigr)^{l} & \text{as } \gamma \ge \mu h,\ 1 \ge \mu^2 h,\\ \gamma^{-\frac{5}{2}}(\mu\gamma)^{\frac{1}{2}} & \text{as } \gamma \le \mu h,\ \mu\gamma \le 1 \end{cases} \tag{9.35}$$

which immediately imply statement (iii) of proposition 9.3.5 below.

Proposition 9.3.5. *(i) Let $\mu h \le 1$. Then under either non-degeneracy condition* (0.7) *or* (0.8)

$$|\mathsf{K}^{\mathsf{MW}}| \le Ch^{-4} + C\mu^2 h^{-\frac{2}{3}(2+\kappa)}; \tag{9.36}$$

(ii) Let $\mu h \ge 1$. Then under either non-degeneracy condition(0.7)′ *or* (0.8)′

$$|\mathsf{K}^{\mathsf{MW}}| \le C\mu^2 h^{-2} + C\mu^2 h^{-\frac{2}{3}(2+\kappa)}; \tag{9.37}$$

(iii) Let $\mu \le h^{-\frac{2}{3}}$. Then under non-degeneracy condition (6.57)

$$|\mathsf{K}^{\mathsf{MW}}| \le Ch^{-4} + C\mu^3 h^{-\frac{2}{3}(1+\kappa)}. \tag{9.38}$$

9.4 Problems

Problem 9.4.1. For $d = 2, 3$ estimate $|\mathsf{J} - \mathsf{J}^{\mathsf{T}}|$, $|\mathsf{J} - \mathscr{J}^{\mathsf{W}}|$ and $|\mathsf{J} - \mathscr{J}^{\mathsf{MW}}|$.

Problem 9.4.2. *For $d = 2, 3$ consider pilot-model approximation.*

Problem 9.4.3. For $d = 2, 3$ compare estimates for $|\mathsf{K}^{\mathsf{W}}|$ and $|\mathsf{J} - \mathscr{J}^{\mathsf{W}}|$ to those for $|\mathsf{K}^{\mathsf{MW}}|$ and $|\mathsf{J} - \mathscr{J}^{\mathsf{MW}}|$.

Problem 9.4.4. *For $d = 2, 3$ as $\mu \le h^{-1}$ get rid off condition (0.5) replacing it by (0.7), or even $(0.8)^+$, or even (0.8).*

Problem 9.4.5. *Get rid off condition (0.6) assuming instead that $|F| + |\nabla F| \ge \epsilon$ for $d = 2$ (which will affect estimates for sure) and $|\mathbf{F}| + |\nabla \otimes \mathbf{F}| \ge \epsilon$ for $d = 3$ (which most likely would not affect estimates)*

Problem 9.4.6. *For $d = 3$ investigate K^{T}, K^{W}, K^{MW} under non-degeneracy conditions (0.7) and $|\nabla_{\perp \mathbf{F}} V/F| \le \epsilon \implies |\operatorname{det}\operatorname{Hess}_{\perp \mathbf{F}} V/F| \asymp 1$.*

Bibliography

[Ivr1] V. Ivrii. *Microlocal Analysis and Precise Spectral Asymptotics*, Springer-Verlag, SMM, 1998, xv+731.

[Ivr2] V. Ivrii. *Microlocal Analysis and Sharp Spectral Asymptotics*, in progress: available online at http://www.math.toronto.edu/ivrii/futurebook.pdf

[Ivr3] Sharp spectral asymptotics for Dirac energy, arXiv:0708.3934 (July 9, 2007), 25pp.

[IRO4] Sharp spectral asymptotics for operators with irregular coefficients. IV. Multidimensional Schrödinger operator with a strong magnetic Field. Full-rank case, arXiv:math/0510327 (Aug. 2, 2004), 83pp.

[MO] ter Morsche, H. and Oonincx, P. J. On the integral representations for metaplectic operators. J. Fourier Analysis and Appls, 8(3):245–258 (2002).

Department of Mathematics,
University of Toronto,
40, St.George Str.,
Toronto, Ontario M5S 2E4
Canada
ivrii@math.toronto.edu
Fax: (416)978-4107